

lncRNAs: Linking RNA to Chromatin

John L. Rinn

Harvard University, Department of Stem Cell and Regenerative Biology, Cambridge, Massachusetts 02138

Correspondence: john_rinn@harvard.edu

Numerous studies over the past decade have identified increasing numbers of long noncoding RNAs (lncRNAs) across many organisms. Research since has shown that lncRNAs constitute an important layer of genome regulation in diverse biological processes and disease. Here, we discuss the common emerging theme of lncRNAs interfacing with epigenetic machinery. This, in turn, modulates the activity and localization of the epigenetic machinery during cell fate specification.

Cellular identity is achieved through a complex choreography of DNA regulatory elements interacting with protein regulatory complexes to shape a myriad of unique epigenetic landscapes. The vast arrays of epigenetic landscapes are established by using ubiquitously expressed chromatin-modifying and -remodeling complexes, to give rise to countless unique combinations of histone posttranslational modifications and DNA methylation patterns. This raises the age-old question: How are the enzymatic complexes being guided to place these marks at a specific combination of sites under different cellular contexts? It has long been suspected that noncoding RNA molecules may provide some specificity to target these complexes to their sites of action. Indeed, it is becoming increasingly clear that a contingent of thousands of long noncoding RNAs (lncRNAs) represent a key layer of epigenetic control (see Fig. 24 of Allis et al. 2014).

A dramatic example of RNA-based epigenetic regulation that operates in mammalian dosage compensation has been known for more than 20 years (Brockdorff and Turner 2014). Specifically, a long intergenic noncoding RNA (lincRNA) termed XIST (X inactive-specific transcript) is expressed from one female X chromosome resulting in the recruitment of Polycomb group complexes (PcG), such as PRC2, to this chromosome with the concomitant

transcriptional silencing across a majority of the X chromosome. In other words, a single lncRNA gene is able to target and silence the majority of a chromosome *in cis*. This is a powerful precedent for RNA-mediated epigenetic regulation. However, the link between the RNA and PRC2 recruitment was, for a long time, elusive. A clue came from studying the epigenetic dynamics of a different lincRNA termed HOTAIR (Hox transcript antisense intergenic RNA; Rinn et al. 2007).

HOTAIR is expressed from one of the four clusters of HOX transcription factors (i.e., HOXC). The HOX family of proteins is a key regulator of body plan during development. HOTAIR was identified because of its distinctive pattern of expression in posterior and distal mouse tissues during development and adult human fibroblasts (Rinn et al. 2007). Intriguingly, HOTAIR expression demarcated a distinct epigenetic boundary between euchromatic and heterochromatic regions within the HOXC cluster. Moreover, the euchromatic-heterochromatic regions were inverted between anterior and posterior cell types; we termed these “diametric” chromatin domains. Collectively, the data led to the initial hypothesis that HOTAIR serves as an epigenetic boundary *in cis* within the HOXC cluster. To our surprise the HOXC chromatin boundary remained unchanged when HOTAIR RNA function was lost; however, the HOXD cluster, located on a separate chromosome, became active. Thus, similar to XIST, the expression of HOTAIR from the HOXC cluster results in the epigenetic silencing, yet the HOX cluster that it regulates (HOXD) is located on a different chromosome, i.e., is regulated *in trans*. This raised the question: How?

The answer was determined by a few critical experiments, which indicated that the HOTAIR lincRNA interacted with PRC2 and that this was required for proper



localization of PRC2. The first experiment, using immunoprecipitation of the PRC2 complex, identified HOTAIR as being physically associated or coprecipitated with PRC2 (Rinn et al. 2007). The reciprocal experiment, which isolated proteins bound to the HOTAIR transcript, also revealed an association between HOTAIR and components of the PRC2 complex but not other chromatin regulatory factors. The final missing link was the demonstration that an RNA–protein interaction between HOTAIR and PRC2 is required for proper localization of PRC2 across the HOXD cluster. Indeed, depletion of HOTAIR resulted in the mislocalization of PRC2 from the HOXD cluster, *in trans*, and concomitant activation of HOXD genes. Together, these findings showed that the physical association between HOTAIR and PRC2 was required for the proper targeting of the chromatin regulatory machinery to their target sites. Thus, a novel layer of RNA-based modulation of epigenetic landscapes was unequivocally revealed.

Since unraveling the fact that PRC2 physically associates with HOTAIR or XIST, so targeting the complex to specific genomic regions, this mechanism has been shown to apply to numerous lncRNAs that physically associate with PRC2.

Many of these associations are, in fact, required for the proper localization of the epigenetic regulatory machinery *in cis* and *in trans* (see Fig. 1). Two independent studies, in particular, determined that in both human and mouse cells, hundreds of lncRNAs, accounting for up to 30% of the transcriptome, coprecipitate with PRC2. Moreover, many of the tested PRC2-bound lncRNAs were required for proper epigenetic and transcriptional regulation of PRC2 targets (Khalil et al. 2009; Zhao et al. 2010). It was further noted from these studies that one RNA could be bound to many different chromatin regulatory proteins, suggesting that it could function as an RNA bridge across multiple complexes (Khalil et al. 2009). Indeed, a detailed biochemical analysis of HOTAIR showed that in addition to binding PRC2, it also bound the histone demethylase LSD1 and NCOR. This pointed to a novel model for epigenetic regulation whereby a single lncRNA recruits several synergistic chromatin regulatory complexes to help guide, dock (NCOR), and facilitate heterochromatin formation (LSD1 and PRC2). Collectively, these studies led to the idea that an RNA scaffold can bridge numerous chromatin and additional regulatory complexes to impart genome target specificity (Fig.




Figure 1. Models of how lncRNAs may function in the epigenetic control of gene expression, both activating and repressing transcription *in cis* and *in trans*. (A) Enhancer RNAs have been shown to play an RNA-mediated role in forming enhancer interactions resulting in long-range *cis* gene regulation by lncRNAs. (B) Activation by the JPX lncRNA as an example of a *trans*-acting lncRNA, which in this case facilitates Xist activation. (C) Xist as an example of an lncRNA that facilitates gene repression *in cis* across the majority of the X chromosome. (D) HOTAIR expression results *in trans* repression of HOX genes.

1D). The recent finding that HOTAIR overexpression is a hallmark in metastatic breast cancer (Gupta et al. 2010) has underscored the importance of HOTAIRs role in epigenetic regulation. In fact, HOTAIR serves as an “onco-lncRNA,” inducing metastasis in breast cancer when overexpressed by remodeling the epithelial epigenome to resemble that of stromal cells. Cumulatively, the lessons learned from HOTAIR over the past five years have shown that lncRNAs play a critical role in interfacing with and modulating chromatin complexes during development and disease.

In the 50 years since RNA was identified as a central component in the flow of genetic information, it has become increasingly clear that RNA is more than a mere messenger and instead performs vast and diverse functions (Amaral et al. 2008). lncRNAs are, in fact, emerging as a critical layer of epigenetic regulation in which different lncRNAs are associated with distinctive epigenetic states, yet share a common mechanism; they physically associate with chromatin-modifying and -remodeling complexes, and guide them to specific genomic loci that are crucial for proper cellular function. However, this is only one facet of lncRNA biology; there is a diversity of other functional roles they play across numerous biological processes alluded to in Amaral et al. (2008).

REFERENCES

*Reference is also in this collection.

- * Allis CD, Jenuwein T, Reinberg D. 2014. Overview and concepts. *Cold Spring Harb Perspect Biol* doi: 10.1101/cshperspect.a018739.
- Amaral PP, Dinger ME, Mercer TR, Mattick JS. 2008. The eukaryotic genome as an RNA machine. *Science* **319**: 1787–1789.
- * Brockdorff N, Turner BM. 2014. Dosage compensation in mammals. *Cold Spring Harb Perspect Biol* doi: 10.1101/cshperspect.a019406.
- Gupta RA, Shah N, Wang KC, Kim J, Horlings HM, Wong DJ, Tsai MC, Hung T, Argani P, Rinn JL, et al. 2010. Long non-coding RNA *HOTAIR* reprograms chromatin state to promote cancer metastasis. *Nature* **464**: 1071–1076.
- Khalil AM, Guttman M, Huarte M, Garber M, Raj A, Rivea Morales D, Thomas K, Presser A, Bernstein BE, van Oudenaarden A, et al. 2009. Many human large intergenic noncoding RNAs associate with chromatin-modifying complexes and affect gene expression. *Proc Natl Acad Sci* **106**: 11667–11672.
- Rinn JL, Kertesz M, Wang JK, Squazzo SL, Xu X, Bruggmann SA, Good-nough LH, Helms JA, Farnham PJ, Segal E, et al. 2007. Functional demarcation of active and silent chromatin domains in human *HOX* loci by noncoding RNAs. *Cell* **129**: 1311–1323.
- Tsai MC, Manor O, Wan Y, Mosammamaparast N, Wang JK, Lan F, Shi Y, Segal E, Chang HY. 2010. Long noncoding RNA as modular scaffold of histone modification complexes. *Science* **329**: 689–693.
- Zhao J, Ohsumi TK, Kung JT, Ogawa Y, Grau DJ, Sarma K, Song JJ, Kingston RE, Borowsky M, Lee JT. 2010. Genome-wide identification of polycomb-associated RNAs by RIP-seq. *Mol Cell* **40**: 939–953.



IncRNAs: Linking RNA to Chromatin

John L. Rinn

Cold Spring Harb Perspect Biol 2014; doi: 10.1101/cshperspect.a018614

Subject Collection [Epigenetics](#)

Metabolic Signaling to Chromatin

Shelley L. Berger and Paolo Sassone-Corsi

Histone and DNA Modifications as Regulators of Neuronal Development and Function

Stavros Lomvardas and Tom Maniatis

Histone Modifications and Cancer

James E. Audia and Robert M. Campbell

Epigenetics and Human Disease

Huda Y. Zoghbi and Arthur L. Beaudet

Induced Pluripotency and Epigenetic Reprogramming

Konrad Hochedlinger and Rudolf Jaenisch

Long-Range Chromatin Interactions

Job Dekker and Tom Misteli

RNAi and Heterochromatin Assembly

Robert Martienssen and Danesh Moazed

Dosage Compensation in *Drosophila*

John C. Lucchesi and Mitzi I. Kuroda

Epigenetic Determinants of Cancer

Stephen B. Baylin and Peter A. Jones

Maintenance of Epigenetic Information

Geneviève Almouzni and Howard Cedar

A Structural Perspective on Readout of Epigenetic Histone and DNA Methylation Marks

Dinshaw J. Patel

The Necessity of Chromatin: A View in Perspective

Vincenzo Pirrotta

Germline and Pluripotent Stem Cells

Wolf Reik and M. Azim Surani

Comprehensive Catalog of Currently Documented Histone Modifications

Yingming Zhao and Benjamin A. Garcia

Epigenetic Regulation of Chromatin States in *Schizosaccharomyces pombe*

Robin C. Allshire and Karl Ekwall

Histone Variants and Epigenetics

Steven Henikoff and M. Mitchell Smith

For additional articles in this collection, see <http://cshperspectives.cshlp.org/cgi/collection/>



All Modifications and
Oligo Types Synthesized

Long Oligos • Fluorescent • Chimeric • DNA • RNA • Antisense

Oligo Modifications?

Your wish is our command.



For additional articles in this collection, see <http://cshperspectives.cshlp.org/cgi/collection/>



**All Modifications and
Oligo Types Synthesized**

Oligo Modifications?
Your wish is our command.

Long Oligos • Fluorescent • Chimeric • DNA • RNA • Antisense

