

Molecular architecture of a miRNA-regulated 3' UTR

DOMINIC DIDIANO and OLIVER HOBERT

Department of Biochemistry and Molecular Biophysics, Howard Hughes Medical Institute, Columbia University Medical Center, New York, New York 10032, USA

ABSTRACT

Animal genomes contain hundreds of microRNAs (miRNAs), small regulatory RNAs that control gene expression by binding to complementary sites in target mRNAs. Some rules that govern miRNA/target interaction have been elucidated but their general applicability awaits further experimentation on a case-by-case basis. We use here an assay system in transgenic nematodes to analyze the interaction of the *Caenorhabditis elegans* *lisy-6* miRNA with 3' UTR sequences. In contrast to many previously described assay systems used to analyze miRNA/target interactions, our assay system operates within the cellular context in which *lisy-6* normally functions, a single neuron in the nervous system of *C. elegans*. Through extensive mutational analysis, we define features in the known and experimentally validated target of *lisy-6*, the 3' UTR of the *cog-1* homeobox gene, that are required for a functional miRNA/target interaction. We describe that both in the context of the *cog-1* 3' UTR and in the context of heterologous 3' UTRs, one or more seed matches are not a reliable predictor for a functional miRNA/target interaction. We rather find that two nonsequence specific contextual features beyond miRNA target sites are critical determinants of miRNA-mediated 3' UTR regulation. The contextual features reside 3' of *lisy-6* binding sites in the 3' UTR and act in a combinatorial manner; mutation of each results in limited defects in 3' UTR regulation, but a combinatorial deletion results in complete loss of 3' UTR regulation. Together with two *lisy-6* sites, these two contextual features are capable of imparting regulation on a heterologous 3' UTR. Moreover, the contextual features need to be present in a specific configuration relative to miRNA binding sites and could either represent protein binding sites or provide an appropriate structural context. We conclude that a given target site resides in a 3' UTR context that evolved beyond target site complementarity to support regulation by a specific miRNA. The large number of 3' UTRs that we analyzed in this study will also be useful to computational biologists in designing the next generation of miRNA/target prediction algorithms.

Keywords: *C. elegans*; miRNA; *lisy-6*; *cog-1*; 3' UTR

INTRODUCTION

Animal miRNAs are a large class of small regulatory RNAs that control the expression of target genes through binding to complementary target sites in mRNA targets (Lee et al. 1993; Reinhart et al. 2000; Bartel 2004). These sites typically have imperfect complementarity to the miRNA. The ability to target imperfectly complementary target sites together with the large number of miRNAs in genomes (as many as 1000 in the human genome) (Bentwich et al. 2005) has led to the prediction that thousands of genes are miRNA-regulated (Rajewsky 2006b). However, despite the apparent abundance of potential miRNA/3' UTR target interactions, few animal miRNA/target pairs have been validated to date

in a physiologically relevant context (Carthew 2006; Stefani and Slack 2008).

The parameters that govern miRNA/3' UTR target recognition have grown increasingly complex as more rigorous sequence analyses and experimentation has examined miRNA/target interactions. Initial reports indicated that G:U wobble-free seed matches between position 2-8 of a miRNA and the target site and the ΔG of the miRNA/target heteroduplex were the primary determinants in miRNA target recognition (Doench et al. 2003; Doench and Sharp 2004; Brennecke et al. 2005; Lewis et al. 2005; Grimson et al. 2007). Additional parameters such as the importance of 3' base-pairing to compensate for weak seed pairing became another factor in identifying potential miRNA targets (Brennecke et al. 2005). However, some other work also gave indications that miRNA/target recognition may be more complex. For example, *lin-4* and *let-7* binding to some of their targets involves both mismatches within the seed, as well as G:U base pairs (Ha et al. 1996; Vella et al. 2004a). Moreover, it was reported that *let-7* requires two unique

Reprint requests to: Oliver Hobert, Department of Biochemistry and Molecular Biophysics, Howard Hughes Medical Institute, Columbia University Medical Center, New York, NY 10032, USA; e-mail: or38@columbia.edu; fax: (212) 342-1810.

Article published online ahead of print. Article and publication date are at <http://www.rnajournal.org/cgi/doi/10.1261/rna.1082708>.

target sites and that the linker region between them in the *lin-41* 3' UTR is required to mediate regulation (Vella et al. 2004a), thereby suggesting that miRNA target sites may require specific 3' UTR contexts to function.

Robins and Padgett (2005) were the first to explicitly explore the role that mRNA structure may play in influencing miRNA/target recognition. They postulated that the folding of mRNAs may mask some potential miRNA target sites and devised a target prediction algorithm that required that at least three consecutive base pairs in the target site complementary to the seed region of the miRNA (nucleotides 2–8) be unpaired in the mRNA structure. Since this initial report, several groups have further advanced miRNA target prediction by including structural considerations in the form of the difference in free energy between the miRNA bound and unbound states ($\Delta\Delta G$) as a means to identify potential target 3' UTRs (Kertesz et al. 2007; Long et al. 2007). Moreover, Zhao et al. (2005, 2007) reported that target sites reside in 3' UTR regions of high free energy based on the ΔG of the 70-base-pair (bp) 5' and 3' of a target site. Another study has added a number of additional considerations that boost site efficacy when attempting to identify potential target 3' UTRs, including target site location relative to the stop codon, AU-richness around target sites, and preferential base pairing to nucleotides 13–16 of the miRNA (Grimson et al. 2007).

Numerous algorithms now exist that attempt to identify potential miRNA target 3' UTRs, including PicTar, TargetScan, Miranda, and Mirbase (Stark et al. 2003; John et al. 2004; Krek et al. 2005; Lewis et al. 2005; Robins and Padgett 2005; Griffiths-Jones et al. 2006; Lall et al. 2006). These algorithms vary in their use of conservation, energetics, and structural considerations. The overlap between target sites predicted by different algorithms is, however, still limited (Rajewsky 2006b). The difficulty in target predictions is further underscored by the observation that only a limited number of mRNAs bound to the miRNP/Ago/AIN complexes are predicted to be targeted by known miRNAs (Beitzinger et al. 2007; Zhang et al. 2007). Part of the difficulty in formulating accurate predictions is the lack of data examining the ability of endogenous miRNAs to down-regulate target 3' UTRs in the cellular context in which such an interaction normally takes place. Another problem is that rules of miRNA/mRNA interactions were built on assay systems that rely on overexpression of the miRNA and its target. Doench and Sharp (2004) reported that endogenously expressed *let-7* was incapable of down-regulating some of their 3' UTR reporter constructs but that some of these same constructs were down-regulated by exogenously supplied *let-7*. Therefore, these authors warned that overexpression of a miRNA in testing for potential target 3' UTR regulation may suggest an interaction that may not occur in vivo.

We have developed an assay system that circumvents at least some of these problems (Didiano and Hobert 2006).

In this assay system, we analyze the ability of the *Caenorhabditis elegans* miRNA *lisy-6* to regulate a given 3' UTR. *lisy-6* is a miRNA that is expressed in a very small number of neurons, and in one of them, the left ASE (ASEL) neuron, it down-regulates expression of the *cog-1* homeobox gene in a 3' UTR dependent manner (Johnston and Hobert 2003). The assay is based on the use of the *ceh-36* promoter to drive the expression of *gfp* in a pair of bilaterally symmetric neurons, the left and right ASE neurons (ASEL and ASER) (Chang et al. 2003). Control and experimental 3' UTRs are placed downstream of the *gfp* coding sequence (Fig. 1A). As *lisy-6* is asymmetrically expressed in ASEL, but not in ASER, this constellation provides an assay for evaluating the ability of *lisy-6* to down-regulate 3' UTR sensor constructs in ASEL, but not ASER (Fig. 1B). Notably, this assay system tests the interaction of a miRNA (*lisy-6*) with its target (*cog-1*) in exactly the same cell in which the interaction normally takes place. Other possible cell-type specific factors involved in the *lisy-6*-mediated regulation of the 3' UTR should therefore be present. Many of the assay systems that have been used in the past to deduce rules of miRNA/target




FIGURE 1. 3' UTR sensor strategy. (A) Schematic representation of 3' UTR sensor constructs. (B) Example of expression patterns of a regulated (*ceh-36::gfp-cog-1* 3' UTR) (top) and an unregulated (*ceh-36::gfp-unc-54* 3' UTR) (bottom) 3' UTR sensor constructs. These images are the same as shown in Didiano and Hobert (2006) and are shown here again merely as an illustration of our assay system. (Reprinted with permission from Didiano and Hobert 2006; © 1998, Nature Publishing Group [http://www.nature.com/].) *lisy-6* is only expressed in ASEL as indicated by arrows. *gfp* expression is dramatically down-regulated in ASEL but not ASER in the *cog-1* 3' UTR sensor construct, but the *unc-54* 3' UTR sensor construct displays equal *gfp* expression in ASEL and ASER. The pair of bright cells above the two ASE cells are the AWCL and AWCR neurons. They serve as internal controls for excluding the scoring of mosaic animals. The degree of regulation can also be quantified using an “asymmetric index” as defined in the Material and Methods and as listed in Supplemental Table 1.

interactions do not display this feature. The second advantage of our assay system is that it performs a direct comparison of *gfp* expression levels in ASEL with ASER in each individual animal (Fig. 1B; see Material and Methods for more details), thereby providing an internal control that is not provided by many of the population-based transfection systems used to deduce features of miRNA/target interactions. Finally, unlike transfection systems, our system relies on the activity of an endogenous miRNA, rather than relying on an overexpression of the miRNA under investigation. As mentioned above, this is an important point as Doench and Sharp (2004) have previously shown that expression levels have an important impact on the ability of a miRNA to regulate a target 3' UTR. We note, however, that our assay does rely on the ectopic expression of a sensor construct, thereby raising the issue that a sensor construct may be expressed at too high a level to allow for regulation by *lisy-6*. We deal with this issue by generating transgenic worms that contain only low copy numbers of the reporter gene, as assessed by *gfp* expression levels that are just above the level of detection (Fig. 1B). These levels of reporter gene expression easily allow for regulation of the wild-type *cog-1* 3' UTR sensor by *lisy-6*, the control to which we compare our mutated 3' UTR versions. Moreover, we always generate multiple independent lines for each construct. If we were indeed operating at levels that are close to a saturated threshold, we would expect to observe some variations between individual lines, which we essentially never observe. We are therefore confident that unregulated sensor constructs in our assay system represent true negatives rather than artifacts of excessive transgene expression.

We have previously used this assay to test the ability of *lisy-6* to regulate mutated versions of the *cog-1* 3' UTR as well as heterologous 3' UTRs that were predicted to be targeted by *lisy-6* (Didiano and Hobert 2006). The key findings of this previous analysis were that even though a seed match to the *lisy-6* miRNA is required for *cog-1* 3' UTR regulation, a seed match is not sufficient to confer regulation. This conclusion was based on (1) an engineered mutation within the *cog-1* 3' UTR that did *not* affect the core *lisy-6* seed match, but led to a loss of regulation, (2) the analysis of heterologous 3' UTRs into which a *lisy-6* site was engineered, and (3) the finding that the 3' UTRs of 13 different genes that contained seed matches to *lisy-6* failed to display *lisy-6*-dependent regulation (Didiano and Hobert 2006). Two independently developed miRNA/target prediction methods accurately predicted the inability of *lisy-6* to regulate these 13 genes, therefore lending further credibility to our experimental findings (Long et al. 2007; Zhao et al. 2007). Nevertheless, it was suggested that the 3' UTR sensors that we tested were expressed at levels too high to allow for regulation by endogenous *lisy-6* (Rajewsky 2006a; Bushati and Cohen 2007). However, it appears improbable to us that each of the 13 predicted targets are

endogenously expressed at low enough levels to allow for regulation by our *lisy-6*, and that in our assay system we overexpressed each single one of these 13 different 3' UTRs, thereby preventing regulation by endogenous *lisy-6*. This also appears unlikely in light of the fact that the *cog-1* gene itself is expressed at low levels in the ASE neurons, a notion derived from reporter gene assays (Palmer et al. 2002). We also tested expression levels of one of the 13 predicted target genes and found it to be not significantly different from that of *cog-1* (J. Etchberger and O. Hobert, unpubl.), thereby making the level argument the least parsimonious explanation of our data.

Rajewsky (2006a) and Bushati and Cohen (2007) also raised the possibility that a second *lisy-6* target site in the *cog-1* 3' UTR, which we had previously reported (Johnston and Hobert 2003) (called site #2 throughout this present paper), may be the explanation for why we do not find a single *lisy-6* site to be sufficient to confer regulation. We find this an unlikely explanation since we reported that two of the 13 predicted *lisy-6* target 3' UTRs contain two predicted *lisy-6* sites but are nevertheless not regulated by *lisy-6* (Didiano and Hobert 2006). Moreover, we had shown that a single *lisy-6* site does function in the context of the *lin-28* 3' UTR. One of the goals of the present paper is to further address the importance of this second site. The findings that we describe here indicate that this second site cannot explain the lack of sufficiency of seed matches to confer miRNA-mediated 3' UTR regulation.

Besides testing the importance of both *lisy-6* complementary sites in the *cog-1* 3' UTR, we describe an extensive mutational analysis of the *cog-1* 3' UTR and describe the analysis of a number of engineered, heterologous 3' UTRs. The findings presented here support the notion that seed matches are not a sufficient predictor for miRNA/target interactions. We show that the sequence context of a 3' UTR is a critical determinant of target site functionality and propose that entire 3' UTRs rather than just individual sites evolve to support regulation by a miRNA. We also show that available prediction algorithms fail to provide an accurate assessment of whether a 3' UTR is regulated by *lisy-6*. Our large number of experimentally tested 3' UTRs provide a data set to improve these algorithms.

RESULTS

Assessing the importance of a *lisy-6* complementary site in the *cog-1* 3' UTR

Our previous analysis of the *cog-1* 3' UTR has demonstrated an absolute requirement of a *lisy-6* complementary site (henceforth called "*lisy-6* site #1") for the correct and differential regulation of the *cog-1* 3' UTR in ASEL versus ASER. Deletion of site #1 causes a complete loss of 3' UTR regulation (Didiano and Hobert 2006). We extended these




FIGURE 2. (Legend on next page)

findings by a systematic point mutational analysis of site #1 (Fig. 2). We find that point mutations in the seed region differentially disrupt 3' UTR regulation. Point mutations only in seed positions 4, 5, and 6, but not in positions 1–3 or 7–11, disrupt 3' UTR function significantly. These findings suggest that sequences outside the seed may contribute to *lsey-6* function. Indeed, a deletion reducing site #1 from an 11mer seed match to an 8mer seed match results, as we have previously reported, in the complete loss of regulation (shown again in Fig. 4, no. 3, below [note: when “no.” is mentioned, this refers to the numbered black circles in a figure]; Didiano and Hobert 2006). The importance of site #1 can also further be illustrated by its conversion into a site that is perfectly complementary to *lsey-6*. Such a perfect site is entirely unable to confer regulation (Fig. 2, no. 14). However, regulation can be re-established through reintroduction of a single mismatch (Fig. 2, no. 15). Taken together, various permutations of *lsey-6* site #1 illustrate the importance of the overall composition of this site.

Insufficiency of the *lsey-6* complementary site #1

We have previously shown that *lsey-6* site #1 from the *cog-1* 3' UTR can confer regulation when transplanted into the context of the *lin-28* 3' UTR but not of the *unc-54* 3' UTR (Didiano and Hobert 2006) (shown again, for comparison, in Fig. 3, nos. 2,4). We extended this analysis and find that *lsey-6* site #1 fails to confer regulation to the 3' UTR of the *actin* gene *act-1* (Fig. 3, no. 10), which is, in contrast to the *unc-54* 3' UTR (885 bp), a small 3' UTR (194 bp) that is also low in GC content (35%).

The *lin-28* 3' UTR is known to be normally regulated by a miRNA, *lin-4* (Moss et al. 1997), and also contains additional *let-7* target sites (Krek et al. 2005), while neither the *actin* nor the *unc-54* 3' UTR are predicted to be miRNA regulated. We therefore hypothesized that there may be some feature(s) common only to miRNA-regulated 3' UTRs that enable them to support miRNA regulation. To test this notion, we engineered *lsey-6* #1 site into the 3' UTR

of the *lin-41* and *lin-14* genes, both known miRNA targets (Wightman et al. 1993; Slack et al. 2000). The *lsey-6* site was engineered into an AU-rich sequence of the *lin-41* 3' UTR and into a *let-7* site of the *lin-14* 3' UTR. Neither construct is subject to *lsey-6* regulation (Fig. 3, nos. 5–8).

Additionally, we find that the *lsey-6* site fails to confer regulation when placed in another position in the *lin-28* 3' UTR, in which it replaces a *lin-4* target site (Fig. 3, no. 3). The *lsey-6* site therefore works in one, but not in another, context of the *lin-28* 3' UTR (Fig. 3, nos. 2,3). It therefore appears unlikely that the *lsey-6* site works in *lin-28* because *lin-28* contains some cryptic features (e.g., cryptic *lsey-6* sites) that permit it to work. If this were the case, the *lsey-6* site should have worked in both contexts in the *lin-28* 3' UTR, which it does not.

The *cog-1* 3' UTR contains two target sites for *lsey-6* but these sites are not sufficient in all 3' UTR contexts

Our previous analysis of the *cog-1/lsey-6* interaction focused on only one *lsey-6* target site, “site #1,” which contains an 11mer seed region (Didiano and Hobert 2006). However, as we already noted when cloning *lsey-6*, the *cog-1* 3' UTR contains a second target site (Johnston and Hobert 2003), which contains only an 8mer seed (Fig. 3) and which is also less conserved in other species (see below). We call the 8mer seed site “site #2.” How important is site #2 for regulation and could this second site explain why a single *lsey-6* site can only work in some, but not all, 3' UTR contexts? We asked whether *lsey-6* site #2 is involved in 3' UTR regulation to an extent that is comparable to that of site #1 by introducing two distinct mutations. A complete deletion of site #2 or introduction of a 4-bp mutation in the seed results in a reduction, but not a complete loss of regulation, which contrasts the complete loss of regulation observed upon deleting site #1 (Fig. 4, nos. 2,4,5; note that numerical values for the degree of regulation, which allow for comparing different constructs, can be found in Supplemental Table 1).

FIGURE 2. Mutational analysis of *lsey-6* site #1. Multiple transgenic lines expressing one given sensor construct are scored for *gfp* expression. Each scored animal is placed into one of three categories, in which relative *gfp* expression in the two ASE neurons is as follows: $ASEL > ASER$ (left black bar), $ASEL = ASER$ (middle gray bar), or $ASEL < ASER$ (right black bar). See Material and Methods for more details on scoring. Each set of three bars represents an independent transgenic line. The gray box shown in each bar graph is a visual “helper” to indicate what is considered to be a minimal level of regulation for the system; this minimal regulation is arbitrarily defined based on the level of regulation seen with the *lsey-6* site #2 deletion, shown in Figure 4, no. 4. The gray box roughly correlates with another quantitative measure for the degree of regulation, called the “asymmetry index,” defined in the Material and Methods and listed in Supplemental Table 1 for each construct analyzed in this paper. Each construct is named based on figure number and number of construct within the figure. The control *unc-54* 3' UTR sensor construct displays characteristic unregulated distribution of *gfp* (no. 1), while the *cog-1* 3' UTR sensors display dramatic regulation with the majority of animals falling into the $ASER > ASEL$ category (no. 2). Shaded boxes indicate location of introduced mismatches. Only mismatches in positions 4, 5, and 6 counting from the 5' end of the miRNA results in complete loss of regulation (nos. 6,7,8). Mismatches in positions 1, 2, 7, 9, 10, and 11 results in little or no loss of regulation (nos. 3,4,9,11–13). Mismatches in positions 3 and 8 results in a moderate loss of regulation (nos. 5,10). If the *lsey-6* target site in the *cog-1* 3' UTR is converted to have perfect complementarity to *lsey-6*, then regulation is completely lost. However, if a single mismatch is reintroduced into position 12, which is unpaired in the wild type, then regulation is restored (nos. 14,15). Note: Construct nos. 1, 2, and 7 have been previously examined (Didiano and Hobert 2006) and a rescoring of newly generated and scored transgenic lines is shown here (rescored lines show the same results as previously reported).




FIGURE 3. (Legend on next page)

Could a lack of site #2 explain the lack of sufficiency of a single *lsey-6* target site to confer regulation to a 3' UTR? To address this issue, we engineered a construct in which we inserted site #1 and site #2 into the *unc-54* 3' UTR. No regulation was observed (Fig. 4, no. 6). Similarly, placing two copies of site #1 into the *unc-54* 3' UTR conferred no regulation (Fig. 4, no. 7; note that the sites are spaced by the same distance as in the *cog-1* 3' UTR). We conclude that even though site #2 has some, comparatively minor, role in 3' UTR regulation, it cannot account for the inability of a *lsey-6* site to confer regulation onto heterologous 3' UTRs. We furthermore note that even within the context of the *cog-1* 3' UTR, the mere presence of two 8 bp seed matches to *lsey-6*, achieved through reduction of the 11mer seed match of site #1 to a 8mer seed (Fig. 4, no. 3), is not sufficient to confer 3' UTR regulation. The presence of one or several seed matches in a 3' UTR is therefore not a sufficient criterion to predict regulation of a 3' UTR by a cognate miRNA.

G:U wobble base pairs in the seed region of the *cog-1* target sites result in impaired regulation

We sought to utilize the minor impact of site #2 on 3' UTR regulation to analyze the issue of the importance of G:U wobble base pairs in miRNA/3' UTR heteroduplexes. Previous work has shown that in a transfection-based cell culture system, G:U wobbles are detrimental for regulation of a synthetic 3' UTR by a nonnatural miRNA (Doench and Sharp 2004). In contrast, we previously reported that G:U wobble base pairs were well tolerated in the seed region of *lsey-6* site #1, even when two G:U wobbles were introduced (Didiano and Hobert 2006). We continued to address this issue using a "sensitized background," that is, the *cog-1* 3' UTR whose regulation is weakened through a loss in *lsey-6* site #2, with the idea being that this may more readily reveal the impact of G:U wobbles on miRNA/target duplex formation. When we combined a G:U wobble base pair in position 6 with a *lsey-6* site #2 deletion, we find that this construct displayed no regulation (Fig. 5, no. 1). Next, we introduced G:U wobbles into both *lsey-6* site #1 and site #2 simultaneously. Three of the five constructs containing G:U wobbles in both target sites displayed regulation, albeit

at levels below wild-type regulation (Fig. 5, nos. 2–6). These results indicate that G:U wobbles can be detrimental to *lsey-6*-mediated regulation in specific circumstances; however, G:U wobbles should not be used as a criterion by which to eliminate potential miRNA/target interactions.

Extra-target site sequences in the *cog-1* 3' UTR influence *lsey-6* mediated regulation

Our analysis has shown that even though *lsey-6* complementary site #1 is an essential feature for *cog-1* 3' UTR regulation, neither single nor multiple *lsey-6* sites are sufficient to confer regulation. An obvious explanation for these observations is that additional features in the *cog-1* 3' UTR are required for miRNA regulation, which, together with the *lsey-6* sites, may also confer regulation on heterologous 3' UTRs. What other features in the *cog-1* 3' UTR could be involved in regulation? One attractive candidate would simply be another miRNA binding site in the *cog-1* 3' UTR. There are 4 potential binding sites for the functionally uncharacterized miRNA *mir-265* in the *cog-1* 3' UTR (Fig. 6). To ask whether *mir-265* is involved in the ASEL-specific down-regulating of the *cog-1* gene, we undertook three separate experiments. First, we assayed the regulation of the wild-type *cog-1* 3' UTR sensor in *mir-265* null mutants (Miska et al. 2007) and found no effect on sensor gene expression (Fig. 6B). Second, we ectopically expressed *mir-265* in ASEL and ASER. The same ectopic expression, when done with *lsey-6*, results in a down-regulation of the *cog-1* target and an ensuing loss of ASER cell fate (Fig. 6C; Johnston and Hobert 2003). However, such an effect cannot be observed upon ectopic expression of *mir-265* (Fig. 6C). Third, we tested whether *cog-1* regulation is disrupted in a *mir-265* null mutant strain by assaying the regulation of a *cog-1* effector gene, *lim-6*. While loss of *lsey-6* leads to a completely penetrant loss of *lim-6* expression, caused by loss of *cog-1* repression (Johnston and Hobert 2003), no such effect can be observed in *mir-265* null mutants (Fig. 6D). We conclude that *mir-265* plays no significant role in regulating the *cog-1* 3' UTR.

As an alternative means to identify potentially relevant sequences in the *cog-1* 3' UTR, we identified phylogenetically conserved sequences in the 3' UTR, utilizing 4

FIGURE 3. Sufficiency of *lsey-6* sites in other 3' UTR contexts. Single copies of the previously described *lsey-6* target site from the *cog-1* 3' UTR were inserted into the *lin-28*, *unc-54*, *lin-41*, *lin-14*, and *actin/act-1* 3' UTRs (Didiano and Hobert 2006). None of the heterologous contexts displayed regulation prior to target site insertion (nos. 1,5,7,9). The primary target site was inserted into two positions of the *lin-28* 3' UTR in place of a presumptive *let-7* target site and a *lin-4* target site (nos. 2,3). Only a single site insertion into the *let-7* site position results in regulation (no. 2) but fails to confer regulation in the second position (no. 3). The target site was also inserted into a *let-7* position in *lin-14* 3' UTR (only *let-7* but not *lin-4* sites are shown for the *lin-14* 3' UTR) but fails to display regulation in this context (no. 8). A single site was insufficient to confer regulation to an *actin* 3' UTR (no. 10). This context was chosen for its short length and high AU content (66% with target site insertion). Additionally, a *cog-1* 3' UTR minimal element, which was defined based on deletion analysis in Figure 8 (nos. 1,2,3), confers regulation to the *actin* 3' UTR but not the *unc-54* 3' UTR (nos. 11,12). Construct nos. 1, 2, and 4 have been previously examined (Didiano and Hobert 2006), and a rescoring of newly generated transgenic lines is shown here (rescored lines show the same results as previously reported). See legend to Figure 2 and Materials and Methods for explanation of data representation and Figure 2, nos. 1 and 2, for regulated and unregulated control 3' UTRs.

different nematode genomes (Fig. 7A). We confirmed that regulation is conserved across phylogeny by generating a sensor construct in which we test the *C. briggsae cog-1* 3' UTR for ASEL-specific down-regulation and find this 3' UTR to indeed be regulated as well (Fig. 7B).

To test the importance of the phylogenetically conserved regions in the 3' UTR, we undertook a deletion analysis of the *C. elegans cog-1* 3' UTR. We find that the first 250 bases of the 394-bp *cog-1* 3' UTR, which contain two phylogenetically conserved sequence motifs (Fig. 7A, dark and light blue boxes), are not required for regulation (Fig. 8, no. 1). Deletion of the last 44 bp (351–394), which contained another, strongly conserved motif (Fig. 7A, purple box), also showed no effect on regulation (Fig. 8, no. 2). Combining the 5' 250-bp deletion with the 3' 44-bp deletion, which removes all phylogenetically conserved motifs aside from the conserved *Isy-6* sites, leads to a modest decrease in regulation, but the 3' UTR still remains well regulated (Fig. 8, no. 3). The remaining 92 bp (258–350) therefore define a minimal regulatable fragment from the *cog-1* 3' UTR, that is not only sufficient to confer regulation in isolation (Fig. 8, no. 3), but also when placed into the context of the *actin* 3' UTR (Fig. 3, no. 11). Notably, though, this minimal element can still not function in the *unc-54* 3' UTR (Fig. 3, no. 12).

Deletions into this 92-bp minimal region lead to a loss of 3' UTR regulation. Specifically, a deletion that removes the last 94 bp (301–394) of the complete *cog-1* 3' UTR (eliminating 50 bp of the 92-bp minimal region) results in the complete loss of regulation (Fig. 8, no. 4). The in vivo importance of this 3' terminal region is underscored by our identification of a mutant allele of the *cog-1* locus in a genetic screen; this mutant allele produces a *cog-1* gain-of-function phenotype, that is, derepression of *cog-1* in ASEL, and harbors a similar deletion at the end of the 3' UTR (105-bp deletion) (Sarin et al. 2007). The last 94 bp of the *cog-1* 3' UTR contain the *Isy-6* site #2; however, the 94-bp deletion has a larger




FIGURE 4. (Legend on next page)

effect on the regulation than a mutation of the *lcy-6* site #2 alone (cf. Fig. 8, no. 4, and Fig. 4, nos. 4,5). This indicates that some sequence in addition to site #2 in the last 94 bp contributes to the regulation. A 50-bp deletion from 301 to 350 bp, encompassing *lcy-6* site #2 (309–324 bp) plus adjacent 25 bp of sequences still resulted in a complete loss of regulation (Fig. 8, no. 5). We term these adjacent 25 bp the “orange box” (see sequence in Fig. 7A). Nineteen out of 25 (76%) base pairs in the orange box are AU base pairs, which is not much above the overall AU content of the entire *cog-1* 3' UTR (65%). A deletion of the 25-bp orange box alone (326–350) results in a modest but significant loss of regulation (Fig. 8, no. 6). Taken together, *lcy-6* site #2 plus an adjacent, modestly conserved 25-bp region, the orange box, are jointly required for *cog-1* regulation; deletion of either sequence results in modest defects, deleting both sites together causes a complete loss of 3' UTR regulation, comparable only to the loss of *lcy-6* site #1. The orange box region does not appear to merely increase the efficacy of the *lcy-6* site #2, as the replacement of site #2 with site #1 in the absence of the orange box does not restore normal regulation (Fig. 8, no. 7).

The orange box acts independently of position and can be substituted with AU- but not GC-rich sequences

One potential function of the orange box could be its binding to a cofactor, either another regulatory RNA or a protein. With the limitation of miRNA target predictions in mind (Rajewsky 2006b), we find that no miRNA is predicted to pair with the orange box. A PUF-type protein was recently shown to be involved in the regulation of a miRNA-controlled 3' UTR (Nolde et al. 2007), and the orange box contains a AUUGUA motif that may constitute a potential PUF-protein binding site (Wharton et al. 1998). However, deleting the AUUGUA motif has no effect on 3' UTR regulation (Fig. 8, no. 8). Next, we deleted chunks of the 25-bp orange box by removing three different 8- to 9-bp regions. None of the smaller deletions recapitulate the effect of the full 25-bp deletion (Fig. 8, nos. 9–11).

FIGURE 4. Asymmetric function of two *lcy-6* sites in the *cog-1* 3' UTR. Construct no. 1 is the same control as shown in Figure 2, no. 2, and is shown for comparison only. A complete deletion (no. 2) or partial deletion resulting in an 8mer seed region of target site #1 (no. 3) each results in a complete loss of regulation, meaning that target site #2 is nonfunctional in isolation. A complete deletion of site #2 (no. 4) or a seed mutant (no. 5) in site #2 each results in a partial loss of regulation. Either two copies of site #1 or a single copy of site #1 in combination with site #2 were inserted into the *unc-54* 3' UTR, but neither of these constructs display regulation (nos. 6,7). miRNA/target heteroduplexes shown here were generated using RNAhybrid (Rehmsmeier et al. 2004). Construct no. 3 has been previously examined (Didiano and Hobert 2006) and a rescoring of newly generated transgenic lines is shown here (rescored lines show the same results as previously reported). See legend to Fig. 2 and Materials and Methods for explanation of data representation.

The addition of a 20-bp insert consisting of only A's or of A's and U's between *lcy-6* site #2 and the orange box does not impair regulation (Fig. 9, nos. 1,2). Intriguingly, the introduction of the AU-insert abrogates the importance of the orange box, that is, deletion of the orange box has no effect on the regulation of the 3' UTR when the heterologous AU-insert is present (Fig. 9, no. 3). In other words, the function of the 25-bp orange box (which contains 25% GC and is therefore not any more or less AU- or GC-rich than the whole *cog-1* 3' UTR) and an arbitrary 20-bp AU-rich sequence are equivalent.

The insertion of a 20-bp GC-rich sequence between *lcy-6* site #2 and the orange box also does not impair regulation (Fig. 9, no. 4). However, in contrast to the inserted AU sequences, the GC sequences cannot substitute for the orange box. That is, a deletion of the orange box in the GC-inserted construct completely abolishes regulation (Fig. 9, no. 5).

At first sight, one interpretation of these findings may be that the 25-bp orange box provides an energetically favorable context for *lcy-6* site #2 to function in. This explanation would be in accordance with the recently deduced nonstructured and AU-rich nature of sequences that flank miRNA binding sites (Zhao et al. 2005; Grimson et al. 2007; Zhao et al. 2007). However, we note that this explanation is rendered unlikely by our finding that the orange box and *lcy-6* site #2 function independently, that is, the effect of the loss of *lcy-6* site #2 is enhanced by loss of the orange box (cf. Fig. 4, no. 4, and Fig. 8, nos. 5,6). Moreover, in the context of the GC insertion (which cannot substitute for the orange box), the orange box can still function from the distance, thereby ruling out that the orange box provides a proximal, local context for site #2 to function.

The linker region between the two target sites is required for wild-type regulation

Are there other regions besides the orange box that are important for *cog-1* 3' UTR function? A previous study on *let-7*-mediated 3' UTR regulation has found that both the composition and the distance between two *let-7* target sites in the *lin-41* 3' UTR can influence miRNA-mediated regulation (Vella et al. 2004a). We tested whether the linker region between the two *lcy-6* target sites in the *cog-1* 3' UTR influences 3' UTR regulation in a number of manners (Fig. 10). To test whether the base sequence composition of the linker affected regulation, we replaced the 26-bp linker between the two sites (35% GC content) with a 26-bp linker consisting entirely of A and U base pairs.




FIGURE 5. G:U wobble base pairing in the seed region impairs but does not necessarily abolish regulation. Construct no. 1 consists of a G:U wobble base pair in position 6 of site #1 in the background of a site #2 deletion and fails to display regulation. Construct nos. 2–6 were engineered to contain G:U wobble base pairs in both *Isy-6* site #1 and site #2. Three of five of these double G:U wobble constructs display intermediate regulation (nos. 2,4,5). Shaded boxes indicate location of G:U wobbles introduced. See legend to Fig. 2 and Materials and Methods for explanation of data representation and Figure 2, nos. 1 and 2, for regulated and unregulated control 3' UTRs.

This mutation results in only a reduction, but not elimination, of regulation (Fig. 10, no. 2). Similarly, an inversion of the linker sequence only slightly affects regulation (Fig. 10, no. 3). A 40-bp extension of the linker also has only minimal effects on regulation (Fig. 10, no. 1). This indicates that the sequence composition and length of the linker region have only a minor impact on regulation, which is in contrast to the *let-7/lin-41* 3' UTR case, in which the linker region was shown to be absolutely required for regulation (Vella et al. 2004a).

Although the composition of the linker does not appear to have a strong impact on 3' UTR regulation, its mere presence does. Deleting 20 bp from the 26-bp linker region, results in a complete loss of regulation (Fig. 10, no. 4). However, two smaller separate 10-bp deletions that cover the essential 20-bp region have small or no effect on regulation (Fig. 10, nos. 5,6). Together with the results of the linker inversion, these results argue that in contrast to the *lin-41/let-7* linker region, which cannot be replaced by heterologous sequences (Vella et al. 2004a), the linker region between the two *Isy-6* sites tolerates significant sequence alterations.

Taken together, we have shown that two regions that flank the two *Isy-6* binding sites, the orange box 3' of *Isy-6* site #2 and the linker region 3' of *Isy-6* site #1, have an impact on 3' UTR regulation. Both flanking regions function independently and together are essential for 3' UTR regulation, as we find that combining the 26-bp linker inversion mutation, which alone has only a minimal effect on regulation (Fig. 10, no. 3), with a deletion of the orange box, which alone has small effects on regulation (Fig. 8, no. 6), results in an almost complete loss of regulation (Fig. 10, no. 7). It is notable that this construct has two intact *Isy-6* target sites #1 and #2, yet it is not regulated. These results indicate that the *Isy-6* target sites are only functional when present within an appropriate 3' UTR context.




FIGURE 6. *mir-265* does not affect *cog-1* activity. (A) Potential target sites for *mir-265* in the *cog-1* 3' UTR shown below 3' UTR schematic, with *Isy-6* target sites shown above. (B) The *cog-1* 3' UTR sensor *otIs185* (Sarin et al. 2007) is down-regulated normally in the ASEL neuron in *mir-265*(*n4534*) null mutant animals. The degree of regulation is the same as shown for the same sensor in a wild-type background (Sarin et al. 2007). (C) The ASE bilateral *ceh-36* promoter was used to drive expression of *Isy-6* and *mir-265* hairpin precursors. *ntlIs1*(*gcy-5::gfp*) was used as a cell-fate marker to test the ability of *Isy-6* and *mir-265* to down-regulate *cog-1*, thereby driving ASEL cell fate in ASER (green circles represent *gcy-5::gfp* expression pattern as ASER-specific [top], ASE bilateral [middle], or loss of expression [bottom]). All three lines misexpressing *Isy-6* results in a “two ASEL phenotype,” as previously reported (Johnston and Hobert 2003). None of the three lines misexpressing *mir-265* results in a phenotype, indicating that *mir-265* is incapable of downregulating *cog-1* via its 3' UTR in ASER. (D) The *mir-265* null mutant *n4534* shows no *Isy* phenotype using the *lim-6::gfp* ASEL specific reporter (*otIs114*). Green circles indicate reporter expression in ASEL and ASER.

Isy-6 target sites are only fully functional in their endogenous positions within the *cog-1* 3' UTR

We further addressed the importance of position and context of the *Isy-6* sites by moving site #1 and site #2 into different positions with the otherwise wild-type *cog-1* 3' UTR. We find that neither site #1 nor site #2 is functional

when placed at a position in which the 3' end of the target site is 70 bp 3' of the stop codon in the *cog-1* 3' UTR. This distance from the stop is much above the recently proposed minimal distance of a miRNA target site to a stop codon and also is located in an “off-center position,” which was also suggested to be favorable for miRNA target site function (Grimson et al. 2007). In spite of these apparently

favorable conditions, moving site #1 in this position leads to a complete loss of regulation (Fig. 11, no. 1). Moving site #2 leads to the same moderate loss of regulation observed upon entirely deleting site #2 (cf. Fig. 11, no. 3, and Fig. 4, no. 4). When moved to another location in the *cog-1* 3' UTR, site #1 confers some limited, but clearly impaired level of regulation (Fig. 11, no. 2), while another location for site #2 has the same effects as having no site #2 at all (Fig. 11, no. 4).

The base-pair composition surrounding the target sites does not influence regulation in a manner correlating with AU or GC richness

What is it exactly about the *lgy-6* target site context that is so important for regulation? Several reports have indicated that miRNA target sites reside in AU-rich regions of 3' UTRs (Robins and Padgett 2005; Grimson et al. 2007). High ΔG values, favored by high AU content, of sequences around the target site have also been proposed to be important for functionality (Zhao et al. 2007). The primary *lgy-6* target site #1 normally resides in a location with 83% AU content (over 30 bp) 5' to the site, while the 26-bp linker sequence 3' to site #1 is only 58% AU in content, which is comparable to the overall AU content of the *cog-1* 3' UTR (65%) (Fig. 7A). When we alter site #1 context by adding 20 bp of GCs 5' to the site #1, thereby dramatically changing local AU content, we observe no significant effect on regulation (Fig. 12, no. 1). As mentioned above in the context of studying the orange box, the introduction of GC sequences 3' of *lgy-6* site #2 also has little effect on regulation (Fig. 9, no. 4).

We addressed this issue further by attempting to “activate” the nonfunctional *lgy-6* target sites in the *unc-54* 3' UTR context (described above in Figs. 3, no. 4; Figs. 4, no. 7), by surrounding a single *lgy-6* site #1 seed region in the context of the *unc-54* 3' UTR with >20-bp AU-rich sequences. This manipulation does not confer regulation on the 3' UTR (cf. Fig. 3, no. 4, and Fig. 12, no. 2). We also changed the construct described above with two *lgy-6* site #1, placed into the *unc-54* 3' UTR, which conferred no regulation (Fig. 4, no. 7). If both sites are embedded into an AU-rich context, regulation is still not observed (Fig. 12, no. 3). Taken together, these results indicate that in the case of the *lgy-6/cog-1* 3' UTR interaction, no simple trend involving the base pair composition around the target sites correlates with the extent of regulation of the 3' UTR.

Finally, we generated a synthetic 3' UTR construct consisting of two copies of site #1 with a 26-bp AU linker and CU bp 5' and 3' to the target sites (Fig. 12, no. 4). The CU flanking sequences were designed to prevent secondary structure around the target sites. This construct displays a small level of regulation, which is not comparable to the level of regulation observed with the intact *cog-1* 3' UTR. Based on these incomplete levels of regulation, we conclude that in those cases in which *lgy-6* sites do not confer




FIGURE 8. (Legend on next page)

regulation (e.g., in *unc-54*, *actin*, *lin-41*, *lin-14* 3' UTR context) this is not because these 3' UTR sequences inhibit *lgy-6* sites from functioning. Rather, in those contexts where *lgy-6* site(s) are functional (i.e., the *cog-1* 3' UTR context), contextual features enable the *lgy-6* site(s) to function.

Predicting experimental results

Based on theoretical considerations and/or cell culture based transfection assays, researchers in several different laboratories have developed means to assess whether a miRNA may exert regulation on a putative target sequence. Zhao et al. (2005, 2007) have proposed that the energetic features (ΔG of folding) of sequences surrounding a miRNA target site are a predictor for miRNA/target interactions. Long et al. (2007) and Kertesz et al. (2007) utilized overall folding properties of the mRNA target to predict miRNA/target interaction. Grimson et al. (2007) reported that local AU content provides a means to assess successful miRNA/target regulation. We do not find any strong correlation between any of these parameters and the presence and strength of regulation of the many constructs we have generated in this study (Supplemental Table 1).

We also tested each 3' UTR sequence for predicted folding patterns and observed no notable correlation between regulation and predicted secondary structure (Supplemental Table 1). That is, we found that in some regulated 3' UTR, the *lgy-6* sites lie within secondary structure elements while in several nonregulated 3' UTR, the sites are exposed (Supplemental Table 1). In sum, the ability to predict whether a 3' UTR is regulated by a miRNA remains limited.

DISCUSSION

We have described here 156 transgenic lines that express 71 different 3' UTR sensor constructs and assessed the ability of an endogenously expressed miRNA, acting within its

normal cellular context, to regulate these sensor constructs. We draw the following conclusions from our studies.

Seed matches are not a sufficient predictor for miRNA/target interactions

This notion stems from the following results. First, through mutating the 5' half (3' in reference to the miRNA) of the first *lgy-6* site, we have generated a 3' UTR that contains two perfect 8mer seed matches to *lgy-6* but is not regulated (Didiano and Hobert 2006). Second, we have shown that 13 3' UTRs with seed matches to *lgy-6* are not regulated by *lgy-6* (Didiano and Hobert 2006). We have extended this finding here by showing that the two *lgy-6* sites do not function in a variety of different contexts, when surrounding sequences are mutated, when transplanted into different, heterologous 3' UTRs, or when moved within the *cog-1* 3' UTR. Our results are consistent with a previous, frequently overlooked study by Vella et al. (2004a) that demonstrated that an engineered 3' UTR with multiple seed matches to the *let-7* miRNA fails to display down-regulation in a natural cellular context.

lgy-6 requires imperfectly complementary target sites to mediate down-regulation of the *cog-1* 3' UTR

A *cog-1* 3' UTR sensor with a perfectly complementary *lgy-6* target site displays no down-regulation, indicating that *lgy-6* is primarily entering the ALG-1/ALG-2 miRNA processing pathway and that the bulged base pairs in the miRNA/target heteroduplex are required by the ALG-1/ALG-2 complex for efficient recognition. These in vivo observations bolster recent biochemical work that has shown that depending on the complementarity of the hairpin precursor structure miRNA or siRNAs get sorted into distinct silencing complexes (Forstemann et al. 2007; Steiner et al. 2007; Tomari et al. 2007). This sorting is not 100% discriminative as we observe that within a heterologous context (*unc-54* 3' UTR), three repeated, perfectly complementary *lgy-6* sites confer a *rde-1*-dependent, though modest, down-regulation of the sensor in ASEL (L. Cochella and O. Hobert, unpubl.). In light of these issues, the previously reported use of "perfect sensors" to detect sites of endogenous miRNA expression (Mansfield et al. 2004) needs to be viewed with a substantial degree of caution.

G:U wobbles can be tolerated in the *lgy-6/cog-1* heteroduplex

We have shown that even though G:U wobble base-pairing can weaken miRNA/target interaction, G:U base-pairing does not abrogate regulation. Even if G:U wobbles are engineered into both *lgy-6* sites, regulation is still preserved. The presence of G:U base pairs should therefore not be

FIGURE 8. Mutational dissection of the *cog-1* 3' UTR. The first 250 bp and last 44 bp of the *cog-1* 3' UTR are not required for regulation (nos. 1,2). and base pairs 258–350 of the *cog-1* 3' UTR alone are sufficient for regulation (no. 3). However, this minimal element is not sufficient in all heterologous contexts (nos. 11,12). A deletion of the last 94 bp of the *cog-1* 3' UTR results in a complete loss of regulation (no. 4). A deletion of 300–350 bp results in a complete loss of regulation (no. 5). Deletion of 325–350 bp results in a partial loss of regulation (no. 6). This region does not affect *lgy-6* target site #2. Replacing site #2 with a second copy of site #1 in the background of a 325- to 350-bp deletion resulted in slightly increased, but sub-wildtype regulation (no. 7). This 25-bp motif contains a potential PUF-protein binding site, but a deletion of this site had no effect on regulation (no. 8). Scanning deletion analysis of the 25-bp motif by two 8-bp and a 9-bp deletion did not recapitulate the 25-bp deletion effect on regulation (nos. 9,10,11). See legend to Figure 2 and Materials and Methods for explanation of data representation and Figure 2, nos. 1 and 2, for regulated and unregulated control 3' UTRs.




FIGURE 9. Functional analysis of the orange box. Distancing the 25-bp orange box from *Lsy-6* site #2 by a 20-bp A or a AU spacer does not affect regulation (nos. 1,2). Deleting the orange box in the presence of the 20-bp AU spacer does not result in a reduction of regulation (no. 3). Distancing the 25-bp orange box from site #2 with a 20-bp GC spacer also does not affect regulation (no. 4), but deleting the orange box in this context results in a complete loss of regulation (no. 5). See legend to Fig. 2 and Materials and Methods for explanation of data representation and Figure 2, nos. 1 and 2, for regulated and unregulated control 3' UTRs.

categorically used to exclude miRNA/target interactions. This notion is also supported by G:U wobbles being present and functionally required in *let-7/lin-41* target sites (Vella et al. 2004b).

Asymmetry of multiple target sites

The *let-7/lin-41* miRNA target interaction was the first experimentally validated example of a synergistic interaction of two miRNA sites (Vella et al. 2004a). Deletion of either of two *let-7* site in the *lin-41* target 3' UTR causes a loss of 3' UTR regulation. More recent studies further underscored the potency with which two miRNA sites can cooperate to induce 3' UTR regulation (Grimson et al.

2007; Saetrom et al. 2007). The *cog-1* 3' UTR also contains two *Lsy-6* sites but their functional importance is highly asymmetric. Only deletion of one site strongly affects 3' UTR regulation; loss of the other site only has a limited effect. Moreover, the spacing of the two sites does not conform with the optimal spacing of cooperating sites; not only are the two *Lsy-6* sites more distantly spaced than previously reported examples (Saetrom et al. 2007), but their spacing can be even further increased without significant loss of 3' UTR regulation. Also, in one other 3' UTR context, that of *lin-28*, *Lsy-6* site #1 can also function alone. *Lsy-6* site #2 likely works via *Lsy-6* binding, rather than by merely providing some structural context, as mutating the seed disrupts its function. One conceivable function of *Lsy-6* site #2 is to provide a recruitment platform for *Lsy-6*. Such a recruitment model has been proposed to be important in opening secondary structural elements and has proven powerful in correctly predicting several in vivo miRNA/target interactions (Long et al. 2007).

Another difference between the multiple target site configuration of the *lin-41* 3' UTR and the *cog-1* 3' UTR is that the linker sequence between the two *let-7* sites needs to be of specific length and sequence content (Vella et al. 2004a). In contrast, we can lengthen or invert the linker sequence between the two *Lsy-6* sites with little effect on regulation.

AU richness or unstructuredness is not a sufficient predictor for *Lsy-6* target site functionality

Recent bioinformatics analysis combined with transfection studies argues that miRNA target site efficacy depends on surrounding AU-rich sequences (Grimson et al. 2007). We find that this is not the case for *Lsy-6*. Decreasing the AU richness of the *Lsy-6* site #1 does not impair 3' UTR regulation, and inclusion of AU-rich flanking sequences to *Lsy-6* sites in a 3' UTR that is normally refractory to regulation does not enable regulation. Moreover, an analysis of all constructs generated in this study does not show any strong correlation between AU-richness and regulation (Supplemental Table 1).

Contextual features that promote miRNA activity

A number of constructs that we have generated in this study argue for the overall importance of 3' UTR features beyond miRNA target sites, which act in conjunction with miRNA target sites and need to be present in a specific configuration relative to miRNA binding sites. We have defined two contextual features that are critical for *cog-1* 3' UTR function, both of which are located 3' to the two *lisy-6* sites, namely, a linker region between *lisy-6* site #1 and #2 and a region located downstream of *lisy-6* site #2, the "orange box." Both sequences are AU-rich, but not more AU-rich than the whole *cog-1* 3' UTR. Neither sequence appears to carry overtly specific sequence information, as they each can be replaced by AU-only sequences (or, in the case of the linker, can be inverted). Also, neither sequence is absolutely required for 3' UTR regulation in isolation, but when affected in combination (inversion of the linker+deletion of the orange box), a complete loss of 3' UTR regulation is observed.

The two *lisy-6* sites and the linker region and orange box are alone sufficient to impart robust regulation on a reporter gene and can also convert the normally nonregulated 3' UTR of the *actin* gene into a miRNA-regulated 3' UTR (Fig. 3). Notably, the two *lisy-6* sites and the linker region and orange box need to be present in a specific configuration, as moving either of the *lisy-6* sites out of their normal context results in defects comparable to the deletion of the respective site. However, at least the orange box (and perhaps also the linker region, which we did not test) need not be in absolute proximity to the *lisy-6* site #2 as the insertion of either AU-only or GC-only sequences between site #2 and the orange box does not affect orange box function.

Given their similar features and relative placement it is conceivable that the linker and orange box regions may fulfill similar functions. As those functions may not be entirely sequence specific and as many RNA-binding proteins display limited specificity in RNA binding (Fierro-Monti and Mathews

2000), it is conceivable that each region may provide a binding platform for an RNA binding protein. Such protein binding is unlikely to function to improve the recruitment of *lisy-6* to its cognate sites, as illustrated by the independent action of site #2 and the orange box. If the orange box were to merely serve to recruit *lisy-6* to site #2, then the function




FIGURE 10. (Legend on next page)

of the orange box would be obsolete if site #2 were removed. To the contrary, we find that the effect of a deletion of site #2 is enhanced by loss of the orange box, arguing for an independent function of the orange box. The binding of miRNAs and other factors may thus be independent events and, in analogy to transcriptional enhancosomes (Merika and Thanos 2001), a joint assembly of multiple *trans*-acting factors, miRNAs, and proteins may be required to exert regulatory control over a 3' UTR. Alternatively, the linker region and orange box may cooperate to provide a structural context that allows for miRNA-mediated regulation.

miRNA target predictions

We find a limited correlation with target prediction algorithms that allow the user to provide 3' UTR sequences (StarMir, Pita). Target genes predicted by other prediction programs also did not show regulation in our assay system (Didiano and Hobert 2006). Other features with proposed predictive values such as target-side surrounding AU-richness or target-side surrounding ΔG values did also not accurately predict our experimental results. Current prediction algorithms calculate target efficacy based on interactions between the mRNA with itself and the mRNA with a miRNA. However, these interactions are occurring in a complex cellular environment in which mRNAs and miRNAs are likely bound by cellular RNA-binding proteins, which is currently impossible to account for *in silico*. Genomes across phylogeny encode multitudes of proteins of unknown function with predicted RNA-binding motifs. One member of a large family (>100 members in *C. elegans*) of RNA binding proteins was in fact recently shown to control miRNA binding efficacy to its target mRNA (Kedde et al. 2007).

Is *lisy-6* an exemplary miRNA?

At this point it is not clear whether lessons learned from the *lisy-6/cog-1* target interaction are generally applicable to other miRNA/target interactions. However, if there is one lesson to be learned from the history of the miRNA field, it is that one should not discard individual case studies as mere oddities (Ruvkun et al. 2004). As our analysis revealed

a striking reliance on some presently rather ill-defined contextual features of the 3' UTR, we consider it of paramount importance to analyze miRNA/target interaction in a cellular environment in which the miRNA and target normally interact. Such cellular environment may impart on the miRNA/target interaction by providing the correct set of cofactors, such as mRNA-binding proteins. Studies that place miRNAs and their target into heterologous context and also often rely on overexpression of the two reactants provide, in our opinion, no more generalizable results than the single miRNA/target interaction that we have dissected here.

CONCLUSIONS

Our study indicates that miRNA/target interactions rely on features beyond the miRNA complementary site in a target gene, a situation that is reminiscent of transcription factor/target recognition. Transcription factor binding sites abound in genome sequences, yet only a limited set are occupied and utilized *in vivo*, apparently due to relatively poorly understood contextual features of genomic DNA sequence (see for example, Etchberger et al. 2007). Our results therefore raise the possibility that the critical role of 3' UTR context may limit the number 3' UTRs that may be regulated by a given miRNA and therefore that less genes in a genome are regulated by miRNAs than currently anticipated.

MATERIAL AND METHODS

DNA constructs

All 3' UTR sensor constructs are based on the same vector platform generated by cloning 1.8 kb of the *ceh-36* promoter (1883 bp to 36 bp upstream of the ATG start codon) into the Hind III/BamH1 sites of the canonical pPD95.75 *gfp* vector, kindly provided by Andy Fire (Stanford University). All tested 3' UTRs were generated by cloning into the EcoRI/EagI restriction sites. This process removes the *unc-54* 3' UTR and replaces it with experimental 3' UTRs. All mutant variants of the *cog-1* 3' UTR derivatives of a cloned PCR product from genomic DNA isolated from the N2 wild-type strain isolate. All heterologous 3' UTRs, with the exception of *unc-54*, which is already present in pPD95.75, were also amplified and cloned from N2 genomic DNA. For the *lin-28* 3' UTR constructs, we did not include the first 50 bp of the 3' UTR since it contained a cryptic *lisy-6* site with a poor seed match (5mer) but relatively favorable energetic pairing. With the exception of those constructs described below, mutations were introduced by PCR fusion. The 3' UTR is amplified as two individual PCR fragments using internal primers, which are overlapping and contain the mutation of interest. These PCR fragments are used as a template in a second round of PCR using the

FIGURE 10. A 26-bp linker between *lisy-6* site #1 and #2 provides contextual information that is length independent. Insertion of a 40-bp AU spacer 3' to the endogenous 26-bp spacer between the two target sites in the *cog-1* 3' UTR results in a partial loss of regulation (no. 1). Replacing the endogenous linker with a 26-bp AU linker (no. 2) or inverting the linker also results in a partial loss of regulation (no. 3). A 20-bp deletion that shortens the linker to 6 bp results in a complete loss of regulation (no. 4). Only one of two separate 10-bp deletions of the linker region results in a reduction of regulation (no. 5). Combining the linker inversion mutation with a deletion of the orange box, examined in Figures 8 and 9, renders the *cog-1* 3' UTR virtually nonfunctional (no. 7). See legend to Figure 2 and Materials and Methods for explanation of data representation and Fig. 2, nos. 1 and 2, for regulated and unregulated control 3' UTRs.




FIGURE 11. *Isy-6* target sites in the *cog-1* 3' UTR are only fully functional in endogenous positions in the 3' UTR. *Isy-6* target site #1 displays no regulation when moved to one position but displays weak regulation when moved to a second position within the *cog-1* 3' UTR (nos. 1,2). Similarly, moving site #2 to the same positions within the *cog-1* 3' UTR results in a level of regulation similar to those displayed by a site #2 deletion (nos. 3,4, see also Fig. 4, no. 4). This indicates that site #1 and site #2 are only capable of achieving wild-type levels of regulation when they are in their endogenous positions. See legend to Figure 2 and Materials and Methods for explanation of data representation.

same outside primers containing the restrictions sites (Hobert 2002). PCR fusion products were subcloned into the EcoRI/EagI sites mentioned above. Some mutations near the 3' end of the 3' UTR were introduced using long primers containing the mutation of interest and the EagI restriction site (e.g., Fig. 8, nos. 5,7,11). In Figure 3, construct no. 12 has the *cog-1* 3' UTR minimal element inserted into a vector linker region 3' of *gfp* and 5' of the *unc-54* 3' UTR in pPD95.75. In Figure 12, construct no. 3 is a shortened *unc-54* 3' UTR; this construct was made by PCR fusion with the internal primers annealing to AU-rich sequences in the *unc-54* 3' UTR, thereby removing the intervening sequence and replacing it with the target site inserts. In Figure 12, construct no. 4 was made by cloning annealed complementary primers containing restriction sites. Constructs involving multiple mutations were made using a cloned 3' UTR containing one of the mutations as a template in a second round of PCR fusion-based mutagenesis. All constructs were sequenced to confirm introduction of mutation. The sequence of all constructs can be found in the Supplementary Information.

For expression of *Isy-6* and *mir-265* in both ASEL and ASER, two constructs, *ceh-36::Isy-6* and *ceh-36::mir-265*, were generated. Both microRNAs hairpins were amplified from N2 genomic DNA

and either directly fused to the *ceh-36* promoter by PCR fusion or first cloned into pPD95.75 containing the *ceh-36* promoter and then PCR-amplified before injection.

Transgenic lines

Transgenic lines were generated by injecting the 3' UTR sensor constructs at 5 ng/ μ L with *rol-6(d)* at 100 ng/ μ L as an injection marker into the gonad of wild-type N2 animals. Stable transgenic lines were selected in the F2 generation.

ceh-36::Isy-6 and *ceh-36::mir-265* were injected as PCR products at 40 ng/ μ L with *rol-6* (100 ng/ μ L) into *ntl-1(gcy-5::gfp)* animals.

Scoring 3' UTR regulation

All strains were maintained at 20°C prior to scoring. All lines were scored blind to genotype under a Zeiss Axioplan 2 microscope. To minimize the inclusion of mosaic animals that may have lost the reporter in either ASEL or ASER or both lineages, only those animals were scored in which expression in the AWCL and AWCR neurons was observed (see *gfp* image in Fig. 1). The fluorescence intensity of the 3' UTR sensor constructs is compared between ASEL and ASER in each individual animal. This assay is quantitative in that it quantifies the percentage of animals whose *gfp* expression pattern falls into one of three categories: ASEL > ASER, ASEL = ASER, and ASEL < ASER. Therefore, our assay quantifies regulation levels over a population based on qualitative differences

in expression levels in individual animals. We express our sensor constructs at a low concentration resulting in dim expression levels that are ideal for discerning regulation. We have spot-checked our qualitative assessment of *gfp* expression using AxioVision imaging software (Zeiss, Inc.) that measures the pixelation intensity of ASEL versus ASER. We found that our visual qualitative scoring of each individual animal is sensitive to 25% differences in *gfp* expression. Importantly, this 25% limit of sensitivity is well below the inherent noise levels of the in vivo system, in which variation of *gfp* expression intensity from an extrachromosomal array varies from transgenic line to transgenic line and from animal to animal within the same transgenic line. This noise is apparent in nonregulated control *unc-54* 3' UTR; ~15%–30% of animals show ASEL > ASER *gfp* expression but about the same number of animals also shows ASEL < ASER *gfp* expression. This noise in the system makes the assay extremely sensitive to regulation. Whenever a 3' UTR is *Isy-6*-regulated, these “noise categories” are transformed into striking levels of differences in *gfp* intensity in ASEL versus ASER; i.e., there are still wild-type levels of fluorescence intensity in ASER, but fluorescence is dramatically down-regulated in ASEL. Additionally, those constructs that display regulation do so consistently across transgenic

discussions and comments on the manuscript. This work was funded by NIH R01 NS050266-01 and 2R01NS039996-05 to O.H. and by an NIH predoctoral fellowship to D.D.

Received March 17, 2008; accepted April 3, 2008.

REFERENCES

- Bartel, D.P. 2004. MicroRNAs. Genomics, biogenesis, mechanism, and function. *Cell* **116**: 281–297.
- Beitzinger, M., Peters, L., Zhu, J.Y., Kremmer, E., and Meister, G. 2007. Identification of human microRNA targets from isolated argonaute protein complexes. *RNA Biol.* **4**: 76–84.
- Bentwich, I., Avniel, A., Karov, Y., Aharonov, R., Gilad, S., Barad, O., Barzilai, A., Einat, P., Einav, U., Meiri, E., et al. 2005. Identification of hundreds of conserved and nonconserved human microRNAs. *Nat. Genet.* **37**: 766–770.
- Brennecke, J., Stark, A., Russell, R.B., and Cohen, S.M. 2005. Principles of microRNA-target recognition. *PLoS Biol.* **3**: e85. doi: 10.1371/journal.pbio.0030085.
- Bushati, N. and Cohen, S.M. 2007. microRNA functions. *Annu. Rev. Cell Dev. Biol.* **23**: 175–205.
- Carthew, R.W. 2006. Gene regulation by microRNAs. *Curr. Opin. Genet. Dev.* **16**: 203–208.
- Chang, S., Johnston Jr., R.J., and Hobert, O. 2003. A transcriptional regulatory cascade that controls left/right asymmetry in chemosensory neurons of *C. elegans*. *Genes & Dev.* **17**: 2123–2137.
- Didiano, D. and Hobert, O. 2006. Perfect seed pairing is not a generally reliable predictor for miRNA-target interactions. *Nat. Struct. Mol. Biol.* **13**: 849–851.
- Doench, J.G. and Sharp, P.A. 2004. Specificity of microRNA target selection in translational repression. *Genes & Dev.* **18**: 504–511.
- Doench, J.G., Petersen, C.P., and Sharp, P.A. 2003. siRNAs can function as miRNAs. *Genes & Dev.* **17**: 438–442.
- Etchberger, J.F., Lorch, A., Sleumer, M.C., Zapf, R., Jones, S.J., Marra, M.A., Holt, R.A., Moerman, D.G., and Hobert, O. 2007. The molecular signature and cis-regulatory architecture of a *C. elegans* gustatory neuron. *Genes & Dev.* **21**: 1653–1674.
- Fierro-Monti, I. and Mathews, M.B. 2000. Proteins binding to duplexed RNA: One motif, multiple functions. *Trends Biochem. Sci.* **25**: 241–246.
- Forstemann, K., Horwich, M.D., Wee, L., Tomari, Y., and Zamore, P.D. 2007. *Drosophila* microRNAs are sorted into functionally distinct argonaute complexes after production by dicer-1. *Cell* **130**: 287–297.
- Griffiths-Jones, S., Grocock, R.J., van Dongen, S., Bateman, A., and Enright, A.J. 2006. miRBase: microRNA sequences, targets and gene nomenclature. *Nucleic Acids Res.* **34**: D140–D144. doi: 10.1093/nar/gkj112.
- Grimson, A., Farh, K.K., Johnston, W.K., Garrett-Engele, P., Lim, L.P., and Bartel, D.P. 2007. MicroRNA targeting specificity in mammals: Determinants beyond seed pairing. *Mol. Cell* **27**: 91–105.
- Ha, I., Wightman, B., and Ruvkun, G. 1996. A bulged *lin-4/lin-14* RNA duplex is sufficient for *Caenorhabditis elegans lin-14* temporal gradient formation. *Genes & Dev.* **10**: 3041–3050.
- Hobert, O. 2002. PCR fusion-based approach to create reporter gene constructs for expression analysis in transgenic *C. elegans*. *Bio-techniques* **32**: 728–730.
- John, B., Enright, A.J., Aravin, A., Tuschl, T., Sander, C., and Marks, D.S. 2004. Human microRNA targets. *PLoS Biol.* **2**: e363. doi: 10.1371/journal.pbio.0020363.
- Johnston, R.J. and Hobert, O. 2003. A microRNA controlling left/right neuronal asymmetry in *Caenorhabditis elegans*. *Nature* **426**: 845–849.
- Kedde, M., Strasser, M.J., Boldajipour, B., Vrieland, J.A., Slanchev, K., le Sage, C., Nagel, R., Voorhoeve, P.M., van Duijse, J., Orom, U.A., et al. 2007. RNA-binding protein Dnd1 inhibits microRNA access to target mRNA. *Cell* **131**: 1273–1286.
- Kertesz, M., Iovino, N., Unnerstall, U., Gaul, U., and Segal, E. 2007. The role of site accessibility in microRNA target recognition. *Nat. Genet.* **39**: 1278–1284.
- Krek, A., Grun, D., Poy, M.N., Wolf, R., Rosenberg, L., Epstein, E.J., MacMenamin, P., da Piedade, I., Gunsalus, K.C., Stoffel, M., et al. 2005. Combinatorial microRNA target predictions. *Nat. Genet.* **37**: 495–500.
- Lall, S., Grun, D., Krek, A., Chen, K., Wang, Y.L., Dewey, C.N., Sood, P., Colombo, T., Bray, N., Macmenamin, P., et al. 2006. A genome-wide map of conserved microRNA targets in *C. elegans*. *Curr. Biol.* **16**: 460–471.
- Lee, R.C., Feinbaum, R.L., and Ambros, V. 1993. The *C. elegans* heterochronic gene *lin-4* encodes small RNAs with antisense complementarity to *lin-14*. *Cell* **75**: 843–854.
- Lewis, B.P., Burge, C.B., and Bartel, D.P. 2005. Conserved seed pairing, often flanked by adenosines, indicates that thousands of human genes are microRNA targets. *Cell* **120**: 15–20.
- Long, D., Lee, R., Williams, P., Chan, C.Y., Ambros, V., and Ding, Y. 2007. Potent effect of target structure on microRNA function. *Nat. Struct. Mol. Biol.* **14**: 287–294.
- Mansfield, J.H., Harfe, B.D., Nissen, R., Obenaus, J., Srineel, J., Chaudhuri, A., Farzan-Kashani, R., Zuker, M., Pasquinelli, A.E., Ruvkun, G., et al. 2004. MicroRNA-responsive “sensor” transgenes uncover Hox-like and other developmentally regulated patterns of vertebrate microRNA expression. *Nat. Genet.* **36**: 1079–1083.
- Merika, M. and Thanos, D. 2001. Enhanceosomes. *Curr. Opin. Genet. Dev.* **11**: 205–208.
- Miska, E.A., Alvarez-Saavedra, E., Abbott, A.L., Lau, N.C., Hellman, A.B., McGonagle, S.M., Bartel, D.P., Ambros, V.R., and Horvitz, H.R. 2007. Most *Caenorhabditis elegans* microRNAs are individually not essential for development or viability. *PLoS Genet.* **3**: e215. doi: 10.1371/journal.pgen.0030215.
- Moss, E.G., Lee, R.C., and Ambros, V. 1997. The cold shock domain protein LIN-28 controls developmental timing in *C. elegans* and is regulated by the *lin-4* RNA. *Cell* **88**: 637–646.
- Nolde, M.J., Saka, N., Reinert, K.L., and Slack, F.J. 2007. The *Caenorhabditis elegans pumilio* homolog, *puf-9*, is required for the 3' UTR-mediated repression of the *let-7* microRNA target gene, *hbl-1*. *Dev. Biol.* **305**: 551–563.
- Notredame, C., Higgins, D.G., and Heringa, J. 2000. T-Coffee: A novel method for fast and accurate multiple sequence alignment. *J. Mol. Biol.* **302**: 205–217.
- Palmer, R.E., Inoue, T., Sherwood, D.R., Jiang, L.I., and Sternberg, P.W. 2002. *Caenorhabditis elegans cog-1* locus encodes GTX/Nkx6.1 homeodomain proteins and regulates multiple aspects of reproductive system development. *Dev. Biol.* **252**: 202–213.
- Rajewsky, N. 2006a. L(ou)sy miRNA targets? *Nat. Struct. Mol. Biol.* **13**: 754–755.
- Rajewsky, N. 2006b. microRNA target predictions in animals. *Nat. Genet. (Suppl)* **38**: S8–13.
- Rehmsmeier, M., Steffen, P., Hochsmann, M., and Giegerich, R. 2004. Fast and effective prediction of microRNA/target duplexes. *RNA* **10**: 1507–1517.
- Reinhart, B.J., Slack, F.J., Basson, M., Pasquinelli, A.E., Bettinger, J.C., Rougvie, A.E., Horvitz, H.R., and Ruvkun, G. 2000. The 21-nucleotide *let-7* RNA regulates developmental timing in *Caenorhabditis elegans*. *Nature* **403**: 901–906.
- Robins, H., Li, Y., and Padgett, R.W. 2005. Incorporating structure to predict microRNA targets. *Proc. Natl. Acad. Sci.* **102**: 4006–4009.
- Ruvkun, G., Wightman, B., and Ha, I. 2004. The 20 years it took to recognize the importance of tiny RNAs. *Cell* **116**: S93–S96.

- Saetrom, P., Heale, B.S., Snove Jr., O., Aagaard, L., Alluin, J., and Rossi, J.J. 2007. Distance constraints between microRNA target sites dictate efficacy and cooperativity. *Nucleic Acids Res.* **35**: 2333–2342. doi: 10.1093/nar/gkml133.
- Sarin, S., O'Meara, M.M., Flowers, E.B., Antonio, C., Poole, R.J., Didiano, D., Johnston Jr., R.J., Chang, S., Narula, S., and Hobert, O. 2007. Genetic screens for *Caenorhabditis elegans* mutants defective in left/right asymmetric neuronal fate specification. *Genetics* **176**: 2109–2130.
- Slack, F.J., Basson, M., Liu, Z., Ambros, V., Horvitz, H.R., and Ruvkun, G. 2000. The *lin-41* RBCC gene acts in the *C. elegans* heterochronic pathway between the *let-7* regulatory RNA and the *LIN-29* transcription factor. *Mol. Cell* **5**: 659–669.
- Stark, A., Brennecke, J., Russell, R.B., and Cohen, S.M. 2003. Identification of *Drosophila* microRNA targets. *PLoS Biol.* **1**: 1–13. doi: 10.1371/journal.pbio.0000060.
- Stefani, G. and Slack, F.J. 2008. Small non-coding RNAs in animal development. *Nat. Rev. Mol. Cell. Biol.* **9**: 219–230.
- Steiner, F.A., Hoogstrate, S.W., Okihara, K.L., Thijssen, K.L., Ketting, R.F., Plasterk, R.H., and Sijen, T. 2007. Structural features of small RNA precursors determine Argonaute loading in *Caenorhabditis elegans*. *Nat. Struct. Mol. Biol.* **14**: 927–933.
- Tomari, Y., Du, T., and Zamore, P.D. 2007. Sorting of *Drosophila* small silencing RNAs. *Cell* **130**: 299–308.
- Vella, M.C., Choi, E.Y., Lin, S.Y., Reinert, K., and Slack, F.J. 2004a. The *C. elegans* microRNA *let-7* binds to imperfect *let-7* complementary sites from the *lin-41* 3' UTR. *Genes & Dev.* **18**: 132–137.
- Vella, M.C., Reinert, K., and Slack, F.J. 2004b. Architecture of a validated microRNA::target interaction. *Chem. Biol.* **11**: 1619–1623.
- Wharton, R.P., Sonoda, J., Lee, T., Patterson, M., and Murata, Y. 1998. The Pumilio RNA-binding domain is also a translational regulator. *Mol. Cell* **1**: 863–872.
- Wightman, B., Ha, I., and Ruvkun, G. 1993. Posttranscriptional regulation of the heterochronic gene *lin-14* by *lin-4* mediates temporal pattern formation in *C. elegans*. *Cell* **75**: 855–862.
- Zhang, L., Ding, L., Cheung, T.H., Dong, M.Q., Chen, J., Sewell, A.K., Liu, X., Yates 3rd, J.R., and Han, M. 2007. Systematic identification of *C. elegans* miRISC proteins, miRNAs, and mRNA targets by their interactions with GW182 proteins AIN-1 and AIN-2. *Mol. Cell* **28**: 598–613.
- Zhao, Y., Samal, E., and Srivastava, D. 2005. Serum response factor regulates a muscle-specific microRNA that targets *Hand2* during cardiogenesis. *Nature* **436**: 214–220.
- Zhao, Y., Ransom, J.F., Li, A., Vedantham, V., von Drehle, M., Muth, A.N., Tsuchihashi, T., McManus, M.T., Schwartz, R.J., and Srivastava, D. 2007. Dysregulation of cardiogenesis, cardiac conduction, and cell cycle in mice lacking *miRNA-1-2*. *Cell* **129**: 303–317.



RNA

A PUBLICATION OF THE RNA SOCIETY

Molecular architecture of a miRNA-regulated 3' UTR

Dominic Didiano and Oliver Hobert

RNA 2008 14: 1297-1317

Supplemental Material <http://rnajournal.cshlp.org/content/suppl/2008/05/07/rna.1082708.DC1>

References This article cites 53 articles, 9 of which can be accessed free at:
<http://rnajournal.cshlp.org/content/14/7/1297.full.html#ref-list-1>

Open Access Freely available online through the *RNA* Open Access option.

License Freely available online through the open access option.

Email Alerting Service Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).

To subscribe to *RNA* go to:
<http://rnajournal.cshlp.org/subscriptions>
