



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 17, 2022 – 07:32 AM JST

PDB ID : 5YXW
Title : Crystal structure of the prefusion form of measles virus fusion protein
Authors : Hashiguchi, T.; Fukuda, Y.; Matsuoka, R.; Kuroda, D.; Kubota, M.; Shirogane, Y.; Watanabe, S.; Tsumoto, K.; Kohda, D.; Plemper, R.K.; Yanagi, Y.
Deposited on : 2017-12-07
Resolution : 2.78 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.27
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.27


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4107 (2.80-2.76)
Clashscore	141614	4575 (2.80-2.76)
Ramachandran outliers	138981	4487 (2.80-2.76)
Sidechain outliers	138945	4489 (2.80-2.76)
RSRZ outliers	127900	4027 (2.80-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	94	
2	B	419	
3	C	2	

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3482 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called glycoprotein F2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	85	662	415	124	119	4	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	19	GLY	-	expression tag	UNP Q786F3

- Molecule 2 is a protein called glycoprotein F1, measles virus fusion protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	367	2771	1749	467	541	14	0	0	0

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	2	28	16	2	10	0	0	0

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	A	1	14	8	1	5	0	0

- Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	2	Total O 2 2	0	0
5	B	5	Total O 5 5	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: glycoprotein F2

Chain A: 



- Molecule 2: glycoprotein F1, measles virus fusion protein

Chain B: 



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C: 



4 Data and refinement statistics

Property	Value	Source
Space group	I 21 3	Depositor
Cell constants a, b, c, α , β , γ	167.39Å 167.39Å 167.39Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	68.34 – 2.78 68.34 – 2.78	Depositor EDS
% Data completeness (in resolution range)	100.0 (68.34-2.78) 100.0 (68.34-2.78)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.77 (at 2.77Å)	Xtrriage
Refinement program	PHENIX (1.11_2567)	Depositor
R, R_{free}	0.202 , 0.249 0.202 , 0.249	Depositor DCC
R_{free} test set	2008 reflections (10.11%)	wwPDB-VP
Wilson B-factor (Å ²)	78.5	Xtrriage
Anisotropy	0.000	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 43.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.021 for -l,-k,-h	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3482	wwPDB-VP
Average B, all atoms (Å ²)	72.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.12% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.58	0/671	0.89	1/910 (0.1%)
2	B	0.51	0/2812	0.75	2/3820 (0.1%)
All	All	0.52	0/3483	0.78	3/4730 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	2

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	48	ARG	NE-CZ-NH2	-7.13	116.73	120.30
2	B	360	ARG	NE-CZ-NH2	-5.19	117.71	120.30
2	B	457	LEU	CB-CG-CD1	-5.14	102.27	111.00

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	291	ILE	Peptide
2	B	370	LEU	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	662	0	696	16	0
2	B	2771	0	2797	36	0
3	C	28	0	25	1	0
4	A	14	0	13	0	0
5	A	2	0	0	1	0
5	B	5	0	0	0	0
All	All	3482	0	3531	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:20:THR:HG23	2:B:436:ARG:HH21	1.41	0.85
1:A:48:ARG:NH1	2:B:264:GLY:O	2.15	0.79
2:B:324:TYR:O	2:B:360:ARG:NH2	2.19	0.75
1:A:92:ASN:O	1:A:96:GLN:HG3	1.87	0.73
1:A:91:LEU:HD23	2:B:225:ILE:HD12	1.75	0.69
2:B:337:MET:HE3	2:B:338:PRO:HD2	1.75	0.68
2:B:212:GLU:HG3	2:B:236:TYR:CG	2.33	0.63
2:B:303:SER:HB3	2:B:375:PHE:CE2	2.34	0.63
2:B:474:LYS:O	2:B:478:GLU:HG3	2.04	0.58
1:A:33:ILE:HB	2:B:313:THR:HG21	1.86	0.56
2:B:398:TYR:HD1	2:B:398:TYR:O	1.89	0.56
2:B:212:GLU:HG3	2:B:236:TYR:CD1	2.43	0.54
5:A:302:HOH:O	2:B:271:HIS:HD2	1.90	0.53
2:B:304:TYR:HB2	2:B:411:ILE:HB	1.90	0.52
2:B:265:ILE:HG23	2:B:283:ILE:HG23	1.92	0.51
1:A:67:ASN:N	1:A:67:ASN:HD22	2.07	0.51
2:B:118:LEU:HD22	2:B:131:ILE:HG23	1.93	0.50
2:B:151:ARG:NH1	2:B:273:ASP:OD1	2.44	0.50
2:B:311:TRP:CZ3	2:B:451:PRO:HB3	2.47	0.50
2:B:333:SER:OG	2:B:345:GLN:HB2	2.12	0.50
1:A:26:HIS:CD2	3:C:1:NAG:H82	2.49	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:69:THR:HG22	2:B:199:GLY:HA2	1.96	0.48
1:A:54:LEU:HD22	2:B:283:ILE:HD11	1.97	0.47
2:B:337:MET:CE	2:B:338:PRO:HD2	2.44	0.47
2:B:398:TYR:HD1	2:B:398:TYR:C	2.18	0.46
2:B:371:VAL:HG13	2:B:375:PHE:HB2	1.97	0.46
2:B:154:LEU:O	2:B:157:THR:HG22	2.17	0.44
1:A:58:LEU:HB2	2:B:277:TYR:O	2.17	0.44
1:A:20:THR:HG22	2:B:383:GLN:NE2	2.33	0.44
1:A:54:LEU:HD12	2:B:173:LEU:HB3	1.99	0.44
2:B:261:GLU:O	2:B:286:PRO:HG2	2.19	0.43
2:B:303:SER:HA	2:B:311:TRP:O	2.18	0.43
2:B:398:TYR:C	2:B:398:TYR:CD1	2.92	0.43
1:A:88:ARG:NH1	2:B:274:THR:OG1	2.52	0.43
2:B:298:ARG:HB2	2:B:318:TYR:CE1	2.54	0.42
2:B:216:LEU:HD21	2:B:234:LEU:HD23	2.01	0.42
1:A:79:LEU:HB3	2:B:210:TYR:CE1	2.54	0.41
1:A:75:GLU:HG3	1:A:78:ARG:HH21	1.86	0.41
2:B:386:LEU:HD12	2:B:423:VAL:HG11	2.03	0.41
1:A:20:THR:HG23	2:B:436:ARG:NH2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	83/94 (88%)	80 (96%)	3 (4%)	0	100	100
2	B	365/419 (87%)	358 (98%)	6 (2%)	1 (0%)	41	70
All	All	448/513 (87%)	438 (98%)	9 (2%)	1 (0%)	47	76

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	371	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	76/83 (92%)	76 (100%)	0	100	100
2	B	305/343 (89%)	303 (99%)	2 (1%)	84	94
All	All	381/426 (89%)	379 (100%)	2 (0%)	88	95

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	143	ASN
2	B	398	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	67	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	C	1	1,3	14,14,15	1.01	1 (7%)	17,19,21	2.65	5 (29%)
3	NAG	C	2	3	14,14,15	1.06	1 (7%)	17,19,21	1.50	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	C	2	3	-	0/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	2	NAG	C7-N2	2.92	1.44	1.34
3	C	1	NAG	C7-N2	2.73	1.43	1.34

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	1	NAG	C1-O5-C5	8.58	123.82	112.19
3	C	1	NAG	C4-C3-C2	-4.26	104.77	111.02
3	C	1	NAG	C2-N2-C7	-2.92	118.75	122.90
3	C	2	NAG	C1-O5-C5	2.80	115.99	112.19
3	C	2	NAG	C8-C7-N2	2.53	120.38	116.10
3	C	1	NAG	O5-C5-C4	2.32	116.47	110.83
3	C	1	NAG	C8-C7-N2	2.27	119.95	116.10
3	C	2	NAG	C4-C3-C2	2.12	114.13	111.02
3	C	2	NAG	O4-C4-C5	2.00	114.27	109.30

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	C	1	NAG	C4-C5-C6-O6
3	C	1	NAG	O5-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	A	203	1	14,14,15	0.98	2 (14%)	17,19,21	2.46	5 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	203	1	-	0/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	203	NAG	C7-N2	2.18	1.41	1.34
4	A	203	NAG	C2-N2	-2.05	1.42	1.46

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	203	NAG	C1-O5-C5	5.65	119.85	112.19
4	A	203	NAG	O5-C1-C2	4.68	118.67	111.29
4	A	203	NAG	C1-C2-N2	-4.64	102.56	110.49
4	A	203	NAG	C8-C7-N2	3.41	121.87	116.10
4	A	203	NAG	C3-C4-C5	-2.17	106.38	110.24

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	85/94 (90%)	0.57	5 (5%) 22 17	54, 69, 104, 122	0
2	B	367/419 (87%)	0.58	21 (5%) 23 18	50, 68, 100, 196	0
All	All	452/513 (88%)	0.58	26 (5%) 23 18	50, 68, 104, 196	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	477	LEU	11.4
2	B	476	LEU	9.4
2	B	481	ASP	8.2
2	B	479	SER	8.1
2	B	480	SER	8.0
2	B	478	GLU	6.6
2	B	473	ALA	6.6
2	B	475	GLU	6.1
1	A	104	VAL	5.8
2	B	470	LEU	5.4
2	B	474	LYS	3.5
2	B	168	GLY	3.4
2	B	472	ASP	3.1
2	B	116	VAL	3.1
1	A	103	SER	3.0
2	B	143	ASN	2.8
1	A	20	THR	2.7
1	A	70	ARG	2.7
2	B	471	GLU	2.5
2	B	147	ILE	2.5
1	A	102	GLN	2.4
2	B	145	GLN	2.3
2	B	375	PHE	2.3
2	B	167	ALA	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	118	LEU	2.1
2	B	198	ILE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q < 0.9
3	NAG	C	2	14/15	0.43	0.35	107,128,135,139	0
3	NAG	C	1	14/15	0.86	0.21	73,93,113,121	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	A	203	14/15	0.95	0.18	78,82,87,90	0

6.5 Other polymers [i](#)

There are no such residues in this entry.