

Management of Children Receiving Antiretroviral Therapy

Updated: June 27, 2024

Reviewed: June 27, 2024

In the United States, most children with HIV are receiving antiretroviral therapy (ART), making treatment-experienced children the norm. Providers may consider antiretroviral (ARV) regimen changes for the following reasons:

- *Treatment simplification:* Modifying ARV regimens in children who are currently receiving effective ART to simplify the regimen
- *Treatment optimization:* Increasing the treatment potency or barrier to resistance of an effective but older or potentially fragile regimen or improving the adverse-event profile
- *Toxicity management:* Recognizing and managing ARV drug toxicity or intolerance (see [Management of Medication Toxicity or Intolerance](#))
- *Treatment failure:* Recognizing and managing treatment failure (see [Recognizing and Managing Antiretroviral Treatment Failure](#))

Modifying Antiretroviral Regimens in Children with Sustained Virologic Suppression on Antiretroviral Therapy

Panel's Recommendations

- Children who have sustained virologic suppression on their current antiretroviral (ARV) regimen should be evaluated regularly for opportunities to change to a new regimen that facilitates adherence, simplifies administration, increases ARV potency or barrier to drug resistance, and decreases the risk of drug-associated toxicity (**AII**).
- Before changing a patient's ARV regimen, clinicians must carefully consider the patient's previous regimens, past episodes of ARV therapy failure, prior drug-resistance test results, drug cost, and insurance coverage, as well as the patient's ability to tolerate the new drug regimen (**AIII**). Archived drug resistance can limit the antiviral activity of a new drug regimen.
- Children should be monitored carefully after a change in treatment. Viral load measurement is recommended 2 to 4 weeks after a change in a child's ARV regimen (**BIII**).

Rating of Recommendations: A = Strong; B = Moderate; C = Optional

Rating of Evidence: I = One or more randomized trials in children[†] with clinical outcomes and/or validated endpoints; I = One or more randomized trials in adults with clinical outcomes and/or validated laboratory endpoints with accompanying data in children[†] from one or more well-designed, nonrandomized trials or observational cohort studies with long-term clinical outcomes; II = One or more well-designed, nonrandomized trials or observational cohort studies in children[†] with long-term outcomes; II* = One or more well-designed, nonrandomized trials or observational studies in adults with long-term clinical outcomes with accompanying data in children[†] from one or more similar nonrandomized trials or cohort studies with clinical outcome data; III = Expert opinion*

[†] Studies that include children or children/adolescents, but not studies limited to postpubertal adolescents

Clinicians choose initial ARV regimens for children with HIV by evaluating the pharmacokinetic (PK), safety, and efficacy data for the drugs that are available in formulations suitable for the child's age and weight at the start of treatment. New ARV drug options may become available as children grow and

learn to swallow pills and as new drugs, drug formulations, and data become available. Even in cases wherein patients have achieved sustained virologic suppression (i.e., suppression for 6–12 months) on their current regimen, clinicians should consider switching patients to new ARV regimens to permit the use of pills instead of liquids; reduce pill burden; allow the use of once-daily medications; reduce the risk of adverse events; minimize drug interactions; and align a child’s regimen with widely used, efficacious adult regimens.¹ These changes often enhance adherence and improve quality of life.²

Treatment Simplification

Many infants and children with HIV initiated treatment with twice-daily dosing (especially prior to the approval of integrase strand transfer inhibitor [INSTI] medications for pediatric use), and regimens included a variety of drug formulations, depending on which formulations were available for a child’s age and weight. Clinicians should regularly review treatment options as children grow, and offer simplified dosing using coformulated drugs and/or once-daily regimens when appropriate (see [Table 18](#) below). Clinicians also should consider a child’s ART history, drug-resistance test results, and ability to swallow tablets. Efforts to increase the availability of coformulated complete ARV regimens have yielded several once-daily options for children that should be considered. The International Maternal Pediatric Adolescent AIDS Clinical Trials (IMPAACT) 2019 study demonstrated safety, efficacy, and appropriate dosing of a fixed-dose combination (FDC) tablet containing abacavir, dolutegravir, and lamivudine (ABC/DTG/3TC) in children aged <12 years, with use of dispersible tablets (Triumeq PD) or an immediate-release tablet (Triumeq) to be swallowed depending on the child’s weight.³ For children weighing ≥14 kg who can swallow pills, additional options include coformulated bictegravir, emtricitabine, and tenofovir alafenamide (BIC/FTC/TAF; Biktarvy) or FTC/TAF (Descovy) plus DTG, which is a two-pill, once-daily regimen. Additional coformulated options are available when children reach 25 kg to 35 kg in weight. See [Table 18](#) below for more information on once-daily options and other coformulated complete ARV regimens. Among treatment-naïve youth in the United States aged 13 to 24 years, some evidence exists that single-tablet regimens (STRs) improve the odds of viral suppression⁴; emerging evidence also supports the safety, efficacy, and tolerability of STRs in younger children.^{5–7} Although these data have not been replicated in treatment-experienced adolescents, clinicians should consider using STRs in children and youth with sustained viral suppression because these regimens reduce pill burden and dosing frequency.

If using an FDC once-daily regimen is not possible, clinicians should determine whether the child’s ARV regimen could be simplified in other ways. For example, small studies have shown that children who achieve virologic suppression using twice-daily dosing for certain ARV drugs (e.g., ABC) maintain virologic suppression when they are switched from twice-daily dosing to once-daily dosing of the same drugs (see the [Abacavir](#) and [Nevirapine](#) sections and FDCs in [Appendix A](#), [Table 1. Antiretrovirals Available in Fixed-Dose Combination Tablets or as a Co-packaged Formulation, by Drug Class](#) and [Appendix A, Table 2. Antiretroviral Fixed-Dose Combination Tablets: Minimum Body Weights and Consideration for Use in Children and Adolescents](#)). However, these studies reported mixed results when switching the dosing for lopinavir/ritonavir (LPV/r) from twice daily to once daily. Therefore, once-daily dosing of LPV/r is not recommended.^{8–11}

Long-acting injectable (LAI) ARV medications may be considered a treatment simplification approach for some virologically suppressed adolescents. The co-packaged, two-drug injectable ARV regimen of cabotegravir and rilpivirine (CAB and RPV; Cabenuva) is approved by the U.S. Food and Drug Administration (FDA) for use in children weighing ≥35 kg and aged ≥12 years, with viral suppression (defined as <50 copies/mL), on a stable ARV regimen, without a history of treatment

failure, and without known or suspected drug resistance to either drug. Studies in adults—such as the First Long-Acting Injectable Regimen (FLAIR) and Antiretroviral Therapy as Long-Acting Suppression (ATLAS) trials—have demonstrated non-inferiority in those receiving monthly CAB and RPV injections compared with adults who stayed on a daily three-drug oral regimen.^{12,13} Similarly, in the ATLAS-2M and SOLAR trials, injections of CAB and RPV every 2 months were found to be non-inferior to monthly injections and once-daily ART, respectively.^{14,15} The IMPAACT 2017 study is currently evaluating CAB and RPV in children aged 12 to 18 years. At 24 weeks of follow-up, injections of CAB and RPV every 2 months maintained viral suppression, showed acceptable PK, and demonstrated an acceptable safety profile in 144 adolescents.¹⁶ Additionally, participating youth and their caregivers reported high acceptability of the treatment and a strong preference for LAI ART over daily oral ART.¹⁷ The Panel on Antiretroviral Therapy and Medical Management of Children Living with HIV (the Panel) notes that questions remain, including whether there are additional adverse effects specific to the pediatric population, whether a two-drug nucleoside-sparing regimen for children with significant ARV treatment history¹⁸ will be effective, and what potential implementation challenges will emerge. A single site in the United States has reported on three adolescents and young adults who experienced viremia while on bimonthly injections which resolved with monthly injections. Two of the individuals also experienced postinjection adverse events that self-resolved.¹⁹ However, given the FDA approval for those as young as 12 years of age, some providers may consider injectable CAB and RPV in adolescents who meet the approved indications and may benefit from a long-acting injectable regimen. See the Cabotegravir and Rilpivirine sections for additional information about these drugs and the dosing and administration of CAB and RPV, and see [Management of the Treatment-Experienced Patient: Optimizing Antiretroviral Therapy in the Setting of Viral Suppression](#) in the Adult and Adolescent ARV Guidelines for practical considerations.

Oral two-drug regimens have some data supporting efficacy in pediatric and adult populations. A two-drug FDC tablet containing DTG/RPV—a nucleoside-sparing, dual-therapy regimen that is marketed as Juluca—is approved by the FDA as a complete regimen to replace the current ARV regimen in adult patients who have been virologically suppressed (HIV RNA <50 copies/mL) on a stable ARV regimen for at least 6 months and who have no history of treatment failure. This approval was based on two Phase 3 clinical trials, SWORD-1 and SWORD-2, in which treatment-experienced adults who were virologically suppressed on three- or four-drug regimens were randomized either to switch to DTG/RPV (early-switch group) or stay on their original regimens through 48 weeks and then switch to DTG/RPV (late-switch group). Results from these trials showed similar rates of virologic suppression in both groups (non-inferiority) through 3 years of follow-up.²⁰⁻²² No equivalent data exist for this drug combination in pediatric patients, although a clinical trial is underway in children aged 6 to 12 years. The Panel usually endorses the use of adult formulations in adolescents, and this product may be appropriate for certain adolescents. DTG/RPV regimens could be useful in patients in whom there is concern for toxicity from nucleoside reverse transcriptase inhibitors (NRTIs). Additionally, findings from the PENTA-17 SMILE study evaluating darunavir/ritonavir (DRV/r) combined with an INSTI, including 318 children aged 6 to 18 years in 11 countries, found that DRV/r plus an INSTI was non-inferior to the standard of care in maintaining virologic suppression at 48 weeks in participants without INSTI or protease inhibitor (PI) resistance.²³ Although the Panel does not recommend this combination for initial treatment, it might be considered in situations in which simplification or avoidance of NRTIs is desired. DTG/3TC (Dovato) also has demonstrated non-inferiority to continuation of three- or four-drug regimens in treatment-experienced adults and those without a history of treatment failure in the TANGO and SALSA studies, respectively.^{24,25} In the ongoing DANCE study, DTG/3TC is being evaluated as an initial regimen in ART-naïve adolescents aged 12 to <18 years and weighing ≥25 kg.

and with HIV RNA of 1,000 copies/mL to ≤500,000 copies/mL. Safety and efficacy of DTG/3TC were comparable to adults, and 22 of 32 participants achieved viral suppression at 96 weeks.²⁶ Based on these findings, the FDA has approved DTG/3TC in adolescents aged ≥12 years and weighing ≥25 kg as an initial regimen or for those on a stable ART regimen with no history of treatment failure and no known drug resistance to the individual drugs. The Panel notes that adolescents may have difficulties adhering to therapy and recommends close monitoring with viral load testing in anyone on oral two-drug regimens. The Panel does not recommend two-drug regimens for initial ART in children (see [What to Start](#)).

Treatment Optimization

The aims of treatment optimization may include improving the potency of the regimen, improving a child's growth or other health outcomes through reduced drug side effects and/or better treated HIV, or maximizing palatability. More studies are directly evaluating treatment optimization in children, and early results support the safety and efficacy of regimen switches for those with viral suppression. Older studies have demonstrated sustained viral suppression and improved growth outcomes in young children who have demonstrated good adherence and no baseline resistance and who were switched from LPV/r-based regimens to an efavirenz (EFV)-based regimen (NEVEREST 3).²⁷⁻²⁹ Replacing LPV/r with Efv may provide some benefits (e.g., once-daily dosing and a different side-effect profile), but most pediatric HIV experts would prefer replacing LPV/r with an equally potent PI (e.g., darunavir [DRV] or atazanavir [ATV]) or an INSTI (e.g., elvitegravir [EVG], raltegravir, DTG, or BIC), based on studies in adults and emerging evidence of non-inferiority or superiority in children.^{30,31} Although not a switch trial, findings from the randomized controlled Once-daily DTG-based ART in Young people vS. Standard thErApY (ODYSSEY) study of more than 700 children aged <18 years in eight countries initiating DTG as first- or second-line therapy showed superior virologic and clinical outcomes in children randomized to optimization with DTG-based ART compared with those in the standard of care (PI- or non-nucleoside reverse transcriptase inhibitor [NNRTI]-based regimens), contributing to evidence supporting optimization with DTG-based regimens.³² Results from the younger ODYSSEY cohort of children weighing between 3 kg and 14 kg also showed superiority of DTG-based ART compared with other regimens, more than 70% of which were PI-based regimens.³³ Additionally, several observational studies in sub-Saharan Africa that are evaluating efforts to optimize pediatric ARV regimens have shown improved viral suppression rates in children that were switched to DTG-based regimens.³⁴⁻³⁶ Similarly, a retrospective study from six African countries reporting on 7,898 children and adolescents aged 0 to ≤19 years demonstrated that 93% remained virologically suppressed after switching from NNRTI- and PI-based regimens to DTG-based regimens, and nearly 80% of those previously unsuppressed achieved viral suppression while on DTG.³⁷ The INSTI-based FDC regimen BIC/FTC/TAF also has shown efficacy and high rates of long-term viral suppression in adolescents and children ≥2 years and weighing 14 kg to <25 kg.^{38,39} Similarly, EVG/cobicistat/FTC/TAF has shown efficacy in adolescents. Early results from small, randomized studies also show potential for switches to newer-generation NNRTI medications—such as RPV⁴⁰ and doravirine⁴¹—in children and adolescents weighing ≥35 kg who have been virologically suppressed on a stable ARV regimen.

Toxicity Management

Several studies of small cohorts of children have demonstrated sustained virologic suppression and reassuring safety outcomes when drugs that have greater long-term toxicity risks are replaced with drugs that are thought to have lower toxicity risks (e.g., replacing stavudine with tenofovir disoproxil fumarate (TDF), TAF, zidovudine, or ABC; replacing PIs with NNRTIs), including improved lipid

profiles.⁴²⁻⁴⁶ Similarly, adolescents who were switched from EFV to RPV, a newer generation of NNRTIs, showed similar rates of viral suppression with improved metabolic profiles and cognitive outcomes.⁴⁰ Additionally, studies in adults have shown improved tolerability, lipid profiles, and insulin sensitivity in patients who were switched from PIs to INSTIs,⁴⁷⁻⁵¹ and adults who were switched from EFV to an INSTI have shown improvement in neuropsychiatric symptoms. One study in South Africa showed that prevalence of hepatic steatosis decreased from 17% to 3% among 30 adolescents who switched to a DTG-containing regimen but increased from 8% to 16% among 38 adolescents who continued a non-DTG-containing regimen. Additionally, cholesterol and triglycerides were lower in those who switched to DTG and in whom no excess weight gain was observed.⁵² In other studies, however, the use of INSTIs, as well as TAF, has been associated with weight gain in adults and adolescents, with emerging data showing an association in children.⁵³⁻⁵⁷ Finally, NRTI-sparing regimens, including the dual-drug oral regimens (DRV and an INSTI or DTG/RPV) and the approved long-acting injectable regimen (CAB with RPV) described above, may be considered in patients with NRTI toxicity who otherwise are eligible for these complete ARV regimens. In a small subgroup analysis of the SWORD study, participants switched to DTG/RPV experienced small but statistically significant improvement in bone mineral density and bone turnover markers compared with those who continued on TDF.⁵⁸ Of note, however, is that, although small in number, more participant adverse events that led to discontinuation were reported in the DTG/RPV arm (3%) than in the arm in which participants stayed on their current regimen (<1%).²⁰

Treatment Failure

Treatment failure is another common reason providers change ARV regimens in children with HIV. This topic is covered in [Recognizing and Managing Antiretroviral Treatment Failure](#).

Regimens That Are Not Recommended for Use in Children

Monotherapy PI regimens (DRV/r, LPV/r, atazanavir/ritonavir)^{59,60} and monotherapy regimens of DTG^{61,62} have been used to simplify or reduce the toxicity of regimens in adult patients who have sustained virologic suppression, but with varying success. These strategies are still being explored, but they are not currently recommended as management strategies in children because of the lack of data.^{60,63-66}

Potential Antiretroviral Drug Switches in Children with Virologic Suppression

Table 18 below contains examples of potential ARV drug changes in children with sustained virologic suppression on their current regimen for the purpose of treatment simplification, optimization, or reduced toxicity. When considering such a change, a clinician should first ensure that a recent viral load test indicates that the child is not experiencing virologic failure and that the child has a reliable history of good adherence (assessed by self and parental report, pharmacy refill, prior viral loads, etc.). Clinicians also must consider ART history, tolerability, ability to swallow tablets, and all prior drug-resistance test results to avoid choosing new ARV drugs for which archived drug resistance would reemerge and limit the activity of the regimen.⁶⁷⁻⁷¹ The evidence that supports many of these ARV changes is indirect (i.e., extrapolated from data about drug performance during initial therapy or follow-up therapy after treatment failure). When such changes are made, careful monitoring (e.g., taking a viral load measurement 2–4 weeks after making the switch to the new regimen) is important to ensure that virologic suppression is maintained.

Table 18. Examples of Changes in Antiretroviral Regimen Components for Children with Sustained Virologic Suppression

This list is not exhaustive and does not necessarily contain all potential treatment options. Instead, it provides examples of changes that could be made. The table includes information only about switching between ARV drugs; **it does not include all the information that clinicians should consider before prescribing these drugs, such as drug cost and the patient's insurance coverage.** Refer to the individual drug sections; [Appendix A, Table 1. Antiretrovirals Available in Fixed-Dose Combination Tablets or as a Co-Packaged Formulation, by Drug Class](#); and [Appendix A, Table 2. Antiretroviral Fixed-Dose Combination Tablets and Co-packaged Formulations: Minimum Body Weights and Considerations for Use in Children and Adolescents](#) in [Appendix A. Pediatric Antiretroviral Drug Information](#) for further information about the use and administration of specific ARV drugs and FDC formulations.

For images of most of the ARV drugs listed in this table, see the [Antiretroviral Medications](#) section of the National HIV Curriculum. In addition, a resource from the United Kingdom illustrates the relative sizes of individual ARV drug FDC tablets (see the [ARV Chart in HIV i-Base](#)). Although most of the drugs listed in that chart are the same as those in the United States, not all formulations available in the United States are included, and there are differences in a few of the brand names.

Current ARV Drug(s)	Age, Weight, and Sexual Maturity Rating Requirements	Potential ARV Drug Switch ^a	Comment
NRTIs			
ABC Twice Daily	Aged ≥3 months ^b	ABC once daily	See the Abacavir^b section for more information.
3TC Twice Daily	Aged ≥3 years	3TC once daily	See the Lamivudine section for more information.
	Any age (starting at full-term birth) Any weight	FTC once daily	See the Emtricitabine section for more information.
ZDV	Aged ≥1 months ^b	ABC	Less long-term mitochondrial toxicity Children aged ≥3 months can take ABC once daily.
	Weighing 17 kg to <25 kg	TDF	TDF is a reasonable, once-daily option for HLA-B*5701-positive children for whom ABC is not recommended and in whom ZDV is not tolerated. TDF is available as an oral powder and as low-strength tablets alone or in combination with FTC.
	Weighing ≥14 kg	TAF ^c	Less long-term mitochondrial toxicity. Once-daily dosing. Only available in coformulation with other ARV drugs; can further reduce pill burden. TAF is preferred over TDF because of the lower risk of bone and renal toxicity, but it may be associated with weight gain and lipid abnormalities.

Table 18. Examples of Changes in Antiretroviral Regimen Components for Children with Sustained Virologic Suppression

Current ARV Drug(s)	Age, Weight, and Sexual Maturity Rating Requirements	Potential ARV Drug Switch ^a	Comment
	Weighing ≥ 14 kg	FTC/TAF ^c (Descovy)	Once-daily dosing. This combination NRTI medication may be more desirable because of smaller pill size and reduced pill burden. Benefits as described for TAF.
Any NRTI	Aged ≥ 12 years Weighing ≥ 35 kg	CAB and RPV co-packaged regimen as Cabenuva	NRTI-sparing regimen. Long-acting injectable, complete ARV regimen requiring two IM injections every 1 to 2 months that together are an alternative to daily oral ARV regimens. Must consider prior history of treatment failure and known or suspected drug resistance to individual drugs. Injection site reactions are common but do not often result in discontinuation of the regimen. See the Cabotegravir section for more information.
	Aged ≥ 12 years Weighing ≥ 35 kg	DTG/RPV (Juluca)	NRTI-sparing FDC that is a complete regimen. In addition to age and weight criteria (based on RPV component because DTG was approved for younger ages/lower weights), must be virologically suppressed (HIV RNA <50 copies/mL) on a stable ARV regimen for at least 6 months and without history of treatment failure. Should be taken with food. No pediatric data.
NNRTIs			
NVP or EFV	Any age (starting at full-term birth) Weighing ≥ 2 kg	RAL ^d	RAL is preferred over NVP in infants from birth to age 4 weeks who weigh ≥ 2 kg. Both are dosed twice daily in children. Note that DTG and BIC have a higher barrier to resistance than RAL. In a child >1 month of age, DTG is preferred. See DTG below.
	Age ≥ 4 weeks Weighing ≥ 3 kg	DTG	DTG is available as a single drug in dispersible and film-coated tablet formulations, or as part of an FDC tablet, all of which can be dosed once daily if no documented resistance or history of failure with INSTI agents exists. DTG plus FTC/TAF (Descovy) in patients weighing at least 14 kg or the weight-appropriate dose of FTC/TDF (Truvada) can be used in children weighing 20 kg to <25 kg. DTG is available as a component of the FDC ABC/DTG/3TC, which is a complete ARV regimen that can be given to infants and children aged ≥ 3 months and weighing ≥ 6 kg to <25 kg in dispersible tablets (Triumeq PD) and to children and adolescents weighing ≥ 25 kg in a single tablet to be swallowed (Triumeq). Higher barrier to resistance, which makes it a good choice for patients who have poor adherence. May improve lipid levels. See the Dolutegravir section for more information.
	Aged ≥ 3 months Weighing ≥ 6 kg to <25 kg	ABC/DTG/3TC (Triumeq PD)	Once-daily dosing. Dispersible tablets with dosage for use in children based on weight. Aligns a child's regimen with an efficacious regimen that is used in adults. See the Dolutegravir section for more information.

Table 18. Examples of Changes in Antiretroviral Regimen Components for Children with Sustained Virologic Suppression

Current ARV Drug(s)	Age, Weight, and Sexual Maturity Rating Requirements	Potential ARV Drug Switch ^a	Comment
	Aged ≥3 months Weighing ≥5 kg	ATV/r	ATV/r has a potentially greater barrier to resistance; however, taking ATV/r may be difficult for some patients, as ATV oral powder must be mixed with food or a beverage before administration, and the palatability of the RTV oral solution is poor.
	Aged ≥3 years Weighing ≥10 kg	DRV/r	DRV/r has a potentially greater barrier to resistance. DRV/r is administered twice daily to patients aged <12 years but may be administered once daily in children aged ≥12 years who do not have any DRV resistance mutations. Note that the palatability of the RTV oral solution is poor when considering administering to children not able to swallow tablets.
	Weighing ≥14 kg	BIC as Biktarvy	Once-daily dosing. BIC is available as a component of the FDC tablet BIC/FTC/TAF (Biktarvy) in two weight-based dose formulations—one formulation for those ≥14 kg to <25 kg and another for those ≥25 kg. This is a complete ARV regimen that can be taken with or without food.
	Weighing ≥25 kg	EVG as Genvoya	EVG is available as a component of the FDC tablet EVG/c/FTC/TAF (Genvoya), which is a complete ARV regimen that must be taken with food.
	Weighing ≥35 kg	DOR	DOR is available in a once-daily FDC tablet DOR/3TC/TDF (Delstrigo). Fewer side effects than reported with EFV. It has continued activity in the setting of some NNRTI mutations.
	Aged ≥12 years Weighing ≥35 kg	CAB and RPV co-packaged regimen as Cabenuva	Long-acting injectable, complete ARV regimen requiring two IM injections every 1 to 2 months that together are an alternative to daily oral ARV regimens. Must consider prior history of treatment failure and known or suspected drug resistance to individual drugs. Injection site reactions are common but do not often result in discontinuation of the regimen. See the Cabotegravir section for more information.
	Aged ≥12 years Weighing ≥35 kg	RPV	Lower incidence of adverse lipid effects. May have fewer sleep disturbances and neuropsychiatric symptoms compared to EFV. RPV has continued activity in the setting of some NNRTI mutations.
PIs			
LPV/r Twice Daily	Any age (starting at full-term birth) Weighing ≥2 kg	RAL ^d	Better palatability. RAL HD can only be given once daily in those weighing ≥40 kg. Unlike LPV/r, the use of RAL is not restricted to infants with a corrected gestational age of ≥42 weeks and a postnatal age of ≥14 days. RAL granules may be difficult to dose for some caregivers.

Table 18. Examples of Changes in Antiretroviral Regimen Components for Children with Sustained Virologic Suppression

Current ARV Drug(s)	Age, Weight, and Sexual Maturity Rating Requirements	Potential ARV Drug Switch ^a	Comment
	Age \geq 4 weeks Weighing \geq 3 kg	DTG	Once-daily dosing if no documented resistance or history of failure with INSTI agents exists. May be better tolerated, and it can be given as a dispersible tablet in young children. DTG is available as a component of the FDC ABC/DTG/3TC, which is a complete ARV regimen that can be given to infants and children aged \geq 3 months and weighing \geq 6 kg to <25 kg in dispersible tablets (Triumeq PD) and to children and adolescents weighing \geq 25 kg in a single tablet to be swallowed (Triumeq). DTG plus FTC/TAF (Descovy) in those weighing at least 14 kg, or the weight-appropriate dose of FTC/TDF (Truvada) can be used in children weighing 20 kg to <25 kg. May improve lipid levels. See the Dolutegravir section for more information.
	Aged \geq 3 months Weighing \geq 6 kg to <25 kg	ABC/DTG/3TC (Triumeq PD)	Once-daily dosing. Dispersible tablets with dosage for use in children based on weight. Aligns a child's regimen with an efficacious regimen that is used in adults. See the Dolutegravir section for more information.
	Aged \geq 3 years Weighing \geq 10 kg	EFV	Once-daily dosing. Better palatability. Lower incidence of adverse lipid effects. Review NNRTI mutations before use. See the Efavirenz section for concerns about Efavirenz dosing for children aged <3 years.
	Aged \geq 3 months Weighing \geq 5 kg	ATV/r	Once-daily dosing. ATV/r may have a lower incidence of adverse lipid effects; however, taking ATV/r may be difficult for some patients, as ATV oral powder must be mixed with food or a beverage before administration, and the palatability of the RTV oral solution is poor.
	Aged \geq 3 years Weighing \geq 10 kg	DRV/r	DRV/r may have a lower incidence of adverse lipid effects. DRV/r is administered twice daily to patients aged <12 years, but it may be administered once daily in children aged \geq 12 years who do not have DRV resistance mutations. Note that palatability of the RTV oral solution is poor when considering administering to children not able to swallow tablets.
	Weighing \geq 14 kg	BIC as Biktarvy	Once-daily dosing. BIC is available as a component of the FDC tablet BIC/FTC/TAF (Biktarvy) in two weight-based dose formulations—one for those \geq 14 kg to <25 kg and another for those \geq 25 kg. This is a complete ARV regimen that can be taken with or without food.
	Weighing \geq 25 kg	EVG as Genvoya	EVG is available as a component of the FDC tablet EVG/c/FTC/TAF (Genvoya), which is a complete ARV regimen that must be taken with food.

Table 18. Examples of Changes in Antiretroviral Regimen Components for Children with Sustained Virologic Suppression

Current ARV Drug(s)	Age, Weight, and Sexual Maturity Rating Requirements	Potential ARV Drug Switch ^a	Comment
	Weighing ≥ 35 kg	DOR	DOR is available in a once-daily FDC tablet DOR/3TC/TDF (Delstrigo). Fewer side effects than reported with EFV. It has continued activity in the setting of some NNRTI mutations.
	Aged ≥ 12 years Weighing ≥ 35 kg	CAB and RPV co-packaged regimen as Cabenuva	Long-acting injectable, complete ARV regimen requiring two IM injections every 1 to 2 months that together are an alternative to daily oral ARV regimens. Must consider prior history of treatment failure and known or suspected drug resistance to individual drugs. Injection site reactions are common but do not often result in discontinuation of the regimen. See the Cabotegravir section for more information.
	Aged ≥ 12 years Weighing ≥ 35 kg	RPV	May be better tolerated. Lower incidence of adverse lipid effects. It has continued activity in the setting of some NNRTI mutations.
INSTIs			
RAL	Age >1 month and weighing <14 kg Weighing ≥ 14 kg	DTG DTG or BIC	Once-daily dosing. Higher barrier to resistance. DTG is available as a single drug in a dispersible tablet for infants and children weighing ≥ 3 kg; in a dispersible FDC for children weighing ≥ 6 kg to 25 kg; in a single-drug film-coated tablet for children weighing ≥ 14 kg; or as an FDC tablet. All of these can be dosed once daily if no documented resistance or history of failure with INSTI agents exists. DTG plus FTC/TAF (Descovy) in those weighing at least 14 kg or the weight-appropriate dose of FTC/TDF (Truvada) can be used in children weighing 20 kg to <25 kg. DTG is available as a component of the FDC ABC/DTG/3TC, which is a complete ARV regimen that can be given to infants and children aged ≥ 3 months and weighing ≥ 6 kg to <25 kg in dispersible tablets (Triumeq PD) and to children and adolescents weighing ≥ 25 kg in a single tablet to be swallowed (Triumeq). See the Dolutegravir section for more information. BIC has once-daily dosing and a higher barrier to resistance. BIC is available as a component of the FDC tablet BIC/FTC/TAF (Biktarvy) in two weight-based dose formulations—one for those ≥ 14 kg to <25 kg and another for those ≥ 25 kg. This is a complete ARV regimen that can be taken with or without food.
	Aged ≥ 12 years Weighing ≥ 35 kg	CAB and RPV co-packaged regimen as Cabenuva	Long-acting injectable, complete ARV regimen requiring two IM injections every 1 to 2 months that together are an alternative to daily oral ARV regimens. Must consider prior history of treatment failure and known or suspected drug resistance to individual drugs. Injection site reactions are common but do not often result in discontinuation of the regimen. See the Cabotegravir section for more information.

Table 18. Examples of Changes in Antiretroviral Regimen Components for Children with Sustained Virologic Suppression

Current ARV Drug(s)	Age, Weight, and Sexual Maturity Rating Requirements	Potential ARV Drug Switch ^a	Comment
EVG/c	Weighing ≥ 14 kg	DTG or BIC	<p>Once-daily dosing. Higher barrier to resistance. DTG is available as a single drug in a dispersible tablet for infants and children weighing ≥ 3 kg; in a dispersible FDC for children weighing ≥ 6 kg to < 25 kg; in a single-drug film-coated tablet for children weighing ≥ 14 kg; or as an FDC tablet. All of these can be dosed once daily if no documented resistance or history of failure with INSTI agents exists. DTG plus FTC/TAF (Descovy) in those weighing at least 14 kg or the weight-appropriate dose of FTC/TDF (Truvada) can be used in children weighing 20 kg to < 25 kg. DTG is available as a component of the FDC ABC/DTG/3TC, which is a complete ARV regimen that can be given to infants and children aged ≥ 3 months and weighing ≥ 6 kg to < 25 kg in dispersible tablets (Triumeq PD) and to children and adolescents weighing ≥ 25 kg in a single tablet to be swallowed (Triumeq). See the Dolutegravir section for more information.</p> <p>BIC has once-daily dosing and a higher barrier to resistance. BIC is available as a component of the FDC tablet BIC/FTC/TAF (Biktarvy) in two weight-based dose formulations—one for those ≥ 14 kg to < 25 kg and another for those ≥ 25 kg. This is a complete ARV regimen that can be taken with or without food.</p>
	Aged ≥ 12 years Weighing ≥ 35 kg	CAB and RPV co-packaged regimen as Cabenuva	Long-acting injectable, complete ARV regimen requiring two IM injections every 1 to 2 months that together are an alternative to daily oral ARV regimens. Must consider prior history of treatment failure and known or suspected drug resistance to individual drugs. Injection site reactions are common but do not often result in discontinuation of the regimen. See the Cabotegravir section for more information.
Other			
Any Multi-Pill and/or Twice-Daily Regimen	Aged ≥ 3 months Weighing ≥ 6 kg to < 25 kg	ABC/DTG/3TC (Triumeq PD)	Once-daily dosing. Dispersible tablets with dosage for use in children based on weight. Aligns a child's regimen with an efficacious regimen that is used in adults. See the Dolutegravir section for more information.
	Weighing ≥ 14 kg	FTC/TAF ^c (Descovy) plus DTG	Once-daily dosing. This regimen may be more desirable because of smaller pill sizes, but it has a higher pill burden (two pills instead of one). Aligns a child's regimen with an efficacious regimen that is used in adults. See the Dolutegravir section for more information.
	Weighing ≥ 14 kg	BIC/FTC/TAF (Biktarvy)	Once-daily dosing. Single pill that can be taken with or without food. Available in two weight-based dose formulations—one for those ≥ 14 kg to < 25 kg and another for those ≥ 25 kg.

Table 18. Examples of Changes in Antiretroviral Regimen Components for Children with Sustained Virologic Suppression

Current ARV Drug(s)	Age, Weight, and Sexual Maturity Rating Requirements	Potential ARV Drug Switch ^a	Comment
	Weighing ≥ 25 kg	ABC/DTG/3TC (Triumeq)	Once-daily dosing. Single pill to be swallowed. Aligns a child's regimen with an efficacious regimen that is used in adults. Large pill size may be a deterrent. See the Dolutegravir section for more information.
	Weighing ≥ 25 kg	EVG/c/FTC/TAF (Genvoya)	Once-daily dosing. Single pill. Alignment with adult ARV regimens. Must be taken with food.
	Weighing ≥ 35 kg	DOR/3TC/TDF (Delstrigo)	Once-daily dosing. Single pill. Aligns a child's regimen with an efficacious regimen that is used in adults. Must be taken with food at a consistent time daily. Renal and bone toxicity of TDF limit its use. Review NNRTI mutations and check for drug-drug interactions before use.
	Weighing ≥ 35 kg SMR 4 or 5	EVG/c/FTC/TDF (Stribild)	Once-daily dosing. Single pill. Aligns a child's regimen with an efficacious regimen that is used in adults. Must be taken with food. Renal and bone toxicity of TDF limit its use.
	Aged ≥ 12 years Weighing ≥ 35 kg	CAB and RPV co-packaged regimen as Cabenuva	Long-acting injectable, complete ARV regimen requiring two IM injections every 1 to 2 months that together are an alternative to daily oral ARV regimens. Must consider prior history of treatment failure and known or suspected drug resistance to individual drugs. Injection site reactions are common but do not often result in discontinuation of the regimen. See the Cabotegravir section for more information.
	Aged ≥ 12 years Weighing ≥ 35 kg	FTC/RPV/TAF (Odefsey)	Once-daily dosing. Single pill. Aligns a child's regimen with an efficacious regimen that is used in adults. Review NNRTI mutations and check for drug-drug interactions before use. Must be taken with food at a consistent time daily.
	Aged ≥ 12 years Weighing ≥ 35 kg SMR 4 or 5	FTC/RPV/TDF (Complera)	Once-daily dosing. Single pill. Aligns a child's regimen with an efficacious regimen that is used in adults. Review NNRTI mutations and check for drug-drug interactions before use. Must be taken with food at a consistent time daily. Renal and bone toxicity of TDF limit its use.
	Aged ≥ 12 years Weighing ≥ 35 kg	DTG/RPV (Juluca)	NRTI-sparing FDC that is a complete regimen. In addition to age and weight criteria (based on RPV component because DTG is approved for younger ages/lower weights), must be virologically suppressed (HIV RNA <50 copies/mL) on a stable ARV regimen for at least 6 months and without history of treatment failure. Should be taken with food. No pediatric data.

Table 18. Examples of Changes in Antiretroviral Regimen Components for Children with Sustained Virologic Suppression

Current ARV Drug(s)	Age, Weight, and Sexual Maturity Rating Requirements	Potential ARV Drug Switch ^a	Comment
	Aged ≥12 years Weighing ≥25 kg	DTG/3TC (Dovato)	Once-daily, two-drug complete regimen approved in adolescents and adults with no known mutations associated with resistance to the individual components who are either ART-naïve or who are virologically suppressed on a stable ART regimen with no history of treatment failure. Because adolescents may have difficulties adhering to therapy, close monitoring with viral load testing is recommended.

^a The possibility of planned and unplanned pregnancy should be considered when selecting an ART regimen for an adolescent. When discussing ART options with adolescents of childbearing potential and their caregivers, it is important to consider the benefits and risks of all ARV drugs and to provide the information and counseling needed to support informed decision-making; refer to the Perinatal Guidelines (see [Recommendations for Use of Antiretroviral Drugs During Pregnancy](#), [Table 7. Situation-Specific Recommendations for Use of Antiretroviral Drugs in Pregnant People and Nonpregnant People Who Are Trying to Conceive](#), and [Appendix C. Antiretroviral Counseling Guide for Health Care Providers](#)).

^b For infants and young children who are being treated with liquid formulations of ABC, initiation with once-daily ABC is not generally recommended. In clinically stable patients with undetectable viral loads who have had stable CD4 T lymphocyte cell counts on twice-daily ABC, the dose can be changed from twice daily to once daily **in those aged ≥3 months**. ABC is not approved by the U.S. Food and Drug Administration for use in neonates and infants aged <3 months. Data from the [IMPAACT P1106 trial](#) and two observational cohorts provide reassuring evidence of the safety of ABC in infants aged <3 months. Based on these data, clinicians may consider the use of **twice daily** ABC in infants aged ≥1 month to <3 months, in consultation with a pediatric HIV specialist (see the [Abacavir](#) section for more information).

^c For children and adolescents weighing ≥14 kg to <35 kg, TAF can be used in combination with an INSTI or an NNRTI, but **not** a boosted PI. For children and adolescents weighing ≥35 kg, TAF can be used in combination with an INSTI, NNRTI, or boosted PI.

^d RAL is recommended for twice-daily use in children. Chewable tablets can be used as dispersible tablets starting at 4 weeks of age. RAL HD once daily is **only recommended** for virologically suppressed children weighing ≥40 kg.

Key: 3TC = lamivudine; ABC = abacavir; ARV = antiretroviral; ATV = atazanavir; ATV/r = atazanavir/ritonavir; BIC = bictegravir; CAB = cabotegravir; DOR = doravirine; DRV = darunavir; DRV/r = darunavir/ritonavir; DTG = dolutegravir; EFV = efavirenz; EVG = elvitegravir; EVG/c = elvitegravir/cobicistat; FDC = fixed-dose combination; FTC = emtricitabine; HD = high dose; HLA = human leukocyte antigen; IM = intramuscular; INSTI = integrase strand transfer inhibitor; LPV/r = lopinavir/ritonavir; NNRTI = non-nucleoside reverse transcriptase inhibitor; NRTI = nucleoside reverse transcriptase inhibitor; NVP = nevirapine; PI = protease inhibitor; RAL = raltegravir; RPV = rilpivirine; RTV = ritonavir; SMR = sexual maturity rating; TAF = tenofovir alafenamide; TDF = tenofovir disoproxil fumarate; ZDV = zidovudine

References

1. Hsu AJ, Neptune A, Adams C, et al. Antiretroviral stewardship in a pediatric HIV clinic: development, implementation and improved clinical outcomes. *Pediatr Infect Dis J.* 2016;35(6):642-648. Available at: <https://pubmed.ncbi.nlm.nih.gov/26906161>.
2. Maiese EM, Johnson PT, Bancroft T, et al. Quality of life of HIV-infected patients who switch antiretroviral medication due to side effects or other reasons. *Curr Med Res Opin.* 2016;32(12):2039-2046. Available at: <https://pubmed.ncbi.nlm.nih.gov/27552553>.
3. Brooks KM, Kiser JJ, Ziembka L, et al. Pharmacokinetics, safety, and tolerability of dispersible and immediate-release abacavir, dolutegravir, and lamivudine tablets in children with HIV (IMPAACT 2019): week 24 results of an open-label, multicentre, phase 1-2 dose-confirmation study. *Lancet HIV.* 2023;10(8):e506-e517. Available at: <https://pubmed.ncbi.nlm.nih.gov/37541705>.
4. Griffith DC, Farmer C, Gebo KA, et al. Uptake and virological outcomes of single- versus multi-tablet antiretroviral regimens among treatment-naive youth in the HIV Research Network. *HIV Med.* 2019;20(2):169-174. Available at: <https://pubmed.ncbi.nlm.nih.gov/30561888>.
5. Natukunda E, Gaur A, Kosalaraksa P, et al. Safety, efficacy, and pharmacokinetics of single-tablet elvitegravir, cobicistat, emtricitabine, and tenofovir alafenamide in virologically suppressed, HIV-infected children: a single-arm, open-label trial. *Lancet Child Adolescent Health.* 2017;1(1):27-34. Available at: <http://www.sciencedirect.com/science/article/pii/S2352464217300093?via%3Dhub>.
6. Liberty A, Strehlau R, Rakhmanina N, et al. Acceptability & palatability of low dose B/F/TAF & E/C/F/TAF in children (≥ 2 y) with HIV. Presented at: International Workshop on HIV & Pediatrics 2020; 2020. Virtual. Available at: <https://academicmedicaleducation.com/meeting/international-workshop-hiv-pediatrics-2020/abstract/acceptability-palatability-low-dose>.
7. Rotsaert A, Nostlinger C, Collin O, et al. Acceptability of a new 4-in-1 abacavir/lamivudine/lopinavir/ritonavir paediatric fixed-dose combination: the caregiver-child dyad's perspective. Presented at: International Workshop on HIV & Pediatrics 2020; 2020. Virtual Available at: <https://academicmedicaleducation.com/meeting/international-workshop-hiv-pediatrics-2020/abstract/acceptability-new-4-1>.
8. Foissac F, Blanche S, Dollfus C, et al. Population pharmacokinetics of atazanavir/ritonavir in HIV-1-infected children and adolescents. *Br J Clin Pharmacol.* 2011;72(6):940-947. Available at: <https://pubmed.ncbi.nlm.nih.gov/21649692>.
9. Chokephaibulkit K, Prasitsuebsai W, Wittawatmongkol O, et al. Pharmacokinetics of darunavir/ritonavir in Asian HIV-1-infected children aged ≥ 7 years. *Antivir Ther.* 2012;17(7):1263-1269. Available at: <https://pubmed.ncbi.nlm.nih.gov/22954687>.
10. Paediatric European Network for Treatment of AIDS. Once vs. twice-daily lopinavir/ritonavir in HIV-1-infected children. *AIDS.* 2015;29(18):2447-2457. Available at: <https://pubmed.ncbi.nlm.nih.gov/26558544>.

11. Gondrie IPE, Bastiaans DET, Fraaij PLA, et al. Sustained viral suppression in HIV-infected children on once-daily lopinavir/ritonavir in clinical practice. *Pediatr Infect Dis J*. 2017;36(10):976-980. Available at: <https://pubmed.ncbi.nlm.nih.gov/28475554>.
12. Orkin C, Bernal Morell E, Tan DHS, et al. Initiation of long-acting cabotegravir plus rilpivirine as direct-to-injection or with an oral lead-in in adults with HIV-1 infection: week 124 results of the open-label Phase 3 FLAIR study. *Lancet HIV*. 2021;8(11):e668-e678. Available at: <https://pubmed.ncbi.nlm.nih.gov/34656207>.
13. Lowenthal E, Chapman J, Calabrese K, et al. Adolescent and parent experiences with long-acting injectables in the MOCHA study. Presented at: Conference on Retroviruses and Opportunistic Infections; 2022. Virtual. Available at: <https://www.croiconference.org/abstract/adolescent-and-parent-experiences-with-long-acting-injectables-in-the-mocha-study>.
14. Jaeger H, Overton ET, Richmond G, et al. Long-acting cabotegravir and rilpivirine dosed every 2 months in adults with HIV-1 infection (ATLAS-2M), 96-week results: a randomised, multicentre, open-label, Phase 3b, non-inferiority study. *Lancet HIV*. 2021;8(11):e679-e689. Available at: <https://pubmed.ncbi.nlm.nih.gov/34648734>.
15. Ramgopal MN, Castagna A, Cazanave C, et al. Efficacy, safety, and tolerability of switching to long-acting cabotegravir plus rilpivirine versus continuing fixed-dose bictegravir, emtricitabine, and tenofovir alafenamide in virologically suppressed adults with HIV, 12-month results (SOLAR): a randomised, open-label, phase 3b, non-inferiority trial. *Lancet HIV*. 2023;10(9):e566-e577. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37567205>.
16. Gaur A, Capparelli E, Baltrusaitis K, et al. Long-acting cabotegravir plus rilpivirine in adolescents with HIV: week 24 IMPAACT 2017(MOCHA) study. Presented at: Conferences on Retroviruses and Opportunistic Infections 2024. Denver, CO. Available at: <https://www.croiconference.org/abstract/long-acting-cabotegravir-plus-rilpivirine-in-adolescents-with-hiv-week-24-impaact-2017mocha-study>.
17. Lowenthal ED, Chapman J, Vaca MZ, et al. IMPACCT 2017 Adolescent/Parent experiences with LA Cabotegravir plus Rilpivirine for HIV treatment. Presented at: Conference on Retroviruses and Opportunistic Infections; 2024. Denver, Colorado. . Available at: https://www.natap.org/2024/CROI/croi_49.htm.
18. Waters L, Sparrowhawk A. Clinical implementation of long-acting antiretroviral treatment in high-income countries: challenges and advantages. *Curr Opin HIV AIDS*. 2022;17(3):121-126. Available at: <https://pubmed.ncbi.nlm.nih.gov/35439786>.
19. Rakhmanina N, Richards K, Adeline Koay WL. Transient viremia in young adults with HIV after the switch to long-acting cabotegravir and rilpivirine: considerations for dosing schedule and monitoring. *J Acquir Immune Defic Syndr*. 2023;92(3):e14-e17. Available at: <https://pubmed.ncbi.nlm.nih.gov/36480701>.
20. Llibre JM, Hung CC, Brinson C, et al. Efficacy, safety, and tolerability of dolutegravir-rilpivirine for the maintenance of virological suppression in adults with HIV-1: phase 3, randomised, non-inferiority SWORD-1 and SWORD-2 studies. *Lancet*. 2018;391(10123):839-849. Available at: <https://pubmed.ncbi.nlm.nih.gov/29310899>.

21. Aboud M, Orkin C, Podzamczer D, et al. Efficacy and safety of dolutegravir-rilpivirine for maintenance of virological suppression in adults with HIV-1: 100-week data from the randomised, open-label, phase 3 SWORD-1 and SWORD-2 studies. *Lancet HIV*. 2019;6(9):e576-e587. Available at: <https://pubmed.ncbi.nlm.nih.gov/31307948>.
22. van Wyk J, Orkin C, Rubio R, et al. Brief report: durable suppression and low rate of virologic failure 3 years after switch to dolutegravir + rilpivirine 2-drug regimen: 148-week results from the SWORD-1 and -2 randomized clinical trials. *J Acquir Immune Defic Syndr*. 2020;85(3):325-330. Available at: <https://pubmed.ncbi.nlm.nih.gov/32675772>.
23. Compagnucci A, Chan MK, Saidi Y, et al. Nucleoside/nucleotide reverse transcriptase inhibitor sparing regimen with once daily integrase inhibitor plus boosted darunavir is non-inferior to standard of care in virologically-suppressed children and adolescents living with HIV –week 48 results of the DA SMILE Penta-17-ANRS 152 clinical trial. *EClinicalMedicine*. 2023;60:102025. Available at: <https://pubmed.ncbi.nlm.nih.gov/37304494>.
24. Libre JM, Brites C, Cheng CY, et al. Efficacy and safety of switching to the 2-drug regimen dolutegravir/lamivudine versus continuing a 3- or 4-drug regimen for maintaining virologic suppression in adults living with human immunodeficiency virus 1 (HIV-1): week 48 results from the phase 3, noninferiority SALSA randomized Trial. *Clin Infect Dis*. 2023;76(4):720-729. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35235656>.
25. Osiyemi O, De Wit S, Ajana F, et al. Efficacy and safety of switching to dolutegravir/lamivudine versus continuing a tenofovir alafenamide-based 3- or 4-drug regimen for maintenance of virologic suppression in adults living with human immunodeficiency virus type 1: results through week 144 from the phase 3, noninferiority TANGO randomized trial. *Clin Infect Dis*. 2022;75(6):975-986. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35079789>.
26. Puthanakit T, Aurpibul L, Lopez M, et al. Efficacy and safety of dolutegravir/lamivudine (DTG/3TC) in antiretroviral therapy (ART)- naive adolescents living with HIV-1: DANCE study week 96 results. Presented at International AIDS Society; July 23-26, 2023, Year; Brisbane, Australia. Available at: https://medinfo.gsk.com/5f95dbd7-245e-4e65-9f36-1a99e28e5bba/450b1aad-cf09-4084-b484-0c909235ec90/450b1aad-cf09-4084-b484-0c909235ec90_viewable_rendition_v.pdf?medcommid=REF--ALL_00509&product=Dolutegravir%2BLamivudine.
27. Kuhn L, Coovadia A, Strehlau R, et al. Switching children previously exposed to nevirapine to nevirapine-based treatment after initial suppression with a protease-inhibitor-based regimen: long-term follow-up of a randomised, open-label trial. *Lancet Infect Dis*. 2012;12(7):521-530. Available at: <https://pubmed.ncbi.nlm.nih.gov/22424722>.
28. Coovadia A, Abrams EJ, Strehlau R, et al. Efavirenz-based antiretroviral therapy among nevirapine-exposed HIV-infected children in South Africa: a randomized clinical trial. *JAMA*. 2015;314(17):1808-1817. Available at: <https://pubmed.ncbi.nlm.nih.gov/26529159>.
29. Murnane PM, Strehlau R, Shiau S, et al. Switching to efavirenz versus remaining on ritonavir-boosted lopinavir in HIV-infected children exposed to nevirapine: long-term

- outcomes of a randomized trial. *Clin Infect Dis*. 2017;65(3):477-485. Available at: <https://pubmed.ncbi.nlm.nih.gov/28419200>.
30. Natukunda E, Rodriguez C, McGrath E, et al. B/F/TAF in virologically suppressed adolescents and children: two-year outcomes in 6 to <18 year olds and six-month outcomes in toddlers. Presented at: 13th International Workshop on HIV Pediatrics 2021. Virtual. Available at: https://www.natap.org/2021/IAS/IAS_80.htm.
31. Anugulruengkitt S, A. Gaur, P. Kosalaraksa, A. Liberty, Y. Shao, et al. . Long-term safety & efficacy of elvitegravir/cobicistat/emtricitabine/tenofovir alafenamide fumarate (E/C/F/TAF) single-tablet regimen in children and adolescents living with HIV. Abstract 4 Presented at: International Workshop on HIV Pediatrics 2021 2021. Virtual. Available at: https://www.natap.org/2021/IAS/IAS_79.htm.
32. Turkova A, White E, Mujuru HA, et al. Dolutegravir as first- or second-line treatment for HIV-1 infection in children. *N Engl J Med*. 2021;385(27):2531-2543. Available at: <https://pubmed.ncbi.nlm.nih.gov/34965338>.
33. Amuge P, Lugemwa A, Wynne B, et al. Once-daily dolutegravir-based antiretroviral therapy in infants and children living with HIV from age 4 weeks: results from the below 14 kg cohort in the randomised ODYSSEY trial. *Lancet HIV*. 2022;9(9):e638-e648. Available at: <https://pubmed.ncbi.nlm.nih.gov/36055295>.
34. Gill MM, Herrera N, Guilaze R, et al. Virologic outcomes and ARV switch profiles 2 years after national rollout of dolutegravir to children less than 15 years in southern Mozambique. *Pediatr Infect Dis J*. 2023;42(10):893-898. Available at: <https://pubmed.ncbi.nlm.nih.gov/37409808>.
35. Kouamou V, Manasa J, Maposphere C, et al. Tenofovir, lamivudine and dolutegravir (TLD) among rural adolescents in Zimbabwe, A cautionary tale. Presented at: International Workshop on HIV Pediatrics 2021. Virtual Available at: <https://academicmedicaleducation.com/meeting/international-workshop-hiv-pediatrics-2021/abstract/tenofovir-lamivudine-and-dolutegravir>.
36. Van de Ven R, Antelman G, Masenge T, Kimambo S. Impact of ARV optimization on HIV viral load suppression among children in Tanzania. Presented at: International AIDS Society 2021. Virtual Available at: <https://theprogramme.ias2021.org/Abstract/Abstract/1498>.
37. Bacha JM, Dlamini S, Anabwani F, et al. Realizing the promise of dolutegravir in effectively treating children and adolescents living with HIV in real-world settings in 6 countries in eastern and southern Africa. *Pediatr Infect Dis J*. 2023;42(7):576-581. Available at: <https://pubmed.ncbi.nlm.nih.gov/36795586>.
38. Rodriguez C. SR, Chokephaibulkit K., , et al. One year outcome of bictegravir/emtricitabine/tenofovir alafenamide (B/F/TAF) in virologically suppressed children ≥ 2 years weighing 14 to < 25kg. Presented at: Internataional Workshop on HIV & Pediatrics 2022. Montreal, Canada. Available at: https://www.natap.org/2022/IAC/IAC_90.htm.

39. Gaur AH, Cotton MF, Rodriguez CA, et al. Fixed-dose combination bictegravir, emtricitabine, and tenofovir alafenamide in adolescents and children with HIV: week 48 results of a single-arm, open-label, multicentre, phase 2/3 trial. *Lancet Child Adolesc Health.* 2021;5(9):642-651. Available at: <https://pubmed.ncbi.nlm.nih.gov/34302760>.
40. Phongsamart W, Jantarabenjakul W, Chantaratin S, et al. Switching efavirenz to rilpivirine in virologically suppressed adolescents with HIV: a multi-centre 48-week efficacy and safety study in Thailand. *J Int AIDS Soc.* 2022;25(1):e25862. Available at: <https://pubmed.ncbi.nlm.nih.gov/35001501>.
41. Melvin AJ, Yee KL, Gray KP, et al. Pharmacokinetics, tolerability, and safety of doravirine and doravirine/lamivudine/tenofovir disoproxil fumarate fixed-dose combination tablets in adolescents living with HIV: Week 24 results from IMPAACT 2014. *J Acquir Immune Defic Syndr.* 2023;92(2):153-161. Available at: <https://pubmed.ncbi.nlm.nih.gov/36215957>.
42. Vigano A, Aldrovandi GM, Giacomet V, et al. Improvement in dyslipidaemia after switching stavudine to tenofovir and replacing protease inhibitors with efavirenz in HIV-infected children. *Antivir Ther.* 2005;10(8):917-924. Available at: <https://pubmed.ncbi.nlm.nih.gov/16430197>.
43. Fabiano V, Giacomet V, Vigano A, et al. Long-term body composition and metabolic changes in HIV-infected children switched from stavudine to tenofovir and from protease inhibitors to efavirenz. *Eur J Pediatr.* 2013;172(8):1089-1096. Available at: <https://pubmed.ncbi.nlm.nih.gov/23636286>.
44. Rosso R, Nasi M, Di Biagio A, et al. Effects of the change from stavudine to tenofovir in human immunodeficiency virus-infected children treated with highly active antiretroviral therapy: studies on mitochondrial toxicity and thymic function. *Pediatr Infect Dis J.* 2008;27(1):17-21. Available at: <https://pubmed.ncbi.nlm.nih.gov/18162932>.
45. Aurpibul L, Puthanakit T, Sirisanthana T, Sirisanthana V. Haematological changes after switching from stavudine to zidovudine in HIV-infected children receiving highly active antiretroviral therapy. *HIV Med.* 2008;9(5):317-321. Available at: <https://pubmed.ncbi.nlm.nih.gov/18331562>.
46. Gonzalez-Tome MI, Amador JT, Pena MJ, et al. Outcome of protease inhibitor substitution with nevirapine in HIV-1 infected children. *BMC Infect Dis.* 2008;8:144. Available at: <https://pubmed.ncbi.nlm.nih.gov/18945352>.
47. Arribas JR, Pialoux G, Gathe J, et al. Simplification to coformulated elvitegravir, cobicistat, emtricitabine, and tenofovir versus continuation of ritonavir-boosted protease inhibitor with emtricitabine and tenofovir in adults with virologically suppressed HIV (STRATEGY-PI): 48 week results of a randomised, open-label, phase 3b, non-inferiority trial. *Lancet Infect Dis.* 2014;14(7):581-589. Available at: <https://pubmed.ncbi.nlm.nih.gov/24908551>.
48. Martinez E, Larrousse M, Llibre JM, et al. Substitution of raltegravir for ritonavir-boosted protease inhibitors in HIV-infected patients: the SPIRAL study. *AIDS.* 2010;24(11):1697-1707. Available at: <https://pubmed.ncbi.nlm.nih.gov/20467288>.

49. Curran A, Martinez E, Saumoy M, et al. Body composition changes after switching from protease inhibitors to raltegravir: SPIRAL-LIP substudy. *AIDS*. 2012;26(4):475-481. Available at: <https://pubmed.ncbi.nlm.nih.gov/22112606>.
50. Bagella P, Squillace N, Ricci E, et al. Lipid profile improvement in virologically suppressed HIV-1-infected patients switched to dolutegravir/abacavir/lamivudine: data from the SCOLTA project. *Infect Drug Resist*. 2019;12:1385-1391. Available at: <https://pubmed.ncbi.nlm.nih.gov/31213857>.
51. Calza L, Colangeli V, Borderi M, et al. Improvement in insulin sensitivity and serum leptin concentration after the switch from a ritonavir-boosted PI to raltegravir or dolutegravir in non-diabetic HIV-infected patients. *J Antimicrob Chemother*. 2019;74(3):731-738. Available at: <https://pubmed.ncbi.nlm.nih.gov/30541118>.
52. Rose PC, De la Rey Nel E, Cotton MF, et al. Decreased hepatic steatosis in South African adolescents with perinatal HIV switching to dolutegravir-containing regimens. *Pediatr Infect Dis J*. 2023;42(7):564-572. Available at: <https://pubmed.ncbi.nlm.nih.gov/36917035>.
53. Eckard AR, McComsey GA. Weight gain and integrase inhibitors. *Curr Opin Infect Dis*. 2020;33(1):10-19. Available at: <https://pubmed.ncbi.nlm.nih.gov/31789693>.
54. Sokhela S, Venter WDF, Bosch B, et al. Final 192-week efficacy and safety results of the ADVANCE trial, comparing 3 first-line antiretroviral regimens. *Open Forum Infect Dis*. 2024;11(3):ofae007. Available at: <https://pubmed.ncbi.nlm.nih.gov/38529213>.
55. Dirajlal-Fargo S, Koay WLA, Levy ME, et al. Effect of integrase inhibitors on weight gain in children and adolescents with HIV. Abstract 826. Presented at: Conference on Retroviruses and Opportunistic Infections; 2020. Boston, MA. Available at: <https://www.croiconference.org/abstract/effect-of-integrase-inhibitors-on-weight-gain-in-children-and-adolescents-with-hiv>.
56. Yeoh DK, Campbell AJ, Bowen AC. Increase in body mass index in children with HIV, switched to tenofovir alafenamide fumarate or dolutegravir containing antiretroviral regimens. *Pediatr Infect Dis J*. 2021;40(5):e215-e216. Available at: <https://pubmed.ncbi.nlm.nih.gov/33847305>.
57. Mallon PW, Brunet L, Hsu RK, et al. Weight gain before and after switch from TDF to TAF in a U.S. cohort study. *J Int AIDS Soc*. 2021;24(4):e25702. Available at: <https://pubmed.ncbi.nlm.nih.gov/33838004>.
58. McComsey GA, Lupo S, Parks D, et al. Switch from tenofovir disoproxil fumarate combination to dolutegravir with rilpivirine improves parameters of bone health. *AIDS*. 2018;32(4):477-485. Available at: <https://pubmed.ncbi.nlm.nih.gov/29239893>.
59. Soriano V, Fernandez-Montero JV, Benitez-Gutierrez L, et al. Dual antiretroviral therapy for HIV infection. *Expert Opin Drug Saf*. 2017;16(8):923-932. Available at: <https://pubmed.ncbi.nlm.nih.gov/28621159>.

60. Arribas JR, Girard PM, Paton N, et al. Efficacy of protease inhibitor monotherapy vs. triple therapy: meta-analysis of data from 2303 patients in 13 randomized trials. *HIV Med.* 2016;17(5):358-367. Available at: <https://pubmed.ncbi.nlm.nih.gov/26709605>.
61. Brenner BG, Thomas R, Blanco JL, et al. Development of a G118R mutation in HIV-1 integrase following a switch to dolutegravir monotherapy leading to cross-resistance to integrase inhibitors. *J Antimicrob Chemother.* 2016;71(7):1948-1953. Available at: <https://pubmed.ncbi.nlm.nih.gov/27029845>.
62. Wijting IEA, Wit F, Rokx C, et al. Immune reconstitution inflammatory syndrome in HIV infected late presenters starting integrase inhibitor containing antiretroviral therapy. *EClinicalMedicine.* 2019;17:100210. Available at: <https://pubmed.ncbi.nlm.nih.gov/31891143>.
63. Rokx C, Schurink CA, Boucher CA, Rijnders BJ. Dolutegravir as maintenance monotherapy: first experiences in HIV-1 patients. *J Antimicrob Chemother.* 2016;71(6):1632-1636. Available at: <https://pubmed.ncbi.nlm.nih.gov/26888910>.
64. Pinnetti C, Lorenzini P, Cozzi-Lepri A, et al. Randomized trial of DRV/r or LPV/r QD monotherapy vs maintaining a PI/r-based antiretroviral regimen in persons with suppressed HIV replication. *J Int AIDS Soc.* 2014;17(4 Suppl 3):19809. Available at: <https://pubmed.ncbi.nlm.nih.gov/25397553>.
65. Santos JR, Llibre JM, Bravo I, et al. Short communication: efficacy and safety of treatment simplification to lopinavir/ritonavir or darunavir/ritonavir monotherapy: a randomized clinical trial. *AIDS Res Hum Retroviruses.* 2016;32(5):452-455. Available at: <https://pubmed.ncbi.nlm.nih.gov/26781004>.
66. Kosalaraksa P, Ananworanich J, Puthanakit T, et al. Long-term lopinavir/ritonavir monotherapy in HIV-infected children. *Pediatr Infect Dis J.* 2013;32(4):350-353. Available at: <https://pubmed.ncbi.nlm.nih.gov/23190774>.
67. Agwu AL, Fairlie L. Antiretroviral treatment, management challenges and outcomes in perinatally HIV-infected adolescents. *J Int AIDS Soc.* 2013;16:18579. Available at: <https://pubmed.ncbi.nlm.nih.gov/23782477>.
68. Wensing AM, Calvez V, Gunthard HF, et al. 2015 Update of the drug resistance mutations in HIV-1. *Top Antivir Med.* 2015;23(4):132-141. Available at: <https://pubmed.ncbi.nlm.nih.gov/26713503>.
69. Dehority W, Deville JG, Lujan-Zilberman J, et al. Effect of HIV genotypic drug resistance testing on the management and clinical course of HIV-infected children and adolescents. *Int J STD AIDS.* 2013;24(7):549-553. Available at: <https://pubmed.ncbi.nlm.nih.gov/23970770>.
70. Tobin NH, Learn GH, Holte SE, et al. Evidence that low-level viremias during effective highly active antiretroviral therapy result from two processes: expression of archival virus and replication of virus. *J Virol.* 2005;79(15):9625-9634. Available at: <https://pubmed.ncbi.nlm.nih.gov/16014925>.

71. Kuritzkes DR. Preventing and managing antiretroviral drug resistance. *AIDS Patient Care STDS*. 2004;18(5):259-273. Available at: <https://pubmed.ncbi.nlm.nih.gov/15186710>.