

Nutrition Profile



DEMOGRAPHICS

Total population (000) 1,281 (2008) Total under-five population (000) 77 (2008) Total number of births (000) 15 (2008) Under-five mortality rate (per 1,000 live births) 10 (2008) Total number of under-five deaths (000) 0 (2008) Infant mortality rate (per 1,000 live births) 9 (2008) Neonatal mortality rate (per 1,000 live births) 4 (2004) HIV prevalence rate (15-49 years, %) - - Population below international poverty line of US\$1.25 per day (%) - -			
Total number of births (000) 15 (2008) Under-five mortality rate (per 1,000 live births) 10 (2008) Total number of under-five deaths (000) 0 (2008) Infant mortality rate (per 1,000 live births) 9 (2008) Neonatal mortality rate (per 1,000 live births) 4 (2004) HIV prevalence rate (15-49 years, %)	Total population (000)	1,281	(2008)
Under-five mortality rate (per 1,000 live births) Total number of under-five deaths (000) Infant mortality rate (per 1,000 live births) Neonatal mortality rate (per 1,000 live births) HIV prevalence rate (15-49 years, %) Population below international	Total under-five population (000)	77	(2008)
births) 10 (2008) Total number of under-five deaths (000) 0 (2008) Infant mortality rate (per 1,000 live births) 9 (2008) Neonatal mortality rate (per 1,000 live births) 4 (2004) HIV prevalence rate (15-49 years, %)	Total number of births (000)	15	(2008)
Infant mortality rate (per 1,000 live births) 9 (2008) Neonatal mortality rate (per 1,000 live births) 4 (2004) HIV prevalence rate (15-49 years, %) Population below international	• •	10	(2008)
Neonatal mortality rate (per 1,000 live births) HIV prevalence rate (15-49 years, %) Population below international	Total number of under-five deaths (000)	0	(2008)
births) 4 (2004) HIV prevalence rate (15-49 years, %) Population below international	Infant mortality rate (per 1,000 live births)	9	(2008)
Population below international	• •	4	(2004)
	HIV prevalence rate (15-49 years, %)		








NUTRITIONAL STATUS

Burden of undernutrition (2008) NCHS reference population


Stunted (under-fives, 000): 6 Underweight (under-fives, 000): 4 Share of developing world stunting burden (%): Wasted (under-fives, 000) Stunting country rank: 130 Severely wasted (under-fives, 000):








INFANT AND YOUNG CHILD FEEDING



No Data



Weaned (not breastfed) Breastfed and

Breastfed and solid/

Qatar

MICRONUTRIENTS


Vitamin A supplementation Percentage of children 6-59 months old receiving two doses of vitamin A during calendar year

No Data

lodized salt consumption trends Percentage of households consuming adequately iodized salt

15,000 newborns are unprotected against IDD (2008)

No Data




ESSENTIAL NUTRITION INTERVENTIONS DURING THE LIFE CYCLE

Pregnancy	Birth	0-5 months	6-23 months	>	24-59 months		
Use of iron-folic acid	Early initiation of breastfeeding	International Code of Marketing of Breastmilk Substitutes		Partial			
supplements	(within 1 hour of birth)	Maternity protection in accordance	ce with ILO Convention 183	No			
Household consumption of adequately iodized salt	Infants not weighed at birth .	Exclusive breastfeeding (< 6 months) 12%	Timely introduction of complementary foods (with continued breastfeeding)	48%			
		Continued breastfeeding at two years	21%				
To increase children	Full coverage of vitamin A supplementation						
development and prevent stunting, nutrition interventions need to be delivered during the mother's pregnancy and the			National guidelines for management of severe acute malnutrition incorporating the community-based approach				
first two years of the	ne chila's lite.	Policy on new ORS formula and zinc for management of diarrhoea*					
	Informa	Policy on community treatment of pneumonia with antibiotics					

MATERNAL NUTRITION/HEALTH WATER AND SANITATION

Maternal mortality ratio, adjusted (per 100,000 live births)	12	(2005)
Maternal mortality ratio, reported (per 100,000 live births)	10	(1996)
Total number of maternal deaths	2	(2005)
Lifetime risk of maternal deaths (1 in :)	2,700	(2005)
Women with low BMI ($< 18.5 \text{ kg/m}^2$, %)		
Anaemia, non-pregnant woman (< 120 g/l, %)	36	(2008)
Antenatal care (at least one visit, %)		•
Antenatal care (at least four visits, %)		
Skilled attendant at birth (%)	99	(1998)
Low birthweight ($<$ 2,500 grams, $\%$)	10	(1999)
Primary school net enrolment or attendance ratio (% female, % male)	95, 94	(2007)
Gender parity index (primary school net enrolment or attendance ratio)	1.01	(2007)





1%

Under-five deaths caused by:

Diarrhoea:

4% Pneumonia:

DISPARITIES IN NUTRITION

	Gender			Residence		Wealth quintile							
Indicator	Male	Female	Ratio of male to female	Urban	Rural	Ratio of urban to rural	Poorest	Second	Middle	Fourth	Richest	Ratio of richest to poorest	Source
Stunting prevalence (WHO Child Growth Standards, %)	-			-									
Underweight prevalence (WHO Child Growth Standards, %)					-								
Wasting prevalence (WHO Child Growth Standards, %)				-	-	-	-					-	
Infants not weighed at birth (%)			-		-								
Early initiation of breastfeeding (%)			-	-	-		-						
Women with low BMI ($<$ 18.5 kg/m², %)	-		-										