

## Method

# CRISPR-based modular assembly of a UAS-cDNA/ORF plasmid library for more than 5500 *Drosophila* genes conserved in humans

Ping Wei,<sup>1</sup> Wen Xue,<sup>2,3</sup> Yun Zhao,<sup>2</sup> Guang Ning,<sup>4</sup> and Jiwu Wang<sup>3,5</sup>

<sup>1</sup>Shanghai Diabetes Institute, Shanghai Key Laboratory of Diabetes Mellitus, Shanghai Clinical Center for Diabetes, Shanghai Jiao Tong University Affiliated Sixth People's Hospital, Shanghai 200233, China; <sup>2</sup>Cyrus Tang Hematology Center, Soochow University, Suzhou 215123, China; <sup>3</sup>Shanghai Institute of Endocrine and Metabolic Diseases, Shanghai 200025, China; <sup>4</sup>Shanghai National Clinical Research Center for Endocrine and Metabolic Diseases, Key Laboratory for Endocrine and Metabolic Diseases of the National Health Commission of the PR China, Shanghai Institute of Endocrine and Metabolic Diseases, Ruijin Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai 200025, China; <sup>5</sup>Department of Anatomy and Physiology, Shanghai Jiao Tong University School of Medicine, Shanghai 200025, China

Construction of a genome-wide transgenic UAS-cDNA/ORF library in *Drosophila* based on the binary GAL4/UAS system has been severely hampered by technical difficulties, although genome-wide cDNA or ORF resources of *Drosophila*, human, and mouse have been publicly available for more than a decade. Here, we developed a new method named CRISPR-based modular assembly (CRISPRmass) for the high-throughput construction of a genome-wide UAS-cDNA/ORF library from publicly available cDNA/ORF resources. Through cleavage of shared vector sequences of cDNA/ORF plasmids by CRISPR/Cas9 and subsequent insertion of UAS modules by Gibson assembly, the procedure of construction of such a library by CRISPRmass is standardized as massively parallel two-step test tube reactions before bacterial transformation. Using CRISPRmass, we generated 5551 UAS-cDNA/ORF constructs covering 83% of the *Drosophila* genes conserved in humans in the *Drosophila* Genomics Resource Center (DGRC) Gold Collection, and among them, 5518 were generated within 3 mo by three people. Our results show that CRISPRmass allows modularization, simplicity, efficiency, and adaptability in the generation of a genome-wide UAS-cDNA/ORF plasmid library by using publicly available cDNA/ORF resources. CRISPRmass can be applied to editing various genome-wide libraries in general and is an alternative to Gateway technology in high-throughput plasmid library editing. Furthermore, the more than 5500 UAS-cDNA/ORF plasmids of *Drosophila* genes serve as a powerful resource for gain-of-function (GOF) screening in cultured cells and for generation of a transgenic UAS-cDNA/ORF library in *Drosophila*.

[Supplemental material is available for this article.]

Genome-wide genetic screens are powerful tools to explore gene function in complex biological processes. In *Drosophila*, genome-wide genetic screens have been relying on loss-of-function (LOF) screens. Based on the binary GAL4/upstream activation sequence (UAS) system (Brand and Perrimon 1993), genome-wide transgenic UAS-RNAi libraries for LOF screens have been constructed for targeted knockdown of most of the *Drosophila* genes (Dietzl et al. 2007; Perkins et al. 2015). However, the majority of *Drosophila* genes show no obvious LOF phenotypes (Miklos and Rubin 1996), therefore the function of such genes cannot be determined by LOF studies. Thus, gain-of-function (GOF) screens may provide a valuable tool to uncover function of such genes (Miklos and Rubin 1996; St Johnston 2002).

In contrast to construction of genome-wide transgenic UAS-RNAi libraries for LOF screens, construction of a genome-wide transgenic UAS-cDNA/ORF library for GOF screens in *Drosophila* has been challenging. Genome-wide cDNA/ORF resources of *Drosophila* (Rubin et al. 2000; Stapleton et al. 2002), human (Yang et al. 2011), and mouse (<https://genecollections.nci.nih.gov/MGC>) have been publicly available for more than a decade,

but the efforts by different research groups to create such a genome-wide cDNA/ORF library using these resources have never ceased. In 2008, a collection of UAS-ORF plasmids of 236 human genes and the corresponding transgenic fly lines have been created based on the conventional cloning technique (Xu et al. 2008). In 2013, a platform of the recombinational (Gateway) cloning technique was established and used successfully in constructing the UAS-ORF plasmids for 655 *Drosophila* genes (Bischof et al. 2013); based on this platform, to date, the UAS-ORF plasmids and transgenic flies mapping to about 2850 *Drosophila* genes have been generated (FlyORF, <http://www.FlyORF.ch>). In 2018, the human ORF collection containing 365 UAS-ORF plasmids for *Drosophila* transgenesis was constructed and deposited to the *Drosophila* Genomics Resource Center (DGRC, <https://dgrc.bio.indiana.edu>). Despite these efforts, there is still a long way to go before a genome-wide transgenic *Drosophila* UAS-cDNA/ORF library is created. The rate-limiting step of construction of a genome-wide transgenic *Drosophila* UAS-cDNA/ORF library is the technical difficulty of constructing a library of UAS-cDNA/ORF plasmids. Currently,

**Corresponding author:** [jiwu@wang-lab.cn](mailto:jiwu@wang-lab.cn)

Article published online before print. Article, supplemental material, and publication date are at <http://www.genome.org/cgi/doi/10.1101/gr.250811.119>.

© 2020 Wei et al. This article is distributed exclusively by Cold Spring Harbor Laboratory Press for the first six months after the full-issue publication date (see <http://genome.cshlp.org/site/misc/terms.xhtml>). After six months, it is available under a Creative Commons License (Attribution-NonCommercial 4.0 International), as described at <http://creativecommons.org/licenses/by-nc/4.0/>.

construction of such a plasmid library has almost exclusively relied on the recombinational (Gateway) cloning technique (Bischof et al. 2013). The recombinational (Gateway) cloning technique requires handling tens of thousands of cDNAs or ORFs by time-consuming individualized manipulations of every single gene including primer design and synthesis, PCR, gel purification, and sequencing. Particularly, construction of UAS-cDNA/ORF plasmids for long cDNAs or ORFs by the recombinational (Gateway) cloning technique is truly problematic. Therefore, the technical difficulties of construction of a library of UAS-cDNA/ORF plasmids have severely hampered construction of such a genome-wide transgenic UAS-cDNA/ORF library in *Drosophila*.

To achieve the goal of performing genome-wide GOF screens, some research groups established alternative approaches that circumvented construction of UAS-cDNA/ORF plasmids. Through random insertion of the P-element transposon bearing a UAS enhancer and an *Hsp70* minimal promoter (EP) into the *Drosophila* genome, about 8500 EP lines have been created (Rørth 1996; Rørth et al. 1998; Mata et al. 2000). Because more than one EP line may overexpress or misexpress the same gene in the presence of GAL4, it is unclear how many genes were covered by these lines, probably because it is very laborious to identify genomic locations of UAS insertions (Mata et al. 2000). Currently a representative set of these EP lines containing about 1700 EP lines which maps to fewer than 1700 genes is available in the Bloomington *Drosophila* Stock Center and covers <12.38% of 13,733 *Drosophila* euchromatic protein-coding genes (Lin et al. 2007). However, this approach has intrinsic drawbacks. If an EP element inserts into a gene in antisense orientation, an antisense transcript may be generated and therefore causes loss of function of the gene. Furthermore, EP lines do not allow expression of genes from other organisms including human. The P-element transposon shows inherent insertional hot spots and orientation bias, and therefore it is impossible to construct a saturated genome-wide GOF library through generation of EP lines (St Johnston 2002). Nevertheless, GOF screens by EP lines have made great contributions to biological research over the past two decades.

CRISPRa is another approach to construct a genome-wide GOF library. During the past two years, two CRISPRa-based methods for construction of a genome-wide GOF library in *Drosophila* were developed (Ewen-Campen et al. 2017; Jia et al. 2018). These two methods serve as useful strategies for GOF screening, especially for genes of which the cDNA/ORFs are not available. However, for every single gene, DNA oligos for at least one sgRNA have to be individually well designed in order to improve success rates of sgRNAs and to reduce their off-target effects, followed by synthesis of DNA oligos and sequencing of sgRNA constructs (Jia et al. 2018). Furthermore, these two CRISPRa-based methods have some other drawbacks due to the intrinsic properties of CRISPRa. For example, CRISPRa-based GOF is sometimes toxic in vivo even in the absence of sgRNAs when crossed with some GAL4 drivers (Jia et al. 2018); sometimes two or more genes share a common promoter region in the *Drosophila* genome, which may cause the expression of more than one gene to be controlled by a single sgRNA, and *Drosophila* CRISPRa does not allow expression of genes from other organisms, including human.

Ideally, the best genome-wide GOF library in *Drosophila* is still the standard cDNA/ORF-based transgenic UAS-cDNA/ORF library, which can overcome all the aforementioned drawbacks. Therefore, there is an urgent need to develop a simple and efficient method of creating a genome-wide UAS-cDNA/ORF plasmid library using cDNA/ORF resources of different species. Here, we developed

a new method named CRISPR-based modular assembly (CRISPRmass) for high-throughput construction of a genome-wide UAS-cDNA/ORF plasmid library from publicly available cDNA/ORF resources, and using CRISPRmass, we created more than 5500 UAS-cDNA/ORF plasmids for *Drosophila* genes conserved in humans.


## Results

### The CRISPRmass pipeline for construction of a UAS-cDNA/ORF plasmid library

The procedure of construction of a genome-wide UAS-cDNA/ORF plasmid library from publicly available cDNA/ORF resources by CRISPRmass was standardized as massively parallel two-step test tube reactions before bacterial transformation (Fig. 1). The first step is that the identical vector backbones of the cDNA/ORF constructs are cleaved by Cas9/sgRNA. The cleavage site is adjacent to the 5' end of cDNA/ORF. The backbones of some vectors are cleaved twice by Cas9/sgRNA in order to remove unnecessary DNA sequence. The cleavage products, without purification, are directly subjected to the subsequent step. The second step is that a vector-specific UAS module is joined into the Cas9-linearized cDNA/ORF constructs through Gibson assembly, resulting in UAS-cDNA/ORF constructs. The vector-specific UAS module comprises a core UAS module and 28–40 bps of 5'- and 3'-terminal sequences overlapping with the backbone cleavage site 5' and 3' ends, respectively. The 5' and 3' overlapping terminal sequences enable Gibson assembly. A core UAS module comprises 10 copies of UAS, an *Hsp70* minimal promoter, an *attB* sequence for  $\phi$ C31-mediated genomic integration, a *mini-white* transformation marker for *Drosophila*, and a positively selectable antibiotic resistance gene distinct from that of the vector. A vector-specific UAS module can be built up by PCR modifying a core UAS module according to the vector backbone cleavage site. After the second step of the two test tube reactions, Gibson assembly products are directly subjected to *Escherichia coli* transformation. Transformants are selected on LB plates containing antibiotic corresponding to the antibiotic resistance gene of the vector-specific UAS module. Theoretically, only the desired UAS-cDNA/ORF colonies can grow.

### Proof of principle for construction of UAS-cDNA/ORF plasmids by CRISPRmass

To show the feasibility of using CRISPRmass to construct a UAS-cDNA/ORF plasmid library from publicly available cDNA or ORF resources such as the DGRC Gold Collection, we first attempted to create a UAS-cDNA/ORF plasmid sublibrary, UAS-DUBs, by CRISPRmass for all the 41 deubiquitinases (DUBs) of *Drosophila* from the DGRC Gold Collection (Tsou et al. 2012). There are three reasons for selecting DUBs to create a UAS-cDNA/ORF plasmid sublibrary as proof of principle. First, DUBs play important roles in diverse biological processes, and altered DUB activity is associated with a number of diseases. However, a collection of UAS-DUBs has not yet been reported. Second, of all the 41 *Drosophila* DUBs, the cDNA/ORF clones for 31 *Drosophila* DUBs are available in the DGRC Gold Collection, which were constructed in four of six total vectors, except pBS SK (–) and pCR2.1 vectors. To build a complete CRISPRmass platform for construction of the UAS-cDNA/ORF library from the DGRC Gold Collection, we only need to further show CRISPRmass also works for cDNA/ORF clones constructed in pBS SK (–) and pCR2.1; therefore, we randomly picked up a



**Figure 1.** The CRISPRmass pipeline for construction of a UAS-cDNA/ORF library. (1) The first step test tube reaction. The identical vector backbones of cDNA/ORF plasmids are cleaved by Cas9/sgRNA. The cleavage site is located in vector backbones adjacent upstream of the cDNA/ORF 5' end. The cleavage products, without purification, are directly subjected to the second step test tube reaction. The sgRNAs used in cleavage reactions are prepared by *in vitro* transcription and can be used directly without purification. Then, 28–40 bp of the 5' end of the cleavage site (yellow box) is defined as “5' end overlap,” and 28–40 bp of the 3' end of the cleavage site (red box) is defined as “3' end overlap.” The vector backbone carries the antibiotic A resistance gene (green box). (2) The second step test tube reaction. A vector-specific UAS module is joined into the Cas9-linearized cDNA/ORF plasmids right upstream of cDNA/ORF 5' end through Gibson assembly, resulting in UAS-cDNA/ORF constructs. Gibson assembly products are directly subjected to *E. coli* transformation. Transformants are selected on LB agar plates containing antibiotic B corresponding to the antibiotic B resistance gene (brown box) of the vector-specific UAS module. Only the desired UAS-cDNA/ORF colonies can grow. A UAS module comprises 10 copies of UAS, a *Hsp70* minimal promoter, an *attB* sequence for  $\phi$ C31-mediated genomic integration, a *mini-white* transformation marker for *Drosophila* transgenesis, a selectable antibiotic B resistance gene for positive selection, and the 5' end overlap and 3' end overlap enabling Gibson assembly. Gibson assembly filters out any potential off-target DNA cleavages caused by CRISPR/Cas9.

clone in which *CG7747* cDNA was cloned in pBS SK(–) and a clone in which *Prdm13* ORF was cloned in pCR2.1. Third, for the rest of the 10 *Drosophila* DUBs, six *Drosophila* DUB ortholog clones are available from the human CCSB-Broad Lentiviral Expression Library (Yang et al. 2011), and one *Drosophila* DUB ortholog clone is available from the mouse Mammalian Gene Collection (MGC). These human and mouse DUB clones allow us to examine whether CRISPRmass can apply to cDNA/ORF libraries of other species. The remaining three *Drosophila* DUBs do not have publicly available cDNA/ORF clones of human or mouse orthologs. Thus, in total, 40 *Drosophila*, human, and mouse cDNA/ORF clones correspond to eight distinct vectors (Supplemental Table S1). All the six human DUB ORFs from the human CCSB-Broad Lentiviral Expression Library were fused with a C-terminal V5 tag in pLX304 vector. To facilitate detection of MYC expression of *Drosophila* and mouse cDNA/ORFs by immunostaining and western blotting, we picked up one cDNA/ORF clone for each of the remaining seven vectors and inserted two or three copies of the MYC epitope tag right before the stop codon (Supplemental Table S1).

To generate *UAS-CG7747*, *UAS-Prdm13*, and 38 UAS-DUB constructs by CRISPRmass, we first determined the optimal Cas9 cleavage sites upstream of cDNA/ORF for all eight vectors. The vector backbones of cDNA/ORF clones in pOT2, pOTB7, pFLC-I, pBS SK(–), pCR2.1, and pDNR-Dual vectors are cleaved once by Cas9 together with a single sgRNA, whereas the vector backbones of cDNA/ORF clones in pLX304 and pCMV-SPORT6 vectors are cleaved twice by Cas9 together with two different sgRNAs to remove unnecessary sequences (Supplemental Fig. S1). The vector backbone of pOTB7\_DraIII is identical to that of pOTB7, so these two vectors can be treated as one in terms of sgRNA design. We designed 3–20 sgRNA target sequences for each cleavage site of the eight vectors and synthesized 53 sgRNAs by *in vitro* transcription (Supplemental Table S2). Then we examined *in vitro* cleavage efficiency of all these sgRNAs and found most of them can be used for CRISPRmass (Supplemental Figs. S2, S3; Supplemental Table S2). We selected one or two sgRNAs for each vector (Supplemental Table S3). We also searched the genome of the corresponding species for sequences with 14



consecutive nucleotides identical to the optimal sgRNA target sequences, and we did not find any potential off-target sites in *Drosophila*, human, or mouse cDNA sequences. Based on the cleavage sites determined by the optimal sgRNAs, we generated a UAS module for each of the eight vectors. After confirmation by DNA sequencing, the vector-specific UAS modules were released by EcoRI digestion and joined into the corresponding Cas9/sgRNA linearized cDNA/ORF plasmids by Gibson assembly. The Gibson assembly products were directly used to transform *E. coli* competent cells, and transformants were selected on LB plates containing ampicillin or chloramphenicol depending on the antibiotic resistance gene of the UAS modules. Subsequent isolation and restriction analysis of plasmids showed that we successfully generated *UAS-CG7747*, *UAS-Prdm13*, and 38 UAS-DUB constructs by CRISPRmass (Fig. 2; Supplemental Table S1). Similar to UAS-cDNA/ORF constructs created by conventional cloning techniques, the UAS-cDNA/ORF constructs generated on eight distinct vectors by CRISPRmass retain the key features of a classic UAS-cDNA/ORF construct of GAL4/UAS system, particularly the UAS and the *Hsp70* minimal promoter. Therefore, we reason that UAS-cDNA/ORF constructs generated by CRISPRmass should have no difference in terms of cDNA/ORF expression with those generated by conventional cloning techniques.


### The UAS-cDNA/ORF constructs generated by CRISPRmass were expressed in the presence of GAL4 in *Drosophila*

We next generated transgenic *UAS-CG7747*, *UAS-Prdm13*, and 38 UAS-DUB lines in *Drosophila* using these 40 constructs created in eight distinct vectors by CRISPRmass. To test whether these transgenes can be expressed in the presence of GAL4 *in vivo*, we selected one tagged transgene for each of eight vectors and crossed to motor neuron-specific *OK6-GAL4* and eye-specific *GMR-GAL4* drivers, respectively. Immunostaining showed these transgenes were expressed in the larval motor neurons (Fig. 3A,B), and western blotting showed these transgenes were expressed in adult fly eyes (Fig. 3C–E). The levels of expressed proteins vary among different transgenes, probably due to differences in translation regulation and protein degradation, but not distinct vectors. Together, these results showed that the UAS-cDNA/ORF constructs in eight distinct vectors generated by CRISPRmass were expressed in the presence of GAL4 in *Drosophila*; therefore, we conclude that CRISPRmass can be used for construction of a genome-wide


UAS-cDNA/ORF library using the publicly available cDNA/ORF resources of *Drosophila*, human, and mouse.

### The UAS-cDNA/ORF constructs generated by CRISPRmass were functional in the presence of GAL4 in *Drosophila*

To further validate whether transgenes generated by CRISPRmass are functional, we screened the *UAS-CG7747*, *UAS-Prdm13*, and 38 UAS-DUB transgenes for phenotypes by crossing to eye-specific *GMR-GAL4*, wing-specific *A9-GAL4*, and muscle-specific *MEF2-GAL4* drivers, respectively. Activation of the *UAS-Prdm13* and several UAS-DUB transgenes elicited distinct phenotypes (Fig. 4A–C),



**Figure 2.** Restriction analysis of 40 UAS-cDNA/ORF constructs generated by CRISPRmass. *UAS-CG7747*, *UAS-Prdm13*, and 38 UAS-DUB constructs were generated by CRISPRmass from the DGRC Gold Collection and analyzed by PstI or PvuII digestion, and the expected restriction patterns for all the UAS-cDNA/ORF constructs were observed (Supplemental Table S1). (M) DNA marker.



**Figure 3.** The UAS-cDNA/ORF constructs generated by CRISPRmass were expressed in the presence of GAL4 in *Drosophila*. Transgenic *UAS-CG7747*, *UAS-Prdm13*, and 38 UAS-DUB lines were generated by CRISPRmass in *Drosophila* using the 40 cDNA/ORF plasmids constructed in eight distinct vectors. An epitope-tagged transgene for each of the eight vectors was chosen and crossed to motor neuron-specific *OK6-GAL4* and eye-specific *GMR-GAL4* drivers, respectively. (DUB) Deubiquitinase. (A,B) Immunostaining showed these transgenes were expressed in the larval motor neurons, and V5 signals are indicated by arrows. Presynaptic active zones are labeled with anti-BRP (nc82) (red). (C–E) Western blotting showed these transgenes were expressed in adult fly eyes. The epitope-tagged proteins and their expected molecular weights are as follows: CG4968-3MYC, 33.9 kD; Rpn8-3MYC, 41.6 kD; Usp5-3MYC, 95.6 kD; CG7747-3MYC, 62.5 kD; Usp20-33-3MYC, 113.2 kD; Prdm13-3MYC, 54.1 kD; Ctbp2-2MYC, 48.3 kD; USP8-V5, 130.0 kD.


indicating the *UAS-CG7747*, *UAS-Prdm13*, and 38 UAS-DUB transgenes generated by CRISPRmass are functional.

### Construction of a UAS-cDNA/ORF plasmid library for more than 5500 *Drosophila* genes conserved in humans by CRISPRmass

We applied CRISPRmass to generate a genome-wide UAS-cDNA/ORF plasmid library using conserved cDNA/ORF clones from the DGRC Gold Collection (Fig. 5). As far as we know, until now the most comprehensive *Drosophila* cDNA/ORF library is the DGRC Gold Collection, which comprises 12,192 clones mapping to 10,066 *Drosophila* genes, and among them, 6687 *Drosophila* genes conserved in humans are available based on OrthoDB v9.1 (Zdobnov et al. 2017). Given the general interest of researchers, we fo-

used on the 6687 *Drosophila* genes conserved in humans to generate a genome-scale UAS-cDNA/ORF library. For the 6687 conserved genes, excluding the clones of which vectors are not clearly specified in the DGRC list and the 33 clones for which UAS-cDNA/ORF plasmids have been previously constructed successfully, 6515 cDNA/ORF clones mapping to 6515 genes were subjected to CRISPRmass. At least one colony grew on LB plates with appropriate antibiotics for 5987 CRISPRmass reactions after transformation. We randomly picked up only one colony from each plate for subsequent restriction analysis. We performed restriction analysis of UAS-cDNA/ORF constructs with PstI- or PvuII-dependent specific vectors, because UAS modules bear at least two recognition sites for either of the two restriction enzymes, and meanwhile all these vectors bear at least one recognition site for either of the two restriction enzymes. Thus, restriction digestions of UAS-cDNA/ORF constructs with PstI or PvuII produce the UAS module-specific fragments and the UAS module-vector assembly-specific fragments. Indeed, we observed both UAS module-specific fragments and UAS module-vector assembly-specific fragments for 5903 constructs, demonstrating that UAS modules were inserted into 98.6% of the 5987 constructs successfully under the condition of randomly analyzing only one colony for each UAS-cDNA/ORF construct, resulting in 5903 UAS-cDNA/ORF constructs (Supplemental Fig. S4; Supplemental Table S4). Using an *Hsp70* minimal promoter-specific primer, we then sequenced the rest of 84 constructs for which no UAS module-specific fragments or UAS module-vector assembly-specific fragments were observed, and we obtained no sequencing signals.

We next analyzed the restriction patterns of the 5903 UAS-cDNA/ORF constructs and found the restriction patterns were as expected for 5397 UAS-cDNA/ORF constructs mapping to 5397 genes. The unexpected restriction patterns for the remaining 506 UAS-cDNA/ORF constructs were probably caused by mislabeled original cDNA/ORF clones from the DGRC Gold Collection or contamination of other UAS-cDNA/ORF constructs. Subsequent sequencing results of the 506 UAS-cDNA/ORF constructs indicated that 354 constructs were not unique in the 5923 UAS-cDNA/ORF constructs and were therefore discarded. The remaining 152 constructs were unique in the 5923 UAS-cDNA/ORF constructs; among them, 121 constructs of which cDNAs or ORFs are conserved in humans were retained and the remaining 31 constructs were discarded. In summary, we created 5518 UAS-cDNA/ORF constructs mapping to 5518 *Drosophila* genes conserved in humans. All 5518 UAS-cDNA/



**Figure 4.** The UAS-cDNA/ORF constructs generated by CRISPRmass were functional in the presence of GAL4 in *Drosophila*. UAS-*CG7747*, UAS-*Prdm13*, and 38 UAS-DUB transgenes were screened for phenotypes by crossing to eye-specific *GMR-GAL4* (A), wing-specific *A9-GAL4* (B), and muscle-specific *MEF2-GAL4* (C) drivers, respectively. Activation of the UAS-*Prdm13* and several UAS-DUB transgenes elicited distinct phenotypes (arrows).

ORF constructs were generated by CRISPRmass successfully within 3 mo by three people (Fig. 5).

We further evaluated the quality of the more than 5500 UAS-cDNA/ORF constructs generated by CRISPRmass. We fully Sanger sequenced the cDNA/ORFs of the first six UAS-cDNA/ORF constructs cloned in each of the four vectors on the list (Supplemental Tables S4–S6) and found that the entire sequences of cDNA/ORFs of the 24 UAS-cDNA/ORF constructs were all as expected, demonstrating that the accuracy rate of the cDNA/ORFs of those 24 constructs is 100%. To examine whether the cDNA/ORFs of those 24 constructs can be expressed in the presence of GAL4, we fused the 24 cDNA/ORFs with a 3xFLAG tag to the N terminus or C terminus and transfected HEK293T cells with each UAS-cDNA/ORF construct together with a GAL4-expressing plasmid, pCMV-GAL4 (Potter et al. 2010). The subsequent western blotting experiments detected specific signals for all the 24 cDNA/ORFs (Supplemental Fig. S5), demonstrating that all the 24 UAS-cDNA/ORFs generated by CRISPRmass were expressed in the presence of GAL4.


Together with the previously generated UAS-cDNA/ORF constructs for 31 *Drosophila* DUBs and two functionally unknown genes, we generated a total of 5551 UAS-cDNA/ORF constructs

mapping to 5551 genes, which cover 83% of the 6687 *Drosophila* genes conserved in humans (Supplemental Table S4).

Thus, CRISPRmass may be even more suitable for high-throughput UAS-cDNA/ORF library construction than the recombinational (Gateway) cloning technique, specifically when a suitable Gateway entry clone set is not available, but a clone set of any other sort does exist. Compared to the Gateway cloning technique, CRISPRmass does not need to generate any entry constructs and thus circumvent all the individualized manipulation for each entry construct, such as primer design and synthesis, PCR, sequencing, gel purification, and so forth. For example, for construction of a UAS-cDNA/ORF library containing 5551 genes from the DGRC Gold Collection, which were cloned in six distinct vectors, the Gateway cloning technique requires at least 5551 individualized manipulations, including primer design and synthesis, PCR, gel purification, and sequencing. In contrast, CRISPRmass allows us to manipulate shared vector sequences; therefore, we only need to design and synthesize primers for preparation of only six sgRNAs and to build up only six UAS modules by PCR modifying the core module according to the six vectors. The remaining manipulations are uniformed test tube reactions followed by *E. coli* transformation (Table 1). Moreover, of 6687 conserved genes in the DGRC Gold Collection, we observed that for the total 17 cDNAs of >7000 bps (the longest one

is 9337 bps), the UAS-cDNA constructs of 12 cDNAs were successfully created, whereas for the remaining five cDNAs, two original cDNA bacterial clones failed to grow and the other three original cDNA bacterial clones were not as expected probably due to mislabeling of the original clones. The size limit for CRISPRmass is not known, because we do not have cDNAs larger than 9337 bps at hand. We infer that the size limit for CRISPRmass depends on the second step reaction, insertion of modules by Gibson assembly, because the first step reaction of CRISPRmass, CRISPR/Cas9-mediated cleavage, is not affected by DNA size. It has been reported that Gibson assembly products as large as 300 kb have been cloned in *E. coli* successfully (Gibson et al. 2009). We therefore conclude that CRISPRmass is also suitable for generating UAS-cDNA/ORF constructs from long cDNAs or ORFs, which could be because CRISPRmass does not manipulate the cDNAs or ORFs themselves but vector backbones and the high cloning capacity of Gibson assembly. Furthermore, even if a suitable Gateway entry clone set is available, the cost of key commercially available kits for construction of the UAS-cDNA/ORF library by CRISPRmass is merely ~40% of that by the Gateway LR reaction (Table 1). In summary, CRISPRmass is faster and cheaper than Gateway for high-throughput construction of desired destination





**Figure 5.** Flow chart of generation of 5518 UAS-cDNA/ORF constructs within 3 mo, in which 6515 DGRC cDNA/ORF plasmids covering 6515 *Drosophila* genes conserved in humans were subjected to CRISPRmass reactions. At least one colony grew on LB agar plates with appropriate antibiotics for 5987 reactions after transformation. Only one colony from each plate was randomly picked up for subsequent restriction analysis. UAS module-specific fragments and UAS module-vector assembly-specific fragments were observed in 5903 constructs, demonstrating that UAS modules were inserted into 98.6% of the 5987 constructs successfully, resulting in 5903 UAS-cDNA/ORF constructs. After a series of analyses of 5903 UAS-cDNA/ORF constructs were conducted, 5518 UAS-cDNA/ORF constructs covering 5518 *Drosophila* genes conserved in humans were retained as a resource.

plasmids, specifically when a starting plasmid library is not Gateway compatible.

## Discussion

Here, we developed a method named CRISPRmass, and to our knowledge, this is the first time that CRISPR/Cas9 has been used in high-throughput plasmid library construction. In addition, by using CRISPRmass, we generated a total of 5551 UAS-cDNA/ORF constructs, covering 83% of the 6687 conserved genes in humans

in the DGRC Gold Collection, and among them, 5518 were generated within 3 mo by three people. The 5551 UAS-cDNA/ORF plasmids of *Drosophila* genes serve as a resource for GOF screening in cultured cells and for generation of transgenic UAS-cDNA/ORF library in *Drosophila*.

Compared to the conventional or recombinational (Gateway) cloning technique, CRISPRmass offers several advantages in generation of a genome-wide UAS-cDNA/ORF plasmid library. First, it offers modularization. We introduced a new concept of modular assembly design and manipulation of vector backbone for construction of a genome-wide library. A combinatory use of CRISPR/Cas9 and Gibson assembly allows editing shared sequence in vector backbone, therefore it modularizes library construction. Once it is constructed for a given vector, a UAS module can fit any cDNA or ORF clones as long as these clones are in the same vector, which avoids individualized design and manipulation for every single cDNA or ORF clone. Another benefit of modularization is that CRISPRmass is also suitable for generating UAS-cDNA/ORF constructs from long cDNAs or ORFs, because CRISPRmass does not manipulate the cDNAs or ORFs themselves but vector backbones. In contrast, construction of UAS-cDNA/ORF plasmids of long cDNAs or ORFs by the conventional or recombinational (Gateway) cloning techniques is difficult. Second, it offers simplicity. The construction of a UAS-cDNA/ORF library from existing cDNA/ORF resources is uniformed and simplified as two steps of test tube reactions followed by bacterial transformation and analysis of only one colony by restriction digestion. For hundreds of thousands of genes, these procedures can be carried out in parallel. The remaining manipulations do not require primer design and synthesis, PCR, gel purification, sequencing, or any advanced experimental techniques (Table 1). Third, it offers efficiency. Introduction of an antibiotic resistance gene into a UAS module enables positive selection of UAS-cDNA/ORF clones. Moreover, Gibson assembly filters out any potential off-target DNA cleavages caused by CRISPR/Cas9, because it requires that both 5' and 3' ends of a Cas9-cleaved vector overlap with those of a UAS module respectively. Fourth, it offers adaptability. A UAS module can fit any vector by modifying a core UAS module, and the expression levels of genes under the control of UAS can be easily modulated by adjusting the UAS copy number.

The core concept of CRISPRmass is to manipulate shared vector sequences for high-throughput plasmid library editing.

**Table 1.** Comparison of construction of a UAS-cDNA/ORF library (5551 plasmids, six vector backbones) from the *Drosophila* DGC Gold Collection by recombinational cloning and CRISPRmass

		Recombinational cloning (Gateway)	CRISPRmass
For generating entry constructs	Number of entry constructs generated	5551	Not applicable
	Pairs of primers designed for PCR amplification of every single cDNA/ORF	Minimum 5551	Not applicable
	PCR reactions for every single cDNA/ORF	Minimum 5551	Not applicable
	Samples for agarose gel electrophoresis for every single cDNA/ORF	Minimum 5551	Not applicable
	Gel purifications for every single cDNA/ORF	Minimum 5551	Not applicable
	Efficiency of BP reaction	> 90% <sup>a</sup>	Not applicable
	Colonies to be analyzed for entry constructs	More than 5551	Not applicable
	Sequencing reactions for every single cDNA/ORF	More than 5551	Not applicable
	ORF-specific primers for sequencing	Required for >2-kb ORF	Not applicable
	For generating UAS-cDNA/ORF constructs	sgRNAs used	Not applicable
UAS modules constructed from a core UAS module		Not applicable	6
Colonies to be analyzed for UAS-cDNA/ORF constructs		5551	5551
Efficiency		> 95% <sup>b</sup>	98.6% <sup>c</sup>
Selection strategy of UAS-cDNA/ORF constructs		Negative selection	Positive selection
Key reactions		LR reaction	CRISPR/Cas9 digestion, Gibson assembly
Materials for key reactions		LR Clonase <sup>d</sup>	Cas9, <sup>e</sup> Gibson assembly master mix <sup>f</sup>
Cost of materials for key reactions		115,203 CNY <sup>g</sup>	45,290 CNY <sup>h</sup>

<sup>a</sup>Based on Gateway Recombination Cloning brochure (Thermo Fisher Scientific; <https://www.thermofisher.com/cn/zh/home/life-science/cloning/gateway-cloning.html>).

<sup>b</sup>This is the LR reaction efficiency based on Gateway Recombination Cloning brochure (Thermo Fisher Scientific; <https://www.thermofisher.com/cn/zh/home/life-science/cloning/gateway-cloning.html>).

<sup>c</sup>Based on this study in which UAS modules were inserted into 98.6% of 5987cDNA/ORF plasmids successfully under the condition of randomly analyzing only one colony for each UAS-cDNA/ORF construct.

<sup>d</sup>LR Clonase (Thermo Fisher Scientific 11791100; <https://www.thermofisher.com/order/catalog/product/11791100?SID=srch-srp-11791100>).

<sup>e</sup>SpCas9 (GenScript Z03386-50; <https://www.genscript.com.cn/search?q=Z03386&search=Search>).

<sup>f</sup>Gibson assembly master mix (NEB E2621X; <http://www.neb-china.com/pshow.asp?id=3196>).

<sup>g</sup>Each LR reaction volume is 1.4  $\mu$ L.

<sup>h</sup>Each CRISPR/Cas9 digestion volume is 2.4  $\mu$ L, and each Gibson assembly reaction volume is 1.4  $\mu$ L.

Theoretically, the first step of CRISPRmass manipulation, which is cleavage of shared vector sequences of library plasmids by CRISPR/Cas9, could be altered to other methods such as zinc-finger nuclease (ZFN) (Kim et al. 1996; Smith et al. 2000) and transcription activator-like effector nuclease (TALEN) (Christian et al. 2010). The second step of CRISPRmass manipulation, which is insertion of modules by Gibson assembly, could be altered to other seamless cloning methods such as cotransformation cloning (Bubeck et al. 1993), seamless ligation cloning extract (SLiCE) (Zhang et al. 2012), ligation-independent cloning (LIC) (Aslanidis and de Jong 1990), sequence- and ligation-independent cloning (SLIC) (Li and Elledge 2007), In-Fusion cloning (Zhu et al. 2007), and T5 exonuclease-dependent DNA assembly (TEDA) (Xia et al. 2019).

In addition to generating UAS-cDNA/ORF plasmid libraries, CRISPRmass has broader applicability to constructing some other genome-scale plasmid libraries, for example, constructing mammalian cDNA/ORF expression libraries by CRISPRmass-assisted insertion of a CMV promoter into upstream of cDNA/ORFs, which is equivalent to transferring cDNA/ORFs into a CMV promoter-containing expression vector. Likewise, CRISPRmass can also be used for high-throughput transferring clone collections that are not compatible with Gateway into the Gateway entry vectors, followed by LR reactions to produce desired destination constructs when multiple different expression vectors are likely to be desirable. Thus, CRISPRmass can be applied to editing various genome-wide plasmid libraries in general and is an alternative to Gateway technology in high-throughput plasmid library editing, paving the way for global studies of the biological networks.

## Methods

### cDNA and ORF clones

The *Drosophila*, mouse, and human cDNA and ORF clones used in this study are from the *Drosophila* Genome Resource Center (<https://dgrc.bio.indiana.edu>) Gold Collection, the mouse Mammalian Gene Collection (MGC, <https://genecollections.nci.nih.gov/MGC/>), and the human CCSB-Broad Lentiviral Expression Library (Yang et al. 2011).

### *Drosophila* strains

GAL4 lines used in this study are the motor neuron driver *OK6-GAL4* (Aberle et al. 2002), eye driver *GMR-GAL4* (Freeman 1996), wing driver *A9-GAL4* (BDSC Stock 8761), and muscle driver *MEF2-GAL4* (BDSC Stock 27390). The UAS-cDNA/ORF transgenic flies were generated using phiC31-mediated integration into the *attP2* landing site, and the UAS-cDNA/ORF constructs used for transgenes were described in Supplemental Table S1.

### sgRNA preparation

To design the sgRNAs used for cleavage of vector backbones of cDNA/ORF plasmids, we used Optimized CRISPR Design (<http://crispr.mit.edu>) to search vector backbone sequences for sgRNA targets. The vector backbone sequences used for sgRNA design are adjacent to the 5' end of cDNAs or ORFs, thus all cDNA/ORF plasmids with an identical vector backbone can be uniformly digested by *Streptococcus pyogenes* Cas9 together with the same



sgRNA. The target candidates of sgRNAs for vector backbones of cDNA/ORF clones are listed in Supplemental Table S2.

To prepare a DNA template for in vitro transcription (IVT) of sgRNA, we synthesize two PAGE-purified oligos, forward primer [5'-TAATACGACTCACTATAGG(N)<sub>20</sub>GTTTATAGAGCTAGAAATAG-3'], where (N)<sub>20</sub> is the target sequence of a sgRNA, and a sgRNA scaffold reverse primer sgRNA-REV (5'-AAAAGCACCGACTCGGTGCCA CTT-3'). Next a 100- $\mu$ L PCR reaction is set up by mixing 5  $\mu$ L template pX330 (Addgene plasmid 42230, a gift from Feng Zhang) (Bischof et al. 2013) (1 ng/ $\mu$ L), 5  $\mu$ L forward primer (10  $\mu$ M), 5  $\mu$ L sgRNA-REV (10  $\mu$ M), 50  $\mu$ L Q5 Hot Start High-Fidelity 2 $\times$  Master Mix (NEB), and 35  $\mu$ L DEPC-treated water (Invitrogen) in a PCR tube. A PCR program on an ABI 2720 PCR Thermal Cycler is run as follows: 30 sec at 98°C; five cycles of 10 sec at 98°C, 10 sec at 51°C, and 5 sec at 72°C; 30 cycles of 10 sec at 98°C and 15 sec at 72°C; and a final extension of 2 min at 72°C. The PCR product is separated by 0.8% agarose gel electrophoresis and a ~120-bp DNA fragment should be visible on the gel. The ~120-bp DNA fragment is then purified by QIAquick Gel Extraction Kit (Qiagen). The gel-purified ~120-bp DNA fragment serves as a template for IVT of sgRNA.

IVT of sgRNA is performed using HiScribe T7 Quick High Yield RNA Synthesis Kit (NEB). A 10- $\mu$ L IVT reaction is set up as follows: 330 ng ~120-bp gel-purified DNA fragment, 3.33  $\mu$ L NTP Buffer Mix, 0.67  $\mu$ L T7 RNA Polymerase Mix, and DEPC-treated water. The 10- $\mu$ L IVT reaction is incubated for 4 h at 37°C, then the in vitro transcribed sgRNA, without purification, is quantified, diluted to 20 ng/ $\mu$ L with DEPC-treated water, aliquoted, stored at -80°C, and is ready for CRISPRmass.

### sgRNA evaluation

To evaluate sgRNAs targeting vector backbone sequences that are adjacent to the 5' end of cDNAs or ORFs in cDNA/ORF plasmids, we performed in vitro CRISPR/Cas9 cleavage analysis of DNA substrates. A 5- $\mu$ L CRISPR/Cas9 cleavage reaction contains 0.2  $\mu$ L 1.22  $\mu$ M *S. pyogenes* Cas9 (GenScript Z03386-50), 0.5  $\mu$ L 20 ng/ $\mu$ L sgRNA, 0.015 pmol DNA substrate, 0.5  $\mu$ L 10 $\times$  Cas9 Buffer, and DEPC-treated water. After the 5- $\mu$ L cleavage reaction is incubated for 1 h at 37°C, cleavage fragments are separated by 0.8% agarose gel electrophoresis and analyzed by Tanon 2500 Gel Imaging System (Tanon). A DNA fragment containing the sgRNA targeting sequences of a vector backbone is released by restriction digestion of a cDNA/ORF plasmid and used as a DNA substrate. DNA substrates and the sizes of substrate cleavage fragments are listed in Supplemental Table S2. The sgRNAs selected for CRISPRmass and their targets on vector backbones of cDNA/ORF clones are listed in Supplemental Table S3.

### Construction of UAS modules

#### UAS module for *pOT2*

A 1141-bp EcoRI/HindIII fragment carrying ampicillin resistance gene (*Amp*) was PCR amplified from pBluescript KS (+) using EcoRI-pOT2G4-Amp-F and HindIII-Ascl-Amp-R primers, digested with EcoRI/HindIII, inserted into the EcoRI/HindIII sites of pMartini vector (Wang et al. 2012), resulting in pMartini-Amp. pMartini-Amp was verified by sequencing.

A 938-bp HindIII/EcoRI fragment carrying *attB-UAS-Hsp70* was PCR amplified from DopR-Tango (a gift from David Anderson) (Inagaki et al. 2012) using HindIII-attB-F and EcoRI-pOT2G4-Hsp70-R primers, digested with EcoRI/HindIII, inserted into the EcoRI/HindIII sites of pBluescript KS (+), resulting in pBS-attB-UAS-Hsp70. pBS-attB-UAS-Hsp70 was verified by sequencing.

A 1127-bp EcoRI/HindIII fragment of pMartini-Amp and a 924-bp attB-UAS-Hsp70 EcoRI/HindIII fragment of pBS-attB-UAS-Hsp70 were ligated together into the EcoRI site of pCR8GW-TOPO vector (Invitrogen), resulting in pCR8GW-Amp-attB-UAS-Hsp70.

A 4127-bp *mini-white* Ascl fragment of pJFRC150-20XUAS-IVS-Flp1::PEST (Addgene plasmid 32132, a gift from Gerald Rubin) (Nern et al. 2011) was inserted into the Ascl site of pCR8GW-Amp-attB-UAS-Hsp70, giving rise to pCR8GW-Amp-W-attB-UAS-Hsp70. The 6178-bp *Amp-W-attB-UAS-Hsp70* fragment was released by EcoRI from pCR8GW-Amp-W-attB-UAS-Hsp70, and this fragment serves as a UAS module for construction of UAS-cDNA/ORF plasmids from pOT2 vector-based cDNA/ORF clones.

#### UAS module for *pFLC-I*

A 866-bp EcoRI/HindIII fragment carrying chloramphenicol resistance gene (*Cam*) was PCR amplified from pOT2-CG4968 using EcoRI-pFLCG3-Cam-F and HindIII-Ascl-Cam-R2 primers, digested with EcoRI/HindIII, inserted into the EcoRI/HindIII sites of pMartini vector, resulting in pMartini-Cam. pMartini-Cam was verified by sequencing.

A 940-bp HindIII/EcoRI fragment carrying *attB-UAS-Hsp70* was PCR amplified from DopR-Tango using HindIII-attB-F and EcoRI-pFLCG3-Hsp70-R primers, digested with EcoRI/HindIII, inserted into the EcoRI/HindIII sites of pBluescript KS (+), resulting in pBS-attB-UAS-Hsp70-2. pBS-attB-UAS-Hsp70-2 was verified by sequencing.

A 852-bp EcoRI/HindIII fragment of pMartini-Cam and a 926-bp attB-UAS-Hsp70 EcoRI/HindIII fragment of pBS-attB-UAS-Hsp70-2 were ligated together into the EcoRI site of pCR8GW-TOPO vector, resulting in pCR8GW-Cam-attB-UAS-Hsp70.

A 4127-bp *mini-white* Ascl fragment of pJFRC150-20XUAS-IVS-Flp1::PEST was inserted into the Ascl site of pCR8GW-Cam-attB-UAS-Hsp70, giving rise to pCR8GW-Cam-W-attB-UAS-Hsp70. The 5905-bp Cam-W-attB-UAS-Hsp70 fragment was released by EcoRI from pCR8GW-Cam-W-attB-UAS-Hsp70, and this fragment serves as a UAS module for construction of UAS-cDNA/ORF plasmids from pFLC-I vector-based cDNA/ORF clones.

#### UAS module for *pBS SK(-)*

A 1805-bp *Cam-attB-UAS-Hsp70* EcoRI fragment was PCR amplified from pCR8GW-Cam-attB-UAS-Hsp70 using EcoRI-pBSG3-Cam-F and EcoRI-pBSG3-hsp70-R primers, digested with EcoRI, inserted into the EcoRI site of pCR8GW-TOPO vector, resulting in pCR8GW-Cam-attB-UAS-Hsp70-2. pCR8GW-Cam-attB-UAS-Hsp70-2 was verified by sequencing.

A 4127-bp *mini-white* Ascl fragment of pJFRC150-20XUAS-IVS-Flp1::PEST was inserted into the Ascl site of pCR8GW-Cam-attB-UAS-Hsp70-2, giving rise to pCR8GW-Cam-W-attB-UAS-Hsp70-2. The 5918-bp Cam-W-attB-UAS-Hsp70-2 fragment was released by EcoRI from pCR8GW-Cam-W-attB-UAS-Hsp70-2, and this fragment serves as a UAS module for construction of UAS-cDNA/ORF plasmids from pBS SK(-) vector-based cDNA/ORF clones.

#### UAS module for *pOTB7* and *pOTB7\_Drill*

A 2070-bp *Amp-attB-UAS-Hsp70* EcoRI fragment was PCR amplified from pCR8GW-Amp-attB-UAS-Hsp70 using EcoRI-pOTB7G9-Amp-F and EcoRI-pOTB7G9-hsp70-R primers, digested with EcoRI, inserted into the EcoRI site of pCR8GW-TOPO vector (Wang et al. 2012), resulting in pCR8GW-Amp-attB-UAS-Hsp70-2. pCR8GW-Amp-attB-UAS-Hsp70-2 was verified by sequencing.

A 4127-bp *mini-white* *Ascl* fragment of pJFRC150-20X UAS-IVS-Flp1::PEST was inserted into the *Ascl* site of pCR8GW-Amp-attB-UAS-Hsp70-2, giving rise to pCR8GW-Amp-W-attB-UAS-Hsp70-2. The 6183-bp Amp-W-attB-UAS-Hsp70 fragment was released by *EcoRI* from pCR8GW-Amp-W-attB-UAS-Hsp70-2, and this fragment serves as a UAS module for construction of UAS-cDNA/ORF plasmids from pOTB7 and pOTB7\_DraIII vectors-based cDNA/ORF clones.

#### UAS module for pCR2.1

A 1805-bp *Cam-attB-UAS-Hsp70* *EcoRI* fragment was PCR amplified from pCR8GW-Cam-attB-UAS-Hsp70 using *EcoRI*-pCR2.1G1-Cam-F and *EcoRI*-pCR2.1G1-hsp70-R primers, digested with *EcoRI*, inserted into the *EcoRI* site of pCR8GW-TOPO vector, resulting in pCR8GW-Cam-attB-UAS-Hsp70-3. pCR8GW-Cam-attB-UAS-Hsp70-3 was verified by sequencing.

A 4127-bp *mini-white* *Ascl* fragment of pJFRC150-20X UAS-IVS-Flp1::PEST was inserted into the *Ascl* site of pCR8GW-Cam-attB-UAS-Hsp70-3, giving rise to pCR8GW-Cam-W-attB-UAS-Hsp70-3. The 5918-bp Cam-W-attB-UAS-Hsp70-3 fragment was released by *EcoRI* from pCR8GW-Cam-W-attB-UAS-Hsp70-3, and this fragment serves as a UAS module for construction of UAS-cDNA/ORF plasmids from pCR2.1 vector-based cDNA/ORF clones.

#### UAS module for pDNR-Dual

A 1805-bp *Cam-attB-UAS-Hsp70* *EcoRI* fragment was PCR amplified from pCR8GW-Cam-attB-UAS-Hsp70 using *EcoRI*-pLX304OriG11-Cam-F and *EcoRI*-pLX304OriG11-hsp70-R primers, digested with *EcoRI*, inserted into the *EcoRI* site of pCR8GW-TOPO vector, resulting in pCR8GW-Cam-attB-UAS-Hsp70-4. pCR8GW-Cam-attB-UAS-Hsp70-4 was verified by sequencing.

A 4127-bp *mini-white* *Ascl* fragment of pJFRC150-20X UAS-IVS-Flp1::PEST was inserted into the *Ascl* site of pCR8GW-Cam-attB-UAS-Hsp70-4, giving rise to pCR8GW-Cam-W-attB-UAS-Hsp70-4. The 5918-bp Cam-W-attB-UAS-Hsp70-4 fragment was released by *EcoRI* from pCR8GW-Cam-W-attB-UAS-Hsp70-4, and this fragment serves as a UAS module for construction of UAS-cDNA/ORF plasmids from pDNR-Dual vector-based cDNA/ORF clones.

#### UAS module for pLX304

A 1800-bp *Cam-attB-UAS-Hsp70* *EcoRI* fragment was PCR amplified from pCR8GW-Cam-attB-UAS-Hsp70 using *EcoRI*-pLX304OriG13-Cam-F and *EcoRI*-pLX304CMVG7-hsp70-R primers, digested with *EcoRI*, inserted into the *EcoRI* site of pCR8GW-TOPO vector, resulting in pCR8GW-Cam-attB-UAS-Hsp70-5. pCR8GW-Cam-attB-UAS-Hsp70-5 was verified by sequencing.

A 4127-bp *mini-white* *Ascl* fragment of pJFRC150-20X UAS-IVS-Flp1::PEST was inserted into the *Ascl* site of pCR8GW-Cam-attB-UAS-Hsp70-5, giving rise to pCR8GW-Cam-W-attB-UAS-Hsp70-5. The 5913-bp Cam-W-attB-UAS-Hsp70-5 fragment was released by *EcoRI* from pCR8GW-Cam-W-attB-UAS-Hsp70-5, and this fragment serves as a UAS module for construction of UAS-cDNA/ORF plasmids from pLX304 vector-based cDNA/ORF clones.

#### UAS module for pCMV-SPORT6

A 1800-bp *Cam-attB-UAS-Hsp70* *EcoRI* fragment was PCR amplified from pCR8GW-Cam-attB-UAS-Hsp70 using *EcoRI*-pLX304OriG13-Cam-F and *EcoRI*-pCMVST6G15-hsp70-R primers, digested with *EcoRI*, inserted into the *EcoRI* site of

pCR8GW-TOPO vector, resulting in pCR8GW-Cam-attB-UAS-Hsp70-6. pCR8GW-Cam-attB-UAS-Hsp70-6 was verified by sequencing.

A 4127-bp *mini-white* *Ascl* fragment of pJFRC150-20X UAS-IVS-Flp1::PEST was inserted into the *Ascl* site of pCR8GW-Cam-attB-UAS-Hsp70-5, giving rise to pCR8GW-Cam-W-attB-UAS-Hsp70-6. The 5918-bp Cam-W-attB-UAS-Hsp70-6 fragment was released by *EcoRI* from pCR8GW-Cam-W-attB-UAS-Hsp70-6, and this fragment serves as a UAS module for construction of UAS-cDNA/ORF plasmids from pCMV-SPORT6 vector-based cDNA/ORF clones.

Primers used for constructing and sequencing UAS modules are listed in Supplemental Table S7.

#### Construction of UAS-cDNA/ORF plasmids by CRISPRmass

Cas9 cleavage of cDNA or ORF clones in pOT2, pOTB7, pOTB7\_DraIII, pFLC-I, pBS SK(-), pCR2.1, and pDNR-Dual vectors is performed as follows: 0.4  $\mu$ L of 0.019  $\mu$ M cDNA or ORF plasmid is cleaved for 1 h at 37°C in a 2.4  $\mu$ L of reaction containing 0.16  $\mu$ L of 1.22  $\mu$ M of Cas9 (GenScript), 0.4  $\mu$ L of 20 ng/ $\mu$ L sgRNA, and 0.24  $\mu$ L 10 $\times$  Cas9 Buffer. cDNA/ORF clones in pLX304 and pCMV-SPORT6 are cleaved twice by Cas9 together with two different sgRNAs to remove unnecessary sequences (Supplemental Fig. S1). Then 0.7  $\mu$ L Cas9-cleaved cDNA/ORF plasmid, without purification, is directly subjected to Gibson assembly together with 0.07  $\mu$ L of 0.006  $\mu$ M UAS module and 0.7  $\mu$ L of NEBuilder HiFi DNA Assembly Master Mix (NEB). Gibson assembly is performed for 1 h at 50°C, then 0.7  $\mu$ L of Gibson assembly product is directly used to transform *E. coli*. Transformants are selected on LB plates containing antibiotic corresponding to the antibiotic resistance gene on the UAS module.

#### Construction of 2 $\times$ MYC or 3 $\times$ MYC-tagged UAS-cDNA/ORF plasmids

A 3 $\times$ MYC epitope sequence (EQKLISEEDLEQKLISEEDLEQKLI SEEDL) was inserted right before the stop codon of a representative *Drosophila* cDNA/ORF clone for each of the six vectors of the DGRC Gold Collection using Q5 Site-Directed Mutagenesis Kit (NEB). Briefly, a 5- $\mu$ L PCR reaction was set up as follows: 0.2 ng cDNA/ORF plasmid, 0.5  $\mu$ M forward primer (Supplemental Table S8), 0.5  $\mu$ M reverse primer (Supplemental Table S8), and 2.5  $\mu$ L Q5 Hot Start High-Fidelity 2 $\times$  Master Mix (NEB). A PCR program on an ABI 2720 PCR Thermal Cycler was performed run as follows: 30 sec at 98°C; five cycles of 10 sec at 98°C, 10 sec at 51°C, and 8 min at 72°C; 30 cycles of 10 sec at 98°C and 8 min at 72°C; and a final extension of 5 min at 72°C. Then 0.2  $\mu$ L of the PCR product is subjected to a KLD reaction together with 0.2  $\mu$ L of 10 $\times$  KLD Enzyme Mix, 1  $\mu$ L of 2 $\times$  KLD Reaction Buffer, and 0.6  $\mu$ L ddH<sub>2</sub>O. KLD reactions were performed for 10 min at room temperature, and the KLD reaction products were used to transform *E. coli*. The resulting plasmids were confirmed by sequencing (Supplemental Table S8). Similarly, a 3 $\times$ MYC epitope sequence was inserted right before the stop codon of the mouse *Ctbp2* cDNA on pCMV-SPORT6 using Q5 Site-Directed Mutagenesis Kit, but subsequent sequencing indicated that a copy of MYC tag was removed from the 3 $\times$ MYC epitope, resulting in a 2 $\times$ MYC-tagged *Ctbp2* plasmid (Supplemental Table S8). 2 $\times$ MYC or 3 $\times$ MYC-tagged cDNA/ORF plasmids and all the primers used for 2 $\times$ MYC or 3 $\times$ MYC epitope tagging are listed in Supplemental Table S8.

#### Construction of 3 $\times$ FLAG-tagged UAS-cDNA plasmids

A 3 $\times$ FLAG epitope sequence (DYKDHDGDYKDHDIDYKDDDDK) was inserted right before the stop codon of the 23 cDNA clones

of the DGRC Gold Collection using Q5 Site-Directed Mutagenesis Kit (NEB). Briefly, a 5- $\mu$ L PCR reaction was set up as follows: 0.2 ng cDNA plasmid, 0.5  $\mu$ M forward primer, 0.5  $\mu$ M reverse primer, and 2.5  $\mu$ L Q5 Hot Start High-Fidelity 2 $\times$  Master Mix (NEB). A PCR program on an ABI 2720 PCR Thermal Cycler was performed run as follows: 30 sec at 98°C; five cycles of 10 sec at 98°C, 10 sec at 51°C or 56°C and 8 min at 72°C; 20 cycles of 10 sec at 98°C, 10 sec at 70°C and 8 min at 72°C; and a final extension of 5 min at 72°C. Then 0.2  $\mu$ L of the PCR product is subjected to a KLD reaction together with 0.2  $\mu$ L of 10 $\times$  KLD Enzyme Mix, 1  $\mu$ L of 2 $\times$  KLD Reaction Buffer, and 0.6  $\mu$ L ddH<sub>2</sub>O. KLD reactions were performed for 10 min at room temperature, and the KLD reaction products were used to transform *E. coli*. The resulting plasmids were confirmed by sequencing. The *Nhe3* gene was fused with a 3xFLAG tag at its N terminus by Gibson assembly. Briefly, because the SphI site overlaps the *Nhe3* start codon, the *Nhe3* cDNA clone was linearized by SphI, resulting in a 6792-bp fragment. The overlapping 3FLAG-*Nhe3*-F and 3FLAG-*Nhe3*-R primers were annealed for 1 min at 50°C and then extended for 2 min at 72°C, resulting in a 102-bp fragment containing a 3xFLAG sequence and partial sequence of *Nhe3* cDNA. The 6792-bp and 102-bp fragments were then joined through Gibson assembly using NEBuilder HiFi DNA Assembly Master Mix (NEB), and the resulting 3xFLAG-tagged *Nhe3* plasmid, pFLC-3FLAG-*Nhe3*, was confirmed by sequencing. All the 3xFLAG-tagged UAS-cDNA plasmids were constructed by CRISPRmass based on the 3xFLAG-tagged cDNA plasmids, and all the 3xFLAG-tagged cDNA and UAS-cDNA plasmids and the primers used for 3xFLAG epitope tagging and sequencing are listed in Supplemental Tables S5 and S6.

### Western blotting and immunohistochemistry

These two methods are available in Supplemental Methods.

### Data access

All processed sequencing data generated in this study have been submitted to the NCBI GenBank (<https://www.ncbi.nlm.nih.gov/genbank/>) under accession numbers MN503478–MN503501 and MN518567–MN518687. The GenBank accession numbers MN503478–MN503501 are for the cDNA/ORFs of 24 UAS-cDNA/ORF constructs from the more than 5500 UAS-cDNA/ORF constructs generated by CRISPRmass. The GenBank accession numbers MN518567–MN518687 are for the 121 UAS-cDNA/ORF constructs that were verified by sequencing from the more than 5500 UAS-cDNA/ORF constructs. The plasmids are available upon request.

### Competing interest statement

Two patents have been filed on this study (China patent application Nos. 2019101104852 and 2019101110552).

### Acknowledgments

We thank the *Drosophila* Genomics Resource Center (DGRC), supported by National Institutes of Health grant 2P40OD010949-10A1, for the DGRC Gold Collection. We thank the Developmental Studies Hybridoma Bank for antibodies and the Bloomington Stock Center for the *Drosophila* stocks. We thank the Core Facility of *Drosophila* Resource and Technology, Shanghai Institute of Biochemistry and Cell Biology, Chinese Academy of Sciences for *Drosophila* transgenesis. We thank Jihui Guo, Yan Wan, Xueyan Song, and Hongshuo Wang for their assis-

tance on the project. This work was sponsored by the National Natural Science Foundation of China (31471010 and 31401013), Shanghai Pujiang Program (14PJ1405900), and Natural Science Foundation of Shanghai (19ZR1446400).

**Author contributions:** J.W. conceived and supervised this project. J.W. and P.W. designed the experiments. J.W., P.W., and W.X. performed the experiments. Y.Z. and G.N. provided suggestions. J.W. and P.W. analyzed the data and wrote the manuscript.

### References

- Aberle H, Haghighi AP, Fetter RD, McCabe BD, Magalhães TR, Goodman CS. 2002. *wishful thinking* encodes a BMP type II receptor that regulates synaptic growth in *Drosophila*. *Neuron* **33**: 545–558. doi:10.1016/S0896-6273(02)00589-5
- Aslanidis C, de Jong PJ. 1990. Ligation-independent cloning of PCR products (LIC-PCR). *Nucleic Acids Res* **18**: 6069–6074. doi:10.1093/nar/18.20.6069
- Bischof J, Bjorklund M, Furger E, Schertel C, Taipale J, Basler K. 2013. A versatile platform for creating a comprehensive UAS-ORFeome library in *Drosophila*. *Development* **140**: 2434–2442. doi:10.1242/dev.088757
- Brand AH, Perrimon N. 1993. Targeted gene expression as a means of altering cell fates and generating dominant phenotypes. *Development* **118**: 401–415.
- Bubeck P, Winkler M, Bautsch W. 1993. Rapid cloning by homologous recombination *in vivo*. *Nucleic Acids Res* **21**: 3601–3602. doi:10.1093/nar/21.15.3601
- Christian M, Cermak T, Doyle EL, Schmidt C, Zhang F, Hummel A, Bogdanove AJ, Voytas DF. 2010. Targeting DNA double-strand breaks with TAL effector nucleases. *Genetics* **186**: 757–761. doi:10.1534/genetics.110.120717
- Dietzl G, Chen D, Schnorrer F, Su KC, Barinova Y, Fellner M, Gasser B, Kinsey K, Oettel S, Scheiblauer S, et al. 2007. A genome-wide transgenic RNAi library for conditional gene inactivation in *Drosophila*. *Nature* **448**: 151–156. doi:10.1038/nature05954
- Ewen-Campen B, Yang-Zhou D, Fernandes VR, González DP, Liu LP, Tao R, Ren X, Sun J, Hu Y, Zirin J, et al. 2017. Optimized strategy for *in vivo* Cas9-activation in *Drosophila*. *Proc Natl Acad Sci* **114**: 9409–9414. doi:10.1073/pnas.1707635114
- Freeman M. 1996. Reiterative use of the EGF receptor triggers differentiation of all cell types in the *Drosophila* eye. *Cell* **87**: 651–660. doi:10.1016/S0092-8674(00)81385-9
- Gibson DG, Young L, Chuang RY, Venter JC, Hutchison CA 3rd, Smith HO. 2009. Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nat Methods* **6**: 343–345. doi:10.1038/nmeth.1318
- Inagaki HK, Ben-Tabou de-Leon S, Wong AM, Jagadish S, Ishimoto H, Barnea G, Kitamoto T, Axel R, Anderson DJ. 2012. Visualizing neuromodulation *in vivo*: TANGO-mapping of dopamine signaling reveals appetite control of sugar sensing. *Cell* **148**: 583–595. doi:10.1016/j.cell.2011.12.022
- Jia Y, Xu RG, Ren X, Ewen-Campen B, Rajakumar R, Zirin J, Yang-Zhou D, Zhu R, Wang F, Mao D, et al. 2018. Next-generation CRISPR/Cas9 transcriptional activation in *Drosophila* using flySAM. *Proc Natl Acad Sci* **115**: 4719–4724. doi:10.1073/pnas.1800677115
- Kim YG, Cha J, Chandrasegaran S. 1996. Hybrid restriction enzymes: zinc finger fusions to Fok I cleavage domain. *Proc Natl Acad Sci* **93**: 1156–1160. doi:10.1073/pnas.93.3.1156
- Li MZ, Elledge SJ. 2007. Harnessing homologous recombination *in vitro* to generate recombinant DNA via SLIC. *Nat Methods* **4**: 251–256. doi:10.1038/nmeth1010
- Lin MF, Carlson JW, Crosby MA, Matthews BB, Yu C, Park S, Wan KH, Schroeder AJ, Gramates LS, St Pierre SE, et al. 2007. Revisiting the protein-coding gene catalog of *Drosophila melanogaster* using 12 fly genomes. *Genome Res* **17**: 1823–1836. doi:10.1101/gr.6679507
- Mata J, Curado S, Ephrussi A, Rørth P. 2000. Tribbles coordinates mitosis and morphogenesis in *Drosophila* by regulating string/CDC25 proteolysis. *Cell* **101**: 511–522. doi:10.1016/S0092-8674(00)80861-2
- Miklos GL, Rubin GM. 1996. The role of the genome project in determining gene function: insights from model organisms. *Cell* **86**: 521–529. doi:10.1016/S0092-8674(00)80126-9
- Nern A, Pfeiffer BD, Svoboda K, Rubin GM. 2011. Multiple new site-specific recombinases for use in manipulating animal genomes. *Proc Natl Acad Sci* **108**: 14198–14203. doi:10.1073/pnas.1111704108
- Perkins LA, Holderbaum L, Tao R, Hu Y, Sopko R, McCall K, Yang-Zhou D, Flockhart I, Binari R, Shim HS, et al. 2015. The transgenic RNAi project



- at Harvard Medical School: resources and validation. *Genetics* **201**: 843–852. doi:10.1534/genetics.115.180208
- Potter CJ, Tasic B, Russler EV, Liang L, Luo L. 2010. The Q system: a repressible binary system for transgene expression, lineage tracing, and mosaic analysis. *Cell* **141**: 536–548. doi:10.1016/j.cell.2010.02.025
- Rørth P. 1996. A modular misexpression screen in *Drosophila* detecting tissue-specific phenotypes. *Proc Natl Acad Sci* **93**: 12418–12422. doi:10.1073/pnas.93.22.12418
- Rørth P, Szabo K, Bailey A, Laverty T, Rehm J, Rubin GM, Weigmann K, Milán M, Benes V, Ansoorge W, et al. 1998. Systematic gain-of-function genetics in *Drosophila*. *Development* **125**: 1049–1057.
- Rubin GM, Hong L, Brokstein P, Evans-Holm M, Frise E, Stapleton M, Harvey DA. 2000. A *Drosophila* complementary DNA resource. *Science* **287**: 2222–2224. doi:10.1126/science.287.5461.2222
- Smith J, Bibikova M, Whitby FG, Reddy AR, Chandrasegaran S, Carroll D. 2000. Requirements for double-strand cleavage by chimeric restriction enzymes with zinc finger DNA-recognition domains. *Nucleic Acids Res* **28**: 3361–3369. doi:10.1093/nar/28.17.3361
- Stapleton M, Liao G, Brokstein P, Hong L, Carninci P, Shiraki T, Hayashizaki Y, Champe M, Pacleb J, Wan K, et al. 2002. The *Drosophila* gene collection: identification of putative full-length cDNAs for 70% of *D. melanogaster* genes. *Genome Res* **12**: 1294–1300. doi:10.1101/gr.269102
- St Johnston D. 2002. The art and design of genetic screens: *Drosophila melanogaster*. *Nat Rev Genet* **3**: 176–188. doi:10.1038/nrg751
- Tsou WL, Sheedlo MJ, Morrow ME, Blount JR, McGregor KM, Das C, Todi SV. 2012. Systematic analysis of the physiological importance of deubiquitinating enzymes. *PLoS One* **7**: e43112. doi:10.1371/journal.pone.0043112
- Wang JW, Beck ES, McCabe BD. 2012. A modular toolset for recombination transgenesis and neurogenetic analysis of *Drosophila*. *PLoS One* **7**: e42102. doi:10.1371/journal.pone.0042102
- Xia Y, Li K, Li J, Wang T, Gu L, Xun L. 2019. T5 exonuclease-dependent assembly offers a low-cost method for efficient cloning and site-directed mutagenesis. *Nucleic Acids Res* **47**: e15. doi:10.1093/nar/gky1169
- Xu R, Deng K, Zhu Y, Wu Y, Ren J, Wan M, Zhao S, Wu X, Han M, Zhuang Y, et al. 2008. A large-scale functional approach to uncover human genes and pathways in *Drosophila*. *Cell Res* **18**: 1114–1127. doi:10.1038/cr.2008.295
- Yang X, Boehm JS, Yang X, Salehi-Ashtiani K, Hao T, Shen Y, Lubonja R, Thomas SR, Alkan O, Bhimdi T, et al. 2011. A public genome-scale lentiviral expression library of human ORFs. *Nat Methods* **8**: 659–661. doi:10.1038/nmeth.1638
- Zdobnov EM, Tegenfeldt F, Kuznetsov D, Waterhouse RM, Simão FA, Ioannidis P, Seppely M, Loetscher A, Kriventseva EV. 2017. OrthoDB v9.1: cataloging evolutionary and functional annotations for animal, fungal, plant, archaeal, bacterial and viral orthologs. *Nucleic Acids Res* **45**: D744–D749. doi:10.1093/nar/gkw1119
- Zhang Y, Werling U, Edelman W. 2012. SLiCE: a novel bacterial cell extract-based DNA cloning method. *Nucleic Acids Res* **40**: e55. doi:10.1093/nar/gkr1288
- Zhu B, Cai G, Hall EO, Freeman GJ. 2007. In-fusion assembly: seamless engineering of multidomain fusion proteins, modular vectors, and mutations. *Biotechniques* **43**: 354–359. doi:10.2144/000112536

Received March 23, 2019; accepted in revised form November 8, 2019.



## CRISPR-based modular assembly of a UAS-cDNA/ORF plasmid library for more than 5500 *Drosophila* genes conserved in humans

Ping Wei, Wen Xue, Yun Zhao, et al.

*Genome Res.* 2020 30: 95-106 originally published online November 13, 2019

Access the most recent version at doi:[10.1101/gr.250811.119](https://doi.org/10.1101/gr.250811.119)

---

**Supplemental Material** <http://genome.cshlp.org/content/suppl/2019/12/14/gr.250811.119.DC1>

**References** This article cites 34 articles, 13 of which can be accessed free at:  
<http://genome.cshlp.org/content/30/1/95.full.html#ref-list-1>

**Creative Commons License** This article is distributed exclusively by Cold Spring Harbor Laboratory Press for the first six months after the full-issue publication date (see <http://genome.cshlp.org/site/misc/terms.xhtml>). After six months, it is available under a Creative Commons License (Attribution-NonCommercial 4.0 International), as described at <http://creativecommons.org/licenses/by-nc/4.0/>.

**Email Alerting Service** Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).

---



---

To subscribe to *Genome Research* go to:  
<https://genome.cshlp.org/subscriptions>

---