



CONTENTS

Bioinformatics and Databases

- TERRA ONTseq: a long-read-based sequencing pipeline to study the human telomeric transcriptome 955^{OA}
Joana Rodrigues, Roberta Alfieri, Silvia Bione, and Claus M. Azzalin

RNA and Gene Expression

- Human T-cell leukemia virus type 1 uses a specific tRNA^{Pro} isodecoder to prime reverse transcription 967^{OA}
Yu-Ci Syu, Joshua Hatterschide, Christina R. Budding, Yingke Tang, and Karin Musier-Forsyth

- Suppression of the *Escherichia coli rnpA49* conditionally lethal phenotype by different compensatory mutations 977^{OA}
Arianne M. Babina, Leif A. Kirsebom, and Dan I. Andersson

- Magnesium ions mitigate metastable states in the regulatory landscape of mRNA elements 992^{OA}
Erdong Ding, Susmit Narayan Chaudhury, Jigneshkumar Dahyabhai Prajapati, José N. Onuchic, and Karissa Y. Sanbonmatsu

- The essential role of architectural noncoding RNA *Neat1* in cold-induced beige adipocyte differentiation in mice 1011
Hikaru Toya, Yuko Okamatsu-Ogura, Saori Yokoi, Misuzu Kurihara, Mari Mito, Shintaro Iwasaki, Tetsuro Hirose, and Shinichi Nakagawa

- The modification landscape of *Pseudomonas aeruginosa* tRNAs 1025
Mariana D. Mandl, Siddhardha S. Maligireddy, Wilfried M. Guiblet, Christina M. Fitzsimmons, Kayla S. McDonald, Delayna L. Warrell, and Pedro J. Batista

- A novel reporter for helicase activity in translation uncovers DDX3X interactions 1041^{OA}
Kevin C. Wilkins, Till Schroeder, Sohyun Gu, Jezrael L. Revalde, and Stephen N. Floor

(continued)

Cover Illustration: Human argonaute microRNA complex (PDB id: 8D71; Xiao Y, Liu TM, MacRae IJ. 2023. A tiny loop in the Argonaute PIWI domain tunes small RNA seed strength. EMBO Rep **24**: e55806). Argonaute proteins use microRNAs (miRNAs) and small interfering RNAs (siRNAs) as guides to regulate gene expression in plants and animals. Human argonaute is shown in this structure to recognize a short region (6–8 nt) of miRNA. microRNA backbone is displayed in NDB colors: A—red, G—green, U—cyan; argonaute protein is displayed as a wheat-colored ribbon. Missing RNA and protein segments are indicated by dotted lines. Cover image provided by the Nucleic Acid Knowledgebase (nakb.org). Image generated using ChimeraX (Protein Sci **30**: 70–82).

Contents (continued)

Functional analysis of the zinc finger modules of the <i>Saccharomyces cerevisiae</i> splicing factor Luc7 Tucker J. Carrocci, Samuel DeMario, Kevin He, Natalie J. Zeps, Cade T. Harkner, Guillaume F. Chanfreau, and Aaron A. Hoskins	1058
Selected humanization of yeast U1 snRNP leads to global suppression of pre-mRNA splicing and mitochondrial dysfunction in the budding yeast Subbaiah Chalivendra, Shasha Shi, Xueni Li, Zhiling Kuang, Joseph Giovinazzo, Lingdi Zhang, John Rossi, Jingxin Wang, Anthony J. Saviola, Robb Welty, Shiheng Liu, Katherine F. Vaeth, Z. Hong Zhou, Kirk C. Hansen, J. Matthew Taliaferro, and Rui Zhao	1070
Transcription factors ER α and Sox2 have differing multiphasic DNA- and RNA-binding mechanisms Wayne O. Hemphill, Halley R. Steiner, Jackson R. Kominsky, Deborah S. Wuttke, and Thomas R. Cech	1089 ^{OA}
RNA: Instructions for contributors	1106

^{OA}Open Access paper