

Table 1.22
U.S. Prevalence Counts, Invasive Cancers Only, January 1, 2010^a
Using Different Tumor Inclusion Criteria^b

| Site | Sex | 5-Year Limited Duration | | | 35-year Limited Duration | |
|--------------------------------|------------|--------------------------------------|--|---|--------------------------------------|--|
| | | 1st Invasive Tumor Ever ^c | 1st Per Site in Previous 35 Years ^d | 1st Per Site in Previous 5 Years ^e | 1st Invasive Tumor Ever ^c | 1st Per Site in Previous 35 Years ^d |
| All Sites | Both Sexes | 4,511,266 | 4,598,344 | 5,044,168 | 12,600,627 | 12,810,577 |
| | Male | 2,332,050 | 2,366,640 | 2,584,260 | 5,990,343 | 6,060,178 |
| | Female | 2,179,216 | 2,231,704 | 2,459,908 | 6,610,284 | 6,750,399 |
| Oral Cavity & Pharynx | Both Sexes | 102,019 | 117,112 | 121,258 | 264,620 | 291,343 |
| | Male | 71,123 | 80,929 | 83,544 | 175,045 | 190,951 |
| | Female | 30,896 | 36,183 | 37,714 | 89,575 | 100,392 |
| Esophagus | Both Sexes | 21,268 | 25,973 | 26,092 | 33,702 | 39,594 |
| | Male | 16,808 | 20,399 | 20,447 | 26,317 | 30,715 |
| | Female | 4,460 | 5,574 | 5,645 | 7,385 | 8,879 |
| Stomach | Both Sexes | 35,420 | 42,280 | 42,609 | 70,821 | 81,142 |
| | Male | 21,286 | 25,765 | 25,901 | 40,560 | 46,844 |
| | Female | 14,134 | 16,515 | 16,708 | 30,261 | 34,298 |
| Colon & Rectum | Both Sexes | 407,833 | 471,251 | 480,046 | 1,129,666 | 1,258,620 |
| | Male | 207,302 | 239,824 | 244,132 | 561,245 | 622,211 |
| | Female | 200,531 | 231,427 | 235,914 | 568,421 | 636,409 |
| Liver & Intrahepatic Bile Duct | Both Sexes | 30,577 | 34,834 | 34,867 | 40,940 | 45,832 |
| | Male | 22,266 | 25,232 | 25,265 | 28,891 | 32,221 |
| | Female | 8,311 | 9,602 | 9,602 | 12,049 | 13,611 |
| Pancreas | Both Sexes | 31,442 | 38,275 | 38,275 | 41,284 | 49,174 |
| | Male | 15,733 | 19,467 | 19,467 | 20,155 | 24,367 |
| | Female | 15,709 | 18,808 | 18,808 | 21,129 | 24,807 |
| Larynx | Both Sexes | 30,947 | 37,152 | 37,518 | 87,135 | 98,063 |
| | Male | 25,107 | 30,036 | 30,356 | 70,117 | 78,586 |
| | Female | 5,840 | 7,116 | 7,162 | 17,018 | 19,477 |
| Lung & Bronchus | Both Sexes | 231,996 | 296,283 | 304,764 | 391,564 | 480,133 |
| | Male | 108,211 | 140,119 | 143,487 | 178,912 | 220,472 |
| | Female | 123,785 | 156,164 | 161,277 | 212,652 | 259,661 |
| Melanoma of the Skin | Both Sexes | 276,312 | 312,334 | 325,566 | 884,259 | 953,581 |
| | Male | 148,749 | 171,783 | 180,200 | 440,298 | 480,917 |
| | Female | 127,563 | 140,551 | 145,366 | 443,961 | 472,664 |
| Breast | Female | 842,193 | 909,424 | 965,835 | 2,770,060 | 2,934,536 |
| Cervix | Female | 38,963 | 40,783 | 40,907 | 208,065 | 213,819 |
| Corpus & Uterus, NOS | Female | 169,126 | 189,983 | 190,106 | 569,448 | 620,803 |
| Ovary ^f | Female | 57,207 | 66,063 | 66,139 | 169,398 | 190,511 |

^a U.S. 2010 cancer prevalence counts are based on 2010 cancer prevalence proportions from the SEER 9 registries and 1/1/2010 U.S. population estimates based on the average of 2009 and 2010 population estimates from the U.S. Bureau of the Census.

^b Prevalence estimates are ambiguous for those with multiple cancers, unless the tumor inclusion criteria are understood. Depending on the application, different inclusion criteria may be appropriate. This table provides three different methods of tumor inclusion:

^c (c) First invasive tumor ever

^d (d) First invasive tumor for each cancer site diagnosed during the previous 35 years (1975-2009)

^e (e) First invasive tumor for each cancer site diagnosed during the previous 5 years (2005-2009)

For definitions (d) and (e) all sites is treated as a separate cancer "site".

Consider a woman who had three invasive cancers: Melanoma in 1981; Breast cancer in 2005; Melanoma in 2006.

In method (c) the melanoma is the woman's first cancer, and is counted for the melanoma and all sites 35-year limited duration prevalence. For 5-year limited duration prevalence, the woman is not counted at all since her first cancer occurred more than 5 years prior to 1/1/2010.

In method (d) the 1981 melanoma is counted for the melanoma and all sites 35-year limited duration prevalence. The 2005 breast cancer is counted for the breast 5-year and 35-year limited duration prevalence.

In method (e) the 2005 breast cancer is counted for the breast cancer and all sites 5-year limited duration prevalence. The 2006 melanoma is counted for 5-year limited duration prevalence for melanoma.

^f Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

Table 1.22 - continued
 U.S. Prevalence Counts, Invasive Cancers Only, January 1, 2010^a
 Using Different Tumor Inclusion Criteria^b

| Site | Sex | 5-Year Limited Duration | | | 35-year Limited Duration | |
|----------------------------|------------|--------------------------------------|--|---|--------------------------------------|--|
| | | 1st Invasive Tumor Ever ^c | 1st Per Site in Previous 35 Years ^d | 1st Per Site in Previous 5 Years ^e | 1st Invasive Tumor Ever ^c | 1st Per Site in Previous 35 Years ^d |
| Prostate | Male | 1,026,391 | 1,107,903 | 1,107,977 | 2,616,190 | 2,787,286 |
| Testis | Male | 42,302 | 43,012 | 43,616 | 205,130 | 207,844 |
| Urinary Bladder | Both Sexes | 198,998 | 248,244 | 253,143 | 551,278 | 641,106 |
| | Male | 151,489 | 190,075 | 194,009 | 411,638 | 479,012 |
| | Female | 47,509 | 58,169 | 59,134 | 139,640 | 162,094 |
| Kidney & Renal Pelvis | Both Sexes | 145,766 | 177,066 | 179,051 | 330,450 | 385,370 |
| | Male | 88,557 | 108,930 | 110,317 | 196,335 | 230,734 |
| | Female | 57,209 | 68,136 | 68,734 | 134,115 | 154,636 |
| Brain & Nervous System | Both Sexes | 43,995 | 46,883 | 47,345 | 128,063 | 132,164 |
| | Male | 24,044 | 25,790 | 26,033 | 68,158 | 70,466 |
| | Female | 19,951 | 21,093 | 21,312 | 59,905 | 61,698 |
| Thyroid | Both Sexes | 165,253 | 183,524 | 184,093 | 494,969 | 528,131 |
| | Male | 37,076 | 42,905 | 43,058 | 107,747 | 117,569 |
| | Female | 128,177 | 140,619 | 141,035 | 387,222 | 410,562 |
| Hodgkin Lymphoma | Both Sexes | 39,048 | 41,589 | 41,611 | 168,309 | 172,937 |
| | Male | 21,122 | 22,557 | 22,568 | 87,084 | 89,522 |
| | Female | 17,926 | 19,032 | 19,043 | 81,225 | 83,415 |
| Non-Hodgkin Lymphoma | Both Sexes | 198,946 | 233,585 | 235,457 | 499,712 | 558,340 |
| | Male | 105,553 | 125,451 | 126,442 | 261,560 | 292,961 |
| | Female | 93,393 | 108,134 | 109,015 | 238,152 | 265,379 |
| Myeloma | Both Sexes | 49,451 | 58,292 | 58,328 | 77,360 | 88,490 |
| | Male | 26,622 | 32,063 | 32,087 | 42,092 | 48,802 |
| | Female | 22,829 | 26,229 | 26,241 | 35,268 | 39,688 |
| Leukemia | Both Sexes | 112,120 | 129,298 | 129,528 | 282,150 | 310,046 |
| | Male | 64,915 | 75,612 | 75,711 | 160,039 | 176,801 |
| | Female | 47,205 | 53,686 | 53,817 | 122,111 | 133,245 |
| Acute Lymphocytic Leukemia | Both Sexes | 16,205 | 16,686 | 16,686 | 66,030 | 66,759 |
| | Male | 9,215 | 9,443 | 9,443 | 36,418 | 36,708 |
| | Female | 6,990 | 7,243 | 7,243 | 29,612 | 30,051 |
| Childhood (Ages 0-19) | Both Sexes | 63,904 | 63,977 | 64,503 | 304,247 | 304,716 |
| | Male | 33,948 | 33,981 | 34,250 | 156,819 | 157,032 |
| | Female | 29,956 | 29,996 | 30,253 | 147,428 | 147,684 |
| Kaposi Sarcoma | Both Sexes | 7,319 | 7,750 | 7,750 | 24,815 | 25,977 |
| | Male | 6,865 | 7,208 | 7,208 | 23,533 | 24,518 |
| | Female | 454 | 542 | 542 | 1,282 | 1,459 |
| Mesothelioma | Both Sexes | 2,968 | 3,926 | 3,938 | 4,538 | 5,658 |
| | Male | 1,998 | 2,761 | 2,773 | 2,601 | 3,467 |
| | Female | 970 | 1,165 | 1,165 | 1,937 | 2,191 |

^a U.S. 2010 cancer prevalence counts are based on 2010 cancer prevalence proportions from the SEER 9 registries and 1/1/2010 U.S. population estimates based on the average of 2009 and 2010 population estimates from the U.S. Bureau of the Census.

^b Prevalence estimates are ambiguous for those with multiple cancers, unless the tumor inclusion criteria are understood. Depending on the application, different inclusion criteria may be appropriate. This table provides three different methods of tumor inclusion:

^c (c) First invasive tumor ever
^d (d) First invasive tumor for each cancer site diagnosed during the previous 35 years (1975-2009)
^e (e) First invasive tumor for each cancer site diagnosed during the previous 5 years (2005-2009)
 For definitions (d) and (e) all sites is treated as a separate cancer "site".

Consider a woman who had three invasive cancers: Melanoma in 1981; Breast cancer in 2005; Melanoma in 2006.

In method (c) the melanoma is the woman's first cancer, and is counted for the melanoma and all sites 35-year limited duration prevalence. For 5-year limited duration prevalence, the woman is not counted at all since her first cancer occurred more than 5 years prior to 1/1/2010. In method (d) the 1981 melanoma is counted for the melanoma and all sites 35-year limited duration prevalence. The 2005 breast cancer is counted for the breast 5-year and 35-year limited duration prevalence. In method (e) the 2005 breast cancer is counted for the breast cancer and all sites 5-year limited duration prevalence. The 2006 melanoma is counted for 5-year limited duration prevalence for melanoma.

Table 1.23
U.S. Complete Prevalence Counts, Invasive Cancers Only, January 1, 2010^a
By Age at Prevalence

| Site/Sex | Age at Prevalence | | | | | | | | |
|-----------------------|-----------------------|--------|--------|--------|---------|---------|-----------|-----------|-----------|
| | All Ages ^c | 0-9 | 10-19 | 20-29 | 30-39 | 40-49 | 50-59 | 60-69 | 70+ |
| All Sites | | | | | | | | | |
| Males | 6,078,974 | 18,390 | 42,087 | 83,873 | 155,608 | 354,801 | 860,143 | 1,578,142 | 2,985,931 |
| Females | 6,948,940 | 16,117 | 36,351 | 86,197 | 226,859 | 636,726 | 1,276,932 | 1,684,808 | 2,984,950 |
| Oral Cavity & Pharynx | | | | | | | | | |
| Males | 181,084 | 68 | 468 | 1,480 | 3,518 | 14,224 | 43,847 | 54,509 | 62,971 |
| Females | 94,109 | 90 | 553 | 1,470 | 3,635 | 8,681 | 17,691 | 22,821 | 39,167 |
| Esophagus | | | | | | | | | |
| Males | 26,388 | 0 | 0 | 22 | 219 | 1,177 | 4,653 | 9,419 | 10,897 |
| Females | 7,451 | 0 | 0 | 0 | 28 | 256 | 1,263 | 1,790 | 4,114 |
| Stomach | | | | | | | | | |
| Males | 41,386 | 0 | 18 | 64 | 607 | 2,589 | 7,095 | 10,242 | 20,773 |
| Females | 30,883 | 4 | 12 | 165 | 568 | 2,307 | 4,398 | 6,312 | 17,118 |
| Colon & Rectum | | | | | | | | | |
| Males | 569,930 | 11 | 117 | 1,190 | 5,632 | 25,029 | 80,675 | 136,002 | 321,273 |
| Females | 584,551 | 0 | 75 | 1,251 | 5,467 | 23,433 | 67,578 | 110,244 | 376,504 |
| Liver & Intrahep | | | | | | | | | |
| Males | 28,967 | 484 | 535 | 554 | 454 | 1,582 | 10,000 | 9,083 | 6,274 |
| Females | 12,437 | 471 | 470 | 312 | 411 | 895 | 2,803 | 3,028 | 4,048 |
| Pancreas | | | | | | | | | |
| Males | 20,293 | 11 | 12 | 127 | 376 | 1,616 | 4,320 | 6,183 | 7,649 |
| Females | 21,316 | 0 | 66 | 155 | 413 | 1,376 | 3,843 | 5,483 | 9,980 |
| Larynx | | | | | | | | | |
| Males | 71,659 | 0 | 0 | 78 | 243 | 1,915 | 10,556 | 20,739 | 38,129 |
| Females | 17,370 | 0 | 0 | 25 | 134 | 1,011 | 3,256 | 4,677 | 8,267 |
| Lung & Bronchus | | | | | | | | | |
| Males | 183,410 | 45 | 90 | 444 | 1,184 | 5,545 | 24,951 | 54,888 | 96,262 |
| Females | 216,021 | 31 | 71 | 396 | 1,435 | 7,883 | 28,898 | 59,047 | 118,261 |
| Melanoma of the Skin | | | | | | | | | |
| Males | 452,540 | 78 | 656 | 5,014 | 18,294 | 46,766 | 95,157 | 122,709 | 163,865 |
| Females | 469,240 | 113 | 993 | 10,894 | 33,972 | 73,514 | 109,961 | 106,450 | 133,343 |

^a U.S. 2010 cancer prevalence counts are based on 2010 cancer prevalence proportions from the SEER 9 registries (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta) and 1/1/2010 U.S. population estimates based on the average of 2009 and 2010 population estimates from the U.S. Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^b Cases diagnosed more than 35 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^c Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.

Table 1.23 - continued
 U.S. Complete Prevalence Counts, Invasive Cancers Only, January 1, 2010^a
 By Age at Prevalence

| Site/Sex | Age at Prevalence | | | | | | | | |
|------------------------|-----------------------|-------|--------|--------|--------|---------|---------|---------|-----------|
| | All Ages ^c | 0-9 | 10-19 | 20-29 | 30-39 | 40-49 | 50-59 | 60-69 | 70+ |
| Breast | | | | | | | | | |
| Males | 14,588 | 0 | 0 | 11 | 89 | 586 | 2,018 | 4,133 | 7,751 |
| Females | 2,829,041 | 0 | 53 | 2,740 | 36,291 | 216,912 | 539,229 | 767,846 | 1,265,969 |
| Cervix | | | | | | | | | |
| Females | 249,496 | 0 | 45 | 2,239 | 16,405 | 42,748 | 59,766 | 55,248 | 73,044 |
| Corpus & Uterus, NOS | | | | | | | | | |
| Females | 600,346 | 0 | 47 | 530 | 5,666 | 24,621 | 87,167 | 157,833 | 324,482 |
| Ovary ^d | | | | | | | | | |
| Females | 186,138 | 84 | 1,018 | 3,453 | 6,883 | 18,736 | 40,700 | 47,632 | 67,632 |
| Prostate | | | | | | | | | |
| Males | 2,617,682 | 34 | 59 | 101 | 324 | 21,033 | 228,561 | 740,428 | 1,627,144 |
| Urinary Bladder | | | | | | | | | |
| Males | 419,648 | 45 | 105 | 611 | 2,438 | 11,535 | 42,898 | 99,644 | 262,372 |
| Females | 143,992 | 23 | 57 | 217 | 964 | 4,266 | 13,698 | 30,030 | 94,737 |
| Kidney & Renal Pelvis | | | | | | | | | |
| Males | 202,516 | 1,454 | 2,505 | 2,659 | 5,172 | 17,901 | 40,060 | 57,511 | 75,253 |
| Females | 138,989 | 1,343 | 2,681 | 2,851 | 4,901 | 12,071 | 23,973 | 33,716 | 57,452 |
| Hodgkin Lymphoma | | | | | | | | | |
| Males | 93,890 | 181 | 2,315 | 9,272 | 16,487 | 24,007 | 21,438 | 13,274 | 6,916 |
| Females | 88,038 | 101 | 1,901 | 9,617 | 16,530 | 23,444 | 18,638 | 10,651 | 7,157 |
| Non-Hodgkin Lymphoma | | | | | | | | | |
| Males | 266,478 | 773 | 3,513 | 7,582 | 13,366 | 29,224 | 52,496 | 67,493 | 92,032 |
| Females | 242,587 | 430 | 1,847 | 4,348 | 8,879 | 20,902 | 41,030 | 56,723 | 108,427 |
| Myeloma | | | | | | | | | |
| Males | 42,185 | 0 | 6 | 84 | 442 | 2,859 | 8,141 | 13,608 | 17,045 |
| Females | 35,432 | 0 | 0 | 39 | 281 | 2,266 | 6,273 | 10,676 | 15,896 |
| Leukemia | | | | | | | | | |
| Males | 162,651 | 6,545 | 12,947 | 12,388 | 10,928 | 13,500 | 22,064 | 32,827 | 51,452 |
| Females | 125,312 | 5,438 | 10,680 | 10,750 | 9,514 | 9,728 | 14,680 | 20,623 | 43,899 |
| Acute Lymphocytic Leuk | | | | | | | | | |
| Males | 38,239 | 5,461 | 11,081 | 9,597 | 6,075 | 3,568 | 1,238 | 775 | 444 |
| Females | 31,328 | 4,527 | 9,014 | 7,859 | 5,163 | 2,755 | 884 | 717 | 408 |

^a U.S. 2010 cancer prevalence counts are based on 2010 cancer prevalence proportions from the SEER 9 registries (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta) and 1/1/2010 U.S. population estimates based on the average of 2009 and 2010 population estimates from the U.S. Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^b Cases diagnosed more than 35 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^c Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.