

1 **IPCC Reasons for Concern regarding climate change risks**

2 Brian C. O'Neill,<sup>1\*</sup> Michael Oppenheimer,<sup>2</sup> Rachel Warren,<sup>3</sup> Stephane Hallegatte,<sup>4</sup> Robert E. Kopp,<sup>5</sup> Hans  
3 O. Pörtner,<sup>6</sup> Robert Scholes,<sup>7</sup> Joern Birkmann,<sup>8</sup> Wendy Foden,<sup>9</sup> Rachel Licker,<sup>2</sup> Katharine J. Mach,<sup>10</sup>  
4 Phillippe Marbaix,<sup>11</sup> Michael Mastrandrea,<sup>10</sup> Jeff Price,<sup>3</sup> Kiyoshi Takahashi,<sup>12</sup> Jean-Pascal van Ypersele<sup>11</sup>  
5 and Gary Yohe<sup>13</sup>

6 <sup>1</sup> National Center for Atmospheric Research, Boulder, CO 80305, USA.

7 <sup>2</sup> Department of Geosciences and the Woodrow Wilson School of Public and International Affairs,  
8 Princeton University, Princeton NJ 08544, USA.

9 <sup>3</sup> Tyndall Centre for Climate Change, School of Environmental Sciences, University of East Anglia,  
10 Norwich, UK.

11 <sup>4</sup> Climate Change Group, World Bank, Washington, DC 20433, USA.

12 <sup>5</sup> Department of Earth and Planetary Sciences and Rutgers Energy Institute, Rutgers University, New  
13 Brunswick, NJ 08901, USA.

14 <sup>6</sup> Marine Biology/Ecological and Evolutionary Physiology, Alfred-Wegener-Institute, D-27570  
15 Bremerhaven, Germany.

16 <sup>7</sup> University of the Witwatersrand, Johannesburg, South Africa.

17 <sup>8</sup> Institute of Spatial and Regional Planning, University of Stuttgart, 70569 Stuttgart, Germany.

18 <sup>9</sup> Department of Botany and Zoology, University of Stellenbosch, Stellenbosch, South Africa.

19 <sup>10</sup> Carnegie Institution for Science, Department of Global Ecology, Stanford, CA 94305, USA.

20 <sup>11</sup> Earth and Life Institute, Université catholique de Louvain, B-1348 Louvain-la-Neuve, Belgium.

21 <sup>12</sup> National Institute for Environmental Studies, Tsukuba, Japan.

22 <sup>13</sup> Department of Economics, Wesleyan University, Middletown, CT 06459, USA.

23 \* email: boneill@ucar.edu

24

25

1 **The Reasons for Concern (RFC) framework communicates scientific understanding about risks in**  
2 **relation to varying levels of climate change. The framework, now a cornerstone of the IPCC**  
3 **assessments, aggregates global risks into five categories as a function of global mean temperature**  
4 **change (GMT). We review the RFC's conceptual basis and the risk judgments made in the most recent**  
5 **IPCC report, confirming those judgments in most cases in the light of more recent literature and**  
6 **identifying their limitations. We point to extensions of the framework that offer complementary**  
7 **climate change metrics to GMT and better account for possible changes in social and ecological**  
8 **system vulnerability. Further research should systematically evaluate risks under alternative scenarios**  
9 **of future climatic and societal conditions.**

10 The RFC framework was developed in the IPCC Third Assessment Report (TAR) to inform discussions  
11 relevant to implementation of Article 2 of the UN Framework Convention on Climate Change (UNFCCC).  
12 Article 2 presents the Convention's long-term objective of avoiding "dangerous anthropogenic  
13 interference with the climate system." The RFC framework and the associated "Burning Embers"  
14 diagram illustrating authors' risk judgments have since been widely discussed and used to inform policy  
15 decisions. For example, they informed a recent dialog between Parties to the UNFCCC and experts<sup>1,2</sup> on  
16 the adequacy of the long-term goal of avoiding a warming of 2°C relative to pre-industrial, contributing  
17 to a strengthening of that goal in the recent Paris Agreement<sup>3</sup>. Elaborations of the Burning Embers have  
18 been used to represent climate impacts and risks at the regional level<sup>4</sup> and for specific systems (e.g.,  
19 ocean systems<sup>5</sup>).

20 This article reviews the conceptual basis for the RFCs (Box 1) and offers an explanation of the reasoning  
21 behind associated risk judgments that is complementary to, but goes beyond, the treatment in the IPCC  
22 Fifth Assessment Report<sup>6</sup>. We focus explicitly on the evidence base for transitions from one risk level to  
23 the next, incorporate post-AR5 literature in those discussions, and offer thoughts about limitations of  
24 the subjective judgments behind each RFC. We also improved the synthesis of RFC-related material  
25 across AR5, and in turn provide both a clearer connection to evidence from AR5 that supports the RFC  
26 judgments, as well as a comparison of the RFCs to similar approaches employing metrics other than  
27 GMT for characterizing risk. Perhaps most importantly, we consider improvements in the framework,  
28 particularly emphasizing the dynamic nature of exposure and vulnerability, two key components of risk  
29 not sufficiently covered in the current approach.

---

30 **TEXT BOX 1: Conceptual Basis**

31 The Reasons for Concern (RFCs) reported in AR5 are:

- 32 1. *risks to unique and threatened systems (indicated by RFC1 below);*
- 33 2. *risks associated with extreme weather events (RFC2);*
- 34 3. *risks associated with the distribution of impacts (RFC3);*
- 35 4. *risks associated with global aggregate impacts (RFC4); and*
- 36 5. *risks associated with large-scale singular events (RFC5).*

37 Types of risk included in each category are discussed in the next section. The categories share an  
38 emphasis on going beyond changes in biophysical systems to possible consequences for society and  
39 ecosystems, including their interdependencies (henceforth "socio-ecological systems"). *Risk* is the

1 potential for negative consequences, whereas *impacts* are the manifestation of that potential<sup>7, 8</sup>.  
2 Climate-related risk depends on the probability of hazardous events or trends and on the consequences  
3 manifested when a physical, climate-related hazard interacts with the exposure and vulnerability of  
4 society and ecosystems. *Hazards* related to climate change include altered occurrence of extreme  
5 events, trends in precipitation or temperature, sea level rise, ocean acidification, deoxygenation or  
6 ocean circulation changes. *Exposure* is the presence of people, ecosystems, or assets in places and  
7 settings that could be adversely affected, and *vulnerability* is their susceptibility and predisposition to  
8 harm<sup>9, 10</sup>. These definitions follow the choices laid out in AR5, although alternatives can be found in the  
9 literature<sup>11</sup>.

10 The process of making judgments about levels of risk for each RFC (Supplementary Text 1) was  
11 underpinned by the identification of “key risks”. Key risks reflect potentially severe adverse  
12 consequences for socio-ecological systems that could be used to inform the interpretation of  
13 “dangerous” in the UNFCCC Article 2 objective. Criteria for identifying key risks include<sup>6, 12, 13</sup>.

- 14 (1) high probability of significant risk materializing, taking into account its timing;
- 15 (2) large magnitude of associated consequences, taking into account the importance of affected  
16 systems;
- 17 (3) persistent vulnerability or exposure contributing to risks, or the irreversibility, at least on human  
18 timescales, of associated impacts; and
- 19 (4) limited potential to reduce risks through adaptation or mitigation.

20 AR5 authors drew on these criteria to characterize climate-related risk for each RFC as a function of  
21 GMT as Undetectable, Moderate, High, or Very High. The transition from Undetectable to Moderate is  
22 defined by the GMT at which there is at least *medium confidence* that impacts associated with a given  
23 RFC are both detectable and attributable to climate change (based on the analysis in ref. 14, 18.6.4\*),  
24 while also accounting for the magnitude of the risk and the other criteria noted above. The transition  
25 from Moderate to High risk is assigned to the GMT at which associated impacts become severe and  
26 widespread. The transition from High to Very High is set at the GMT at which risk is high according to all  
27 criteria and in particular the ability to adapt is limited. In each case, variations in regional climate  
28 outcomes for a given GMT are accounted for and the likelihood of the associated hazardous event or  
29 trend is judged.

30 Defining the risk levels this way enables integration within each RFC across different but related risks  
31 and many different types of evidence. The scale is inherently nonlinear and qualitative, even if  
32 quantified evidence enters the judgments.

33

## 34 **REASONS FOR CONCERN**

35 Risk judgments for each RFC are based on the key risk criteria (Box 1) but the relative importance of  
36 each varies across RFCs depending on the quality and quantity of information available in the literature.

---

\* To be more explicit about the source of information in chapters of IPCC reports, we provide specific sub-sections or figure/table numbers following the citation number.

1 It is also not possible to rely on a single quantitative metric of risk for a given RFC since each one  
2 aggregates over a number of different risks. An enhanced Burning Embers diagram (Fig. 1) summarizes  
3 the evidence, indicating both individual risks that played important roles in identifying particular risk  
4 transitions, as well as overarching key risks relevant in broader terms to each RFC (Table 1). These  
5 overarching key risk categories were developed in AR5<sup>6</sup> from risks identified as being of high concern by  
6 chapter authors from across IPCC Working Group (WG) II (Supplementary Text 1). Unless otherwise  
7 specified, we refer to GMT relative to pre-industrial (1850-1900). Note that conversions from units used  
8 in AR5 can give the appearance of overly precise temperature levels (Supplementary Text 2).

9

## 10 **RFC1: Risks to unique and threatened systems**

11 Unique and threatened systems encompass ecological and human systems that (1) have restricted  
12 geographic ranges that are constrained by climate-related conditions, and (2) have high endemism or  
13 other distinctive properties. Many of these systems also face exceptional human-driven threats.  
14 Examples include tropical glacier systems, coral reefs, mangrove ecosystems, biodiversity hotspots<sup>15</sup>,  
15 and unique indigenous communities<sup>16</sup>.

16 AR5 located the transition from Undetectable to Moderate risk below recent temperatures based on the  
17 detection and attribution (with at least *medium confidence*) of impacts on Arctic, mountain, and warm-  
18 water coral reef systems (ref. 14, 18.6.4), with indirect support from impacts on other systems  
19 (Supplementary Text 3.1). In the Arctic, impacts include the observed decline in sea ice extent<sup>17</sup>,  
20 warming and thawing of permafrost in Alaska and associated land-sliding<sup>14, 18, 19</sup>, substantial changes in  
21 ecosystems and ecological dynamics, including signs of broad-scale boreal forest encroachment into  
22 tundra<sup>20, 21, 22</sup>, and livelihood impacts on indigenous Arctic peoples<sup>14</sup>. In mountain systems, there is  
23 evidence of shrinking or receding glaciers from all continents<sup>14</sup>. There is also *high confidence* that  
24 climate change has contributed to widespread and frequent coral bleaching and mortality due to high  
25 temperatures<sup>23, 24, 25, 26</sup>.

26 A transition from Moderate to High risk occurs over the range ~1.1-1.6 °C above pre-industrial. In broad  
27 terms, this transition is placed halfway between the Undetectable-to-Moderate transition and High-to-  
28 Very High transition (discussed next) to reflect the generally increasing risks over this range. However,  
29 specific projected impacts for Arctic and coral reef systems also informed the judgment (Supplementary  
30 Text 3.2).

31 A transition to Very High risk is located around 2.6°C to reflect very high risks and limited ability to adapt  
32 for a wide range of unique and threatened ecosystems<sup>27, 28</sup> (Supplementary Text 3.3.1). Substantial  
33 impacts to unique and threatened systems are projected at or even below this level of warming<sup>29, 30</sup>.  
34 These systems include both major ecoregions and biodiversity hotspots containing unique (including  
35 endemic) and threatened systems. They include the Cerrado in South America, the Fynbos and  
36 Succulent Karoo ecoregions in South Africa, Australian rainforest ecoregions, the Caribbean, Indo-  
37 Burma, Mediterranean Basin, Southwest Australia, and the Tropical Andes<sup>30, 31, 32</sup>. Risks to Arctic, coral  
38 reef, and mountain systems also escalate above this level of warming (Supplementary Text 3.3.2). For  
39 example, large-scale coral reef dissolution may occur if CO<sub>2</sub> concentrations reach approximately 560

1 ppm due to the combined effects of warming and ocean acidification (ref. 33, 5.4.2.4; ref. 25), consistent  
2 with a warming of approximately 2.5°C (ref. 34, Fig. 3 and Table 2).

3 More comprehensive impact assessments are needed that consider more fully the human dimensions of  
4 impacts on unique and threatened systems. Most projections of impacts on species and ecosystems fail  
5 to consider how adaptation may ameliorate or exacerbate existing pressures and threats and introduce  
6 new ones<sup>35</sup> (Supplementary Text 3.3.3; see Box 2 for general description of adaptation level assumed in  
7 RFCs). Also, whether species will be able adapt or move fast enough to keep up with their changing  
8 environments will be crucial to the resilience of ecological systems<sup>36</sup> but remains poorly studied<sup>37</sup>.

9 **RFC2: Risks associated with extreme weather events**

10 RFC2 encompasses risk to human health, livelihoods, assets, and ecosystems from extremes such as  
11 heat waves, heavy rain, drought and associated wildfires, and coastal flooding.

12 The transition from Undetectable to Moderate risk is located at recent temperatures based primarily on  
13 evidence for the detection and attribution of impacts of extreme events on coral reefs and human  
14 health (Supplementary Text 4.1). Bleaching of warm-water corals has resulted from periods of elevated  
15 near-surface ocean temperature to levels attributed to climate change (ref. 38; ref. 14, 18.6.4, Table 18-  
16 10; ref. 6, 19.6.3.2). For human health impacts, there has been detection and attribution of mortality  
17 impacts of temperature extremes in some regions (ref. 39, 11.4.1). Additional support for this transition  
18 comes from the detection and attribution of extreme heat and precipitation events, including post-AR5  
19 analyses at the global scale<sup>40</sup>, along with the widespread occurrence of high vulnerability and exposure  
20 and abnormal levels of mortality in some events<sup>41</sup>.

21 The transition to High risk is located at ~1.6°C, relying primarily on projections of large, near-term  
22 changes in the magnitude and likelihood of extremes of temperature and precipitation. The choice is  
23 somewhat subjective due to the paucity of literature projecting the impacts of changes in heat  
24 extremes. By about 2035 (during which time the increase in model- and scenario- averaged GMT  
25 remains below ~1.6°C), 25-30% of daily maximum temperatures are projected to exceed the historical  
26 (1961-1990) 90th percentile value (ref. 42, Fig. 11-17). Duration, intensity and spatial extent of heat  
27 waves and warm spells also increase in the near term. We chose 2035 as a benchmark for the transition  
28 to high risk because the potential impacts from changes in temperature extremes are large and AR5  
29 indicates such changes are likely<sup>42</sup>. Furthermore, there is high confidence in projected mean changes  
30 through 2035 because they are not strongly dependent on future emissions. In addition, on average,  
31 the frequency and intensity of heavy precipitation events over land will likely increase over much of the  
32 world (Supplementary Text 4.2). A reduction in return period for historical once-in-20-yr precipitation  
33 events globally (land only) to about once-in-14-yr or less by 2046-65 is also expected<sup>43</sup>.

34 A key limitation is that changing exposure has been quantified for very few types of events, e.g.,  
35 exposure to tropical cyclones<sup>44, 45</sup> or heat waves<sup>46, 47</sup>, and quantification of future vulnerability is also  
36 rare<sup>4, 48, 49</sup>. Lower mean age, greater wealth, and increased penetration of air conditioning could  
37 ameliorate risk. Recent experience in France<sup>50</sup> and Bangladesh (ref. 43, 9.2.5) provides evidence for the  
38 potential for reductions in vulnerability in both developed and developing countries. In contrast, risks  
39 could increase in the future even if the temperature change remains moderate, since exposure to

1 climate-influenced hazards is increasing significantly in various world regions<sup>48</sup>, particularly in Asia and  
2 Africa due to population-growth, urbanization<sup>51, 52</sup>, and migration.

### 3 **RFC3: Risks associated with the uneven distribution of impacts**

4 This category of risk reflects climate change impacts that disproportionately affect particular groups due  
5 to uneven distribution of physical climate change hazards, exposure or vulnerability. Unevenness can be  
6 with respect to geographic location, income and wealth, gender, age, or other physical and  
7 socioeconomic characteristics.

8 The transition from Undetectable to Moderate risk is located at recent temperatures based primarily on  
9 the detection and attribution with at least medium confidence of negative impacts on wheat yields in  
10 Europe and South Asia (ref. 14, Table 18.9) and evidence of negative agricultural impacts in other  
11 regions as well (ref. 53, Figs. 7.2, 7.7; ref. 54; Supplementary Text 5.1). Some positive impacts on crop  
12 yield have also been detected, for example in Northern Europe and South America (ref. 14, Table 18.9).  
13 AR5 authors took yield impacts as an early warning sign of attributable risk to food security<sup>6</sup>.

14 The transition to High risk occurs between ~1.6 and ~2.6°C based on risks of increased water stress and  
15 reductions in crop production in some regions (Supplementary Text 5.2). Without adaptation, losses in  
16 production of wheat, rice and maize are expected by 2.6 °C of local warming (and therefore typically a  
17 lower level of global warming) although individual locations may benefit<sup>53, 55</sup>. Projections of yield loss are  
18 greatest in low latitudes and tropical regions such as Africa, S. Asia and Central and S. America<sup>53, 55, 56, 57</sup>.  
19 Substantial decreases in water resources are projected for warming of 2.3°C<sup>58, 59</sup>.

20 A transition to Very High risk occurs around 4.6°C based primarily on projected large impacts on crop  
21 yields and water resources in many regions combined with limited scope for agricultural adaptation<sup>53, 55,</sup>  
22 <sup>57, 58, 59</sup>, although other risks contribute (Supplementary Text 5.3). Poorer populations in less developed  
23 countries would be at highest risk of malnutrition, for example in sub-Saharan Africa<sup>60</sup> where food  
24 security is projected to be at risk even under high adaptation levels (ref. 61, 22.5). .

25 A principal limitation to the judgments for this RFC is the sparseness of literature on impacts that can be  
26 linked to levels of GMT in sectors beyond food and water (such as health, energy, civil conflict, urban  
27 areas, and migration<sup>62, 63</sup>) that also have distributional consequences, especially for the poor<sup>64</sup>  
28 (Supplemental Text 5.4). In addition, the food and water literature focuses primarily on biophysical  
29 impacts (such as crop yields or water supply) as opposed to societal impacts (such as food and water  
30 security). The agronomic limits to adaptation considered in the judgment of Very High Risk do not  
31 account for additional means of offsetting yield changes<sup>65</sup> such as changes in cropland and pasture area,  
32 reductions in food waste<sup>66, 67</sup>, and changes in diet<sup>68</sup> or international trade. Biophysical impact studies are  
33 also subject to substantial uncertainties, including the strength of the CO<sub>2</sub> fertilization effect on crop  
34 yields<sup>53</sup>, and the yield effects of extreme events, neither well accounted for (Supplementary Text 5.4).

### 35 **RFC4: Risks associated with global aggregate impacts**

36 This category of risk reflects impacts to socio-ecological systems that can be aggregated globally  
37 according to a single metric such as lives affected, monetary damage, number of species at risk of  
38 extinction, or degradation and loss of a number of ecosystems at a global scale. Ecosystem degradation

1 may be caused by wholesale transformation of biomes, by large scale extirpation of species induced by  
 2 climatic range loss, and by the disruption of ecosystem functioning as interacting species respond  
 3 differently to climate change<sup>29</sup>.

4 AR5 concluded that global aggregate impacts on socio-ecological systems by any of the metrics listed  
 5 above have not yet been detected and attributed to climate change with sufficient confidence to locate  
 6 the transition from Undetectable to Moderate risk at recent temperatures<sup>14</sup> (Supplementary Text 6.1).

7 A Moderate risk level occurs at warming of ~1.6-2.6°C based on projected impacts to biodiversity and  
 8 the global economy (and therefore a transition from Undetectable to Moderate risk between current  
 9 temperatures and ~1.6°C). A global assessment of 16,857 species of all birds, amphibians and corals  
 10 found that with approximately 2°C of warming above preindustrial (A1B, 2050s), 24-50% of birds, 22-  
 11 44% of amphibians and 15-32% of corals were at increased risk of extinction (Supplementary Text 6.2.1)  
 12 due to their vulnerability to climate change<sup>27</sup>. Other studies found increasing extinction risks with  
 13 warming and project range losses exceeding 50%<sup>69</sup> for large fractions of species globally at 2°C warming  
 14 (Supplementary Text 6.2.2). Estimates of global economic damages transition from generally small,  
 15 negative projected impacts around 1°C warming<sup>70</sup> to central estimates of impacts ranging from 0 to 3%  
 16 of global Gross Domestic Product for levels of warming between 1.9 and 3.0°C (Supplementary Text  
 17 6.2.3).

18 The transition to High risk around 3.6°C reflects an increase in the magnitude and likelihood of extensive  
 19 loss of biodiversity (including losses in range, equating to local extirpations) and concomitant loss of  
 20 ecosystem services (Supplementary Text 6.3). There are too few studies of aggregate economic damages  
 21 to provide support for the judgment of risks above 3°C.

22 Limitations of the judgments for this RFC include the limited number of studies that assess global  
 23 aggregate economic impacts that can be associated with specific levels of warming. In addition, global  
 24 estimates of economic damages are incomplete, generally inadequately represent the possibility of  
 25 abrupt and irreversible changes, ignore some impacts that are difficult to monetize, and depend in part  
 26 on value-based judgments that can mask differential impacts through space and time<sup>71</sup> (Supplementary  
 27 Text 6.4). Finally, assessments of impacts on ecosystems insufficiently consider how biotic interactions  
 28 between species may be disrupted by climatic change<sup>72</sup>.

## 29 **RFC 5: Risks associated with large-scale singular events**

30 Large-scale singular events (sometimes called “tipping points”, or critical thresholds) are relatively large,  
 31 abrupt and sometimes irreversible changes in physical, ecological, or social systems in response to  
 32 smooth variations in driving forces (accompanied by natural variability)<sup>73,74</sup>. AR5 focused on two types  
 33 of such events in assessing this risk: disintegration of the Greenland and West Antarctic ice sheets (GIS  
 34 and WAIS, respectively) leading to a large and rapid sea level rise and major regime shifts in ecosystems  
 35 such as degradation of coral reef and Arctic systems. In each case, there is *low confidence* in the precise  
 36 temperature changes at which thresholds might exist for these phenomena (ref. 6, 19.6.3.6; ref. 75,  
 37 12.4.5, 12.5.5; ref. 76, 13.4). For coral reefs, the distinction between the “regime shift” criterion here  
 38 and the systematic degradation indicated under RFC1 resides in the likelihood of abrupt change. While  
 39 the long term outcome for coral reefs under each of the two categories of risk may be similar, RFC5 is

1 concerned with a rapid undermining of system function (where “rapid” and “abrupt” are relative terms;  
2 see discussion below and Supplementary Text 7.2).

3 The transition from Undetectable to Moderate risk between  $\sim 0.6$  and  $\sim 1.6^\circ\text{C}$  warming is based on  
4 potential regime shifts in the Arctic and in coral reef systems. Impacts on the Arctic and on warm water  
5 coral reef systems are already observed (see RFC1), but for RFC5, the detection and attribution criterion  
6 applies to a large and sudden change. There is robust evidence of early warning signals that a  
7 biophysical regime shift already may be underway in Arctic ecosystems, including impacts on human  
8 livelihoods (ref. 14, 18.6.4), and observed increases in mass coral bleaching are considered to be a  
9 strong warning signal for the irreversible loss of an entire biome (ref. 14, 18.6.4).

10 The transition to High risk over the  $\sim 1.6$ - $4.0^\circ\text{C}$  warming range (slightly revised from AR5) is based on ice  
11 sheet responses and the resulting sea level rise. The warming level associated with eventual, near-  
12 complete loss of the Greenland ice sheet is greater than about  $1^\circ\text{C}$  (*low confidence*) but less than about  
13  $4^\circ\text{C}$  (*medium confidence*) (ref. 77, based on ref. 78, 5.8, and ref. 76, 13.4, 13.5). The difference between  
14 the risk range and the ice sheet loss range arises because the risk range implicitly incorporates a  
15 quantification of the implications of the qualitative confidence levels presented by IPCC WGI<sup>77</sup>. Within  
16 this range, a more rapid increase in risk is judged to occur as temperature rises between  $\sim 1.6^\circ\text{C}$  and  
17  $\sim 2.6^\circ\text{C}$ , reflecting additional risk of a very large sea level rise due to ice loss from both ice sheets as  
18 occurred during the Last Interglacial (Supplementary Text 7.1), when GMT was no more than  $2^\circ\text{C}$   
19 warmer than preindustrial levels<sup>79</sup>.

20 Due to the large uncertainty in *timing* of ice sheet loss (which affects the probability of it occurring  
21 sufficiently slowly to allow effective adaptation, e.g. over a millennium, as well as the probability that  
22 action during the next centuries may reduce the warming sufficiently early to limit the melting), RFC5 is  
23 not judged to attain Very High risk in the temperature range below  $\sim 5.6^\circ\text{C}$ , the maximum warming  
24 considered in Figure 1.

25 Improved prognostic modeling of continental ice sheets is a necessity for significantly sharpening this  
26 risk assessment. Post-AR5 literature on such models<sup>80, 81, 82, 83</sup>, observations<sup>84</sup>, and additional lines of  
27 evidence<sup>85</sup> indicate the possibility of large, very fast (decade-to-century scale) responses providing  
28 further support for a tipping point (Supplementary Text 7.2).

29

### 30 **ADDITIONAL METRICS**

31 The RFCs and associated Burning Embers diagram use GMT rise as the proxy indicator for climate-  
32 related hazards. This approach has the benefit of simplifying the communication of risk. However, there  
33 are important climate-related hazards that are inadequately captured by the temperature indicator  
34 alone. We discuss three metrics that were incorporated in complementary ember diagrams in the AR5  
35 Synthesis Report<sup>81</sup> as illustrations of ways in which the analysis of key risks could be extended (Fig. 2)  
36 and informed a recent UNFCCC policy dialog on long-term targets<sup>1</sup>.

#### 37 **Rate of climate change**



1 For many socio-ecological systems the *rate* of climate change determines the success or failure to adapt.  
2 Theory as well as paleo-ecological and paleo-climatic data indicate that adaptation of organisms to  
3 climate change through geographic movement has limits (Supplementary Text 8.1). The 'rate of climate  
4 change' ember assigns risk levels as a function of the rate of climate change during the 21<sup>st</sup> century,  
5 translated into a velocity at which climate zones move across the landscape. A range of species  
6 movement rates was estimated for a number of groups of species by authors of ref. 86, using data from  
7 fossil records, dispersal studies, and models of species movement (see listing of the primary sources in  
8 the caption of fig 4.5 in ref. 86).

9 The relationship between 'climate velocity' (the rate of movement of climate zones) and the rate of  
10 GMT change depends on topography (Supplementary Text 8.2). Thus, at a given rate of GMT change,  
11 risks to species vary depending on location. In addition, there are geographical barriers to species-range  
12 shifts, such as coasts, mountaintops, or habitat fragmentation breaking connections to cooler areas<sup>87</sup>.  
13 Rate of change considerations supplement amount of change rather than replacing it; for instance there  
14 are situations (such as mountaintops) where potentially fast-moving species have nowhere to go.

15 Authors of the IPCC Synthesis Report compared the estimated rates of species movement with  
16 estimates of the climate velocity during past<sup>88, 89, 90</sup> and projected future<sup>91, 92, 93</sup> climate change. Since  
17 trees and herbaceous plants form the productive basis of most terrestrial ecosystems, and flat  
18 landscapes occupy a large part of the land surface, moderate risk was assigned to commence when the  
19 climate velocity exceeded the lower end of the range of observed movement rates (trees in flat  
20 landscapes) and end at the median movement rate for rodents and primates. The risk was assessed as  
21 High beginning where the movement rate exceeded the upper end of the range for trees and ending at  
22 the upper limit for herbs and rodents, beyond the upper limit for primates, and at the median for  
23 freshwater molluscs. Substantial biotic community and ecosystem disruption over large areas could be  
24 anticipated in this range. Very High risks were assigned when the median movement rate was exceeded  
25 in all assessed groups (which included carnivores and split-hoofed animals in addition to the groups  
26 described above). The impact on species assemblages and thus ecosystem function would, with high  
27 likelihood, be large, persistent and difficult to adapt to for this rate of climate change.

## 28 **Anthropogenic CO<sub>2</sub> causing ocean acidification**

29 This ember diagram depicts the increasing risk for the well-being and survival of marine organisms due  
30 to accumulating CO<sub>2</sub> in seawater causing ocean acidification (OA). Since pre-industrial times,  
31 atmospheric CO<sub>2</sub> levels have risen from 280 to presently about 400 ppm, paralleled by a drop in ocean  
32 pH of approximately 0.1 units<sup>94</sup>. Anthropogenic OA occurs on a background of natural temporal and  
33 spatial variability of pH, CO<sub>2</sub>, and aragonite and calcite saturation levels, for example in upwelling areas,  
34 where oxygen-deficient and CO<sub>2</sub>-enriched deep water is brought to the surface.

35 Risks of harmful ecosystem effects of OA are considered Moderate around CO<sub>2</sub> levels of 380 ppm. This  
36 judgment is based on observed declines in calcification of foraminifera and pteropods attributed to  
37 anthropogenic OA<sup>95</sup>. In addition, negative impacts on pteropods and oyster cultures along the west  
38 coast of North America have been attributed to upwelling of acidified water shifted closer to shore  
39 combined with anthropogenic acidification<sup>96</sup>.

1 Under OA only, warming excluded, the transition to High risk occurs at a CO<sub>2</sub> level of about 500 ppm,  
2 beyond which studies reflect onset of significantly negative effects and High risk in 20 to 50 % of extant  
3 calcifying taxa (corals, echinoderms, molluscs). The negative effects comprise declines in physiological  
4 performance, indicated by changes in characteristics such as standard metabolic rate, aerobic scope,  
5 growth, morphology, calcification, acid-base regulation, immune response, fertilization, sperm motility,  
6 developmental time, changes in gene expression patterns, behavioral changes and abundance<sup>81, 95, 97</sup>.  
7 Risks are judged to be Very High with limited capability to adapt beyond about 700 ppm, based on a  
8 rising percentage of the calcifying taxa being negatively affected. For the calcifying invertebrate taxa,  
9 these conclusions are confirmed by observations at natural analogues (volcanic CO<sub>2</sub> seeps, upwelling  
10 systems) and by the similarity of sensitivity distributions among taxa during paleo-periods<sup>97</sup>.

11 Current knowledge indicates that the combined pressures of ocean warming extremes and acidification  
12 lead to a shift in sensitivity thresholds to lower CO<sub>2</sub> concentrations, as seen in corals and crustaceans<sup>95</sup>.  
13 For corals this comes with the risk that OA will increasingly contribute to the reduction in areal extent of  
14 coral ecosystems, already underway as a result of interacting stressors (extreme events, increased  
15 predation, bleaching<sup>98</sup>). Knowledge on the long-term persistence of acidification impacts presently relies  
16 on findings in the paleo-records. Therefore, evidence that changes in extant ecosystems will persist is  
17 limited, especially for fishes. Additionally, knowledge is scarce on compensatory mechanisms and their  
18 capacity and associated limits to long-term evolutionary adaptation under ocean warming and  
19 acidification.

## 20 **Sea-level rise**

21 While sea level change is driven by temperature change, the relationship is uncertain and involves  
22 delays, so that coastal risks are not directly and linearly related to temperature. Accounting for  
23 variability in sea level is also important, because a change in average sea level can disproportionately  
24 increase the likelihood of water levels that exceed the coping capacity of socio-ecological systems.

25 For this ember, the detection and attribution of impacts on society or ecosystems was not used for  
26 judging risk levels due to the difficulty of attributing such impacts. Impact attribution is difficult because  
27 observed increases in impacts are overwhelmingly due to population and socio-economic changes (ref.  
28 33, 5.4.4) or non-climatic, anthropogenic stress (ref. 33, 5.2), and also influenced by historical  
29 investments in coastal protection for which data are lacking. Therefore attribution of sea level rise itself  
30 was used.

31 The transition to Moderate risk starts before the recent period, given that global sea level rise over the  
32 past several decades is attributable to climate change (ref. 17, 10.4.3) and increases the risk of coastal  
33 flooding, soil salinization, and saltwater intrusion. The risk is estimated to reach the Moderate level at  
34 about 10 cm above the 1986-2005 level, which authors of the Synthesis Report estimated to be the level  
35 at which increased flood risks become significant and require changes in coastal management.

36 At this level, the transition to High risks starts and risks are expected to become High at around 100 cm  
37 above the same reference level. High risk is defined for this RFC as the risk of losses that, in the absence  
38 of adaptation, would reach levels that are at least an order of magnitude higher than today, and cause  
39 coastal ecosystem losses that are visible and widespread. High risks may occur before the 100 cm level is

1 reached, since some evidence suggests the risk would increase rapidly even before this value  
 2 (Supplementary Text 8.3). For example, for sea level rise of 40-130 cm, 1.3 to 2.9% of the world  
 3 population could be flooded every year<sup>99</sup>.

4 The transition to Very High risk is expected over the range of 100 – 200 cm above the 1986-2005 level.  
 5 This transition starts where adaptation limits for ecosystems and human systems are reached in many  
 6 places. Limited evidence suggests that only a small number of adaptation options are available for  
 7 specific coastal areas if sea level exceeds 100 cm at the end of the century (ref. 33, 5.5.6). There are also  
 8 biophysical limits to the adaptation of ecosystems and natural areas, which vary greatly depending on  
 9 the rate of change, location and other stressors (ref. 33, 5.2).

10

## 11 **TEXT BOX 2: RFCS AND THE VULNERABILITY OF SOCIO-ECOLOGICAL SYSTEMS**

12 The Burning Embers diagram does not explicitly account for differences in the exposure and  
 13 vulnerability of socio-ecological systems over time, including those changes arising from adaptation. In  
 14 AR5, judgments about risks reflected in the Burning Embers diagram were based on the varied  
 15 assumptions in the underlying literature about future societal conditions that would affect vulnerability  
 16 and exposure, including income and poverty, technology, demography, institutions, and other factors.  
 17 These assumptions range from complete disregard of future societal conditions, to central or middle-of-  
 18 the-road expectations, to differing societal futures across studies which were then aggregated by IPCC  
 19 authors. Only autonomous adaptation (i.e., adaptation that does not require coordinated planning) is  
 20 reflected in the impacts used to make risk judgments.

21 At the same time, a growing number of examples in the impact literature demonstrate the dependence  
 22 of impacts on societal conditions, especially the differential vulnerability of people and ecosystems  
 23 exposed<sup>100</sup> (Supplementary Text 9). AR5 concluded with *high confidence* that risks vary substantially  
 24 across plausible alternative development pathways, and that both climate change and societal  
 25 development are important to understanding possible future risks (ref. 6, 19.6.2.2). AR5 also introduced  
 26 an alternative version of a burning ember with an additional axis for exposure and vulnerability.  
 27 However, the figure was conceptual, illustrating how risks for a particular RFC might vary by societal  
 28 conditions as well as by the level of climate change.

29 Here we illustrate how a vulnerability-dependent version of a burning ember diagram could be  
 30 developed, drawing on impact studies<sup>101, 102, 103</sup> that project the number of people at risk of hunger  
 31 under alternative assumptions about future vulnerability and climate change. There is substantial  
 32 uncertainty about estimates of hunger risk for any given societal and climate future, due to, for  
 33 example, uncertainties in crop modeling, the effects of CO<sub>2</sub> fertilization, economic models of food  
 34 consumption, and factors affecting access to food. This uncertainty precludes judgments about the  
 35 *absolute* level of risk for any given climate and vulnerability outcome, and therefore the production of a  
 36 burning-ember style diagram. However, judgments about changes in risk if climate or vulnerability varies  
 37 from a given outcome in the future are possible. Fig. 3a shows changes in the number of people at risk  
 38 of hunger due to climate change relative to the number at risk for a particular set of conditions used as a  
 39 benchmark (medium vulnerability, and about 2.6 °C of GMT). The general pattern confirms that lower

1 vulnerability development pathways minimize risk, while increases in level of future warming result in a  
2 larger risk. Exceptions occur in studies in which relatively large CO<sub>2</sub> concentration increases improve  
3 access to food in scenarios with a further warming of 3°C and high vulnerability (figure 3). As additional  
4 literature accumulates, it may be possible to use the type of approach illustrated here to produce a  
5 fuller assessment of vulnerability-dependent RFCs, including explicit treatment of adaptation  
6 (Supplementary Text 10.1).

7 A complementary view of future risks is provided by Figure 3b, which shows that the total number of  
8 people at risk of hunger is much more sensitive to the development pathway than to the level of climate  
9 change.

10

## 11 **DISCUSSION AND FUTURE DIRECTIONS**

12 The RFCs were designed to categorize and depict increasing risks from warming of the climate system  
13 and thereby inform (but not determine) judgments about danger from climate change. Within the  
14 current limits of the framework, the RFC assessment provides a number of insights relevant to Article 2.  
15 First, continued high emissions would lead to High or Very High risk of severe, widespread, and in some  
16 cases irreversible impacts globally within this century. Risks to unique and threatened systems, among  
17 the most sensitive natural and human systems, increase most quickly with additional warming. Risks  
18 associated with global aggregate impacts increase most slowly.

19 In addition, the RFCs can communicate the specific nature of current and future risks. For RFCs 1-3, risks  
20 from anthropogenic climate change are currently Moderate, based primarily on detection and  
21 attribution of associated impacts on Arctic ecosystems and coral reefs (RFC1); extreme heat and  
22 precipitation events and their impacts on human health and coral reefs (RFC2); and impacts on crop  
23 production in some regions (RFC3). In terms of future risk, at 2°C above preindustrial, High risks are  
24 based on increasing risks to Arctic systems and coral reefs, as well as increasing species extinction risks  
25 (RFC 1), and projected increasing magnitude and likelihood of extreme weather events (RFC 2).  
26 Moderate-to-High risks are based on projections of increasing risks to crop production and water  
27 resources (RFC 3), and to the risks associated with ice sheet disintegration and very large sea level rise  
28 (RFC5). Limiting warming to 1.5°C would reduce the risks for RFCs 1 and 2 from High to the Moderate-to-  
29 High transition.

30 At 3°C above pre-industrial, risks are at least High, or nearly so, for all RFCs. In addition to the basis for  
31 High risk judgments that apply to 2°C, additional factors include a higher risk of species extinction (RFCs  
32 1 and 4), limited ability to adapt to impacts on coral reefs and Arctic systems (leading to Very High risk  
33 for RFC 1), and the higher risk of very large sea level rise associated with eventual ice sheet loss (RFC 5).

34 Judgments, choices, and decisions informed by the RFCs should take into account key challenges faced  
35 by this framework. First, the assessment of risk levels across the RFCs has been based primarily on  
36 impacts to physical and ecological systems, given a literature on consequences for society that is either  
37 thin or difficult to relate to specific levels of climate change and future societal conditions. Extensions to  
38 the framework to explicitly account for the vulnerability of socio-ecological systems (Fig. 3) offer a  
39 means of incorporating new knowledge of this type.

1 Second, aggregating risks across affected sectors and systems necessarily suppresses the detail and  
2 variation of associated risks. Communicating the specific key risks informing the RFC assessment is  
3 important to prevent misinterpretation of risk judgments and to better inform discussion of response  
4 options. It may also be possible to extend the burning embers diagram to better represent individual  
5 risks (Supplementary Fig. 2).

6 Third, the perceived seriousness of risks will vary by stakeholder and among authors carrying out the  
7 assessment. Some may value the existence of species and ecosystems – beyond their role in providing  
8 ecosystem services – more highly than others, and therefore perceive particular RFCs (such as RFC1) as  
9 more important. Others may prioritize aggregate damages or may consider equity and distributional  
10 impacts as paramount.

11 Finally, additional dimensions of climate change beyond GMT, such as the rate of climate change, ocean  
12 acidification, and sea-level rise, can be important metrics of hazard, sometimes more directly linked to  
13 impacts than global mean temperature.

14 These caveats, and the review of the RFCs as a whole, suggest a number of research needs. More  
15 systematic evaluation of key risks and impacts at varying levels of climate change is needed to inform a  
16 more complete, specific and quantitative understanding of the differential impacts across possible  
17 climate futures for a larger number of key risks. A deeper literature of this kind would avoid imbalances  
18 in the role of specific risks, such as the large role of agricultural risks in RFC3 and the role of risks to coral  
19 reefs across several RFCs. It would also improve understanding of the uncertainty in the level of GMT  
20 associated with risk transitions. There is an equally strong need for research on socioeconomic  
21 dimensions of risks, to improve on the current common use of physical climate system outcomes as a  
22 proxy for societal impacts. In particular, more work is needed on how alternative societal development  
23 pathways, implying different levels of vulnerability to climate change and possibilities for adaptation,  
24 affect the risks of any given level of warming.

25 Beyond improving the research base, modifications to or extensions of the RFC framework itself may be  
26 called for, especially as new evidence accumulates, while also recognizing the value of simplicity in  
27 communicating risk. Efforts should be continued to make the methods for producing the RFCs and the  
28 associated burning embers diagram more systematic, transparent and comparable across generations.  
29 Improvements in these aspects of the RFCs will also make them more effective tools for informing  
30 decisions related to avoiding dangerous climate change.

31

## 32 **Corresponding Author**

33 Correspondence to: boneill@ucar.edu.

## 34 **Acknowledgements**

35 K.T. gratefully acknowledges research support of the “Global Environmental Research Fund” (S-10-1)  
36 provided by the Ministry of the Environment, Japan.

1 **Author Contributions**

2 B.C.O. and M.O. led the design of the study. B.C.O. led, and M.O. contributed to, the coordination of the  
3 paper. M.O., R.W., S.H., R.E.K., B.C.O., H.O.P., and B.S. led the drafting of subsections of the paper.

4 B.C.O., P.M., R.L., K.J.M., M.M., and K.T. led the development of figures. All authors contributed to  
5 writing and/or editing the paper.

6

1 **TABLES**

2 **Table 1:** Eight overarching key risks representative of the range of key risks identified by WG II authors  
 3 as of highest concern to their chapters (ref. 6, 19.6.2.1, based on Table 19-4). These risks inform  
 4 judgments regarding the indicated RFCs.

Overarching Key Risk	Description	Reason for Concern				
		1	2	3	4	5
i	Risk of death, injury, ill-health, or disrupted livelihoods in low-lying coastal zones and small island developing states and other small islands, due to storm surges, coastal flooding, and sea level rise.	✓	✓	✓	✓	✓
ii	Risk of severe ill-health and disrupted livelihoods for large urban populations due to inland flooding in some regions.		✓	✓		
iii	Systemic risks due to extreme weather events leading to breakdown of infrastructure networks and critical services such as electricity, water supply, and health and emergency services.		✓	✓	✓	
iv	Risk of mortality and morbidity during periods of extreme heat, particularly for vulnerable urban populations and those working outdoors in urban or rural areas.		✓	✓		
v	Risk of food insecurity and the breakdown of food systems linked to warming, drought, flooding, and precipitation variability and extremes, particularly for poorer populations in urban and rural settings.		✓	✓	✓	
vi	Risk of loss of rural livelihoods and income due to insufficient access to drinking and irrigation water and reduced agricultural productivity, particularly for farmers and pastoralists with minimal capital in semi-arid regions.		✓	✓		
vii	Risk of loss of marine and coastal ecosystems, biodiversity, and the ecosystem goods, functions, and services they provide for coastal livelihoods, especially for fishing communities in the tropics and the Arctic.	✓	✓		✓	
viii	Risk of loss of terrestrial and inland water ecosystems, biodiversity, and the ecosystem goods, functions, and services they provide for livelihoods.	✓		✓	✓	

5

1 **FIGURES**


2

3 **Figure 1: The enhanced burning embers diagram, providing a global perspective on climate-related**  
4 **risks.** Levels of risk associated with 5 different reasons for concern are illustrated for increasing global  
5 mean temperature and are the same as those presented in the IPCC Working Group II report. Icons  
6 indicate selected risks that played an important role in locating transitions between levels of risks.  
7 Colored dots indicate overarching key risk categories that were considered in the assessment for each  
8 RFC (see Table 1). Confidence in the judgments of risk transitions is indicated as provided in ref. 104. For  
9 example, RFC1 is underpinned by overarching key risks i, vii, and viii from Table 1; there is medium  
10 confidence in the location of the transition from Undetectable to Moderate risk, which is informed by  
11 impacts to coral reef, Arctic and mountain systems; and there is high confidence in the location of the  
12 transition from High to Very High risk, which is informed by impacts to coral reef and Arctic systems as  
13 well as to species associated with unique and threatened systems.

14



1



2

**Selected Key Risks**

- Biodiversity
- Mountain systems
- Agriculture
- Water stress
- Coral reefs
- Heat waves
- Economic damages
- Greenland ice sheet
- Arctic systems
- Extreme precip.
- Human health
- Antarctic ice sheet

3


4

5

6


1 **Figure 2: Additional burning embers diagrams from the AR5 Synthesis Report<sup>81</sup>.** These figures use (a)  
 2 rate of climate change, (b) atmospheric CO<sub>2</sub> and associated ocean acidification as well as (c) sea level  
 3 rise as the metric of climate-related hazard, rather than global mean temperature (for further  
 4 explanations see text).

**(a) Risk for terrestrial and freshwater species impacted by the rate of warming**





5

**(b) Risk for marine species impacted by ocean acidification only, or additionally by warming extremes**



6

(c) Risk for coastal human and natural systems impacted by sea level rise




1

2

1 **Figure 3, panel a: Illustrative version of a vulnerability-dependent burning embers diagram.** The figure  
2 uses results for one type of climate change impact (additional population at risk of hunger due to  
3 climate change) based on three studies<sup>101, 102, 103</sup>. The x-axis categorizes scenarios of societal  
4 development by trends in exposure and vulnerability based on ref. 34. Each colored circle indicates the  
5 difference between the number of people at risk of hunger due to climate change according to one  
6 scenario and the number at risk as calculated under benchmark outcomes. Benchmark conditions are  
7 defined as those associated with a medium vulnerability scenario with about 2.6 °C warming relative to  
8 preindustrial (Box 2, and Supplementary Text 10 for further description). Results for this benchmark  
9 outcome are plotted as zero (a white square) in the figure. Green circles indicate lower risk than this  
10 benchmark outcome (values <0), and generally occur for lower levels of climate change and/or lower  
11 levels of societal vulnerability. Yellow, red and purple circles indicate greater risk (values >0), and  
12 generally occur for more climate change and/or higher societal vulnerability. The figure incorporates 40  
13 scenarios with a range of economic, crop and climate models and assumptions about CO<sub>2</sub> fertilization  
14 and adaptation (Supplementary Table 1). The medium vulnerability, 2.6° C (benchmark) outcomes span  
15 a range of -12 to 151 million additional people at risk of hunger, illustrating the relatively large  
16 uncertainty in estimates of this risk. The Exposure and Vulnerability (E&V) axis indicates relative trends  
17 over time rather than absolute levels, with current conditions defined as “Medium” E&V. For example, a  
18 future development path in which E&V remains “Medium” is assumed to change over time (and  
19 therefore also with changes in GMT along the y-axis) at a moderate rate, driven by trends in  
20 socioeconomic conditions that are in the middle of the range of future scenarios. “Low” and “High” E&V  
21 indicate futures that are substantially more optimistic or pessimistic, respectively, regarding trends in  
22 exposure and vulnerability.

1 a



2



1 **Supplementary Information**

2 **IPCC Reasons for Concern regarding climate change risks**

3 Brian C. O'Neill, Michael Oppenheimer, Rachel Warren, Stephane Hallegatte, Robert E. Kopp, Hans O.  
4 Pörtner, Robert Scholes, Joern Birkmann, Wendy Foden, Rachel Licker, Katharine J. Mach, Phillippe  
5 Marbaix, Michael Mastrandrea, Jeff Price, Kiyoshi Takahashi, Jean-Pascal van Ypersele and Gary Yohe

6 Corresponding author: [boneill@ucar.edu](mailto:boneill@ucar.edu)

7  
8 This document contains Supplementary Discussion for the following topics appearing in the main paper:

- 9 **1. Process for making judgments about RFC risk levels**  
10 **2. Units of global mean temperature change (GMT)**  
11 **3. RFC1: Unique and Threatened Systems**  
12 **4. RFC2: Extreme events**  
13 **5. RFC3: Distributional Impacts**  
14 **6. RFC4: Global Aggregate Impacts**  
15 **7. RFC 5: Large-scale singular events**  
16 **8. Additional Metrics: Rate of climate change**  
17 **9. RFCs and the vulnerability of socio-ecological systems**  
18 **10. Figure captions**

19 It also contains the following supplemental tables and figures:

20 **Supplementary Table 1: Scenario results informing Figure 3.**

21 **Supplementary Figure 1: Process for assessing risks associated with Reasons for Concern in the IPCC**  
22 **Fifth Assessment Report.**

23 **Supplementary Figure 2: Decomposition of burning ember for RFC4 into two primary key risks.**

24 Note that as in the main text, all undesigned temperatures are with respect to preindustrial.

25

1 **1. Process for making judgments about RFC risk levels**

2 The assessment based on the criteria for key risks listed in the main text proceeded in two streams,  
 3 which eventually merged. In the first, AR5, Working Group II, Chapter 19 authors defined varying  
 4 discrete levels of risk in terms of the key risk criteria, and made a preliminary assessment of risks as a  
 5 function of GMT for each RFC based on relevant impact studies. At the same time, regional and sectoral  
 6 chapters of the assessment used the criteria in combination with their own expert judgment to identify  
 7 key risks in their domains (ref. 6, Table KR-1, Table 19.4). For example, risk of loss of rural livelihoods is a  
 8 broader concern than loss of urban livelihoods, based on the observed large and pervasive rural-to-  
 9 urban migration and the expected intensification of climate-related factors underlying this migration.  
 10 Similarly, the risk from water stress is greater for rural than urban areas due to the limited potential to  
 11 adapt for the former areas. The risk from heat-related morbidity and mortality is widespread and  
 12 imminent, and increases in extreme heat are virtually certain by 2100. In contrast, the relationship of  
 13 climate changes to incidence of vector-borne diseases is less direct and more difficult to quantify.

14 These regional and sectoral key risks were synthesized into 8 overarching risk categories, each of which  
 15 was then associated with one or more of the RFCs and integrated with the preliminary assessment of  
 16 risk levels for each RFC (Table 1, main text; ref. 6, 19.6.2.1). For example, the fourth key risk,

17 **“risk of mortality and morbidity during periods of extreme heat, particularly for vulnerable**  
 18 **urban populations and those working outdoors in urban or rural areas”**,

19 is associated with RFC2 and RFC3 because it is directly connected with outcomes of extreme events and  
 20 is unevenly distributed given geographic differences in hazards and variations in the vulnerability and  
 21 exposure of populations.

22 Levels of risk were associated with GMT for each RFC by Chapter 19 authors based on an aggregation of  
 23 the relevant key risks and on individual key risks for which the underlying scientific literature allowed  
 24 clear association with particular levels of climate change. Although this assessment expressed additional  
 25 risk due to climate change as a function of GMT, it required direct or indirect consideration of exposure  
 26 and vulnerability since these dimensions of risk are part of the key risk criteria.

27 Due to the limits of current information, the assignment of risk levels for each RFC comes down to an  
 28 assessment of risk of only some of the impacts pertinent to each RFC. Ideally, the approach described  
 29 here would provide a systemization that may allow consistency and transparency, at least in the basis of  
 30 judgment used by the same or different groups of experts in future assessments.

31

32 **2. Units of global mean temperature change (GMT)**

33 In AR5<sup>6</sup>, risk transitions were located using GMT relative to recent temperatures (defined as the 1986-  
 34 2005 average), and the approximate nature of these transitions was signaled in part by using round  
 35 numbers (1 C, 2 C, etc.). To convert to warming relative to pre-industrial (defined as the 1850-1900  
 36 average), we subtracted 0.6 C from these values, following ref. 105 which evaluates it as 0.61±0.06 °C  
 37 cooler than the recent period. This can lead to the appearance of overly precise locations. We indicate



1 the approximate nature of these transitions either by explicitly stating that locations are “around” or  
2 “approximately” a certain value, or using the “~” symbol (e.g., ~1.6 C).

3

### 4 **3. RFC1: Unique and Threatened Systems**

#### 5 **3.1. Transition from Undetected to Moderate risk at below recent temperatures.**

6 Climate change impacts have also been detected and attributed in numerous ecosystems around the  
7 world, including those usually considered unique and threatened (see RFC4). Regarding mountain  
8 systems, a range of climate change-induced ecosystem impacts such as increased mountain forest  
9 mortality and wildfire, and tree line shifts have been observed in Europe, Asia, Africa and elsewhere<sup>89,</sup>  
10 <sup>106, 107, 108, 109</sup>. Socioecological systems dependent on glacier melt are unique, and there is high confidence  
11 that glaciers have continued to shrink almost worldwide<sup>77</sup>.

#### 12 **3.2. Transition to High risk at 1.1 – 1.6 °C above preindustrial.**

13 As indicated in the main text, this judgment was supported by projected impacts on Arctic and coral reef  
14 systems. The mean projected loss of Arctic sea ice at around 1.6 °C of warming above preindustrial is  
15 more than 2 million km<sup>2</sup> (30%) relative to recent extent (ref. 75, Figure 12.28b and Table 12.2, based on  
16 RCP2.6 for the period 2046-2065 when global temperatures have reached 1.6°C ± 0. 3°C). Reductions in  
17 sea ice affect the Inuit culture, which subsists on sea ice-dependent ecosystems, as well as species  
18 dependent on sea ice, such as polar bears<sup>110</sup>, which have recently been listed as Vulnerable on the IUCN  
19 Red List due to climate change driven habitat loss<sup>111</sup>. Recent work shows polar bears have exhibited little  
20 adaptive capacity to the loss of sea ice (e.g., they are generally not changing their behavior by feeding in  
21 terrestrial areas<sup>28</sup>).

22 Most warm-water coral reefs are projected to be in rapid and terminal decline from warming and  
23 acidification at atmospheric CO<sub>2</sub> levels of 450 ppm<sup>112</sup> and a corresponding global temperature rise of  
24 around 1.6°C (as in RCP2.6 in the 2050s). Similarly, if there is no change in corals’ thermal tolerance, a  
25 median of 95% of coral reefs is projected to be subject to long-term degradation for a global mean  
26 temperature rise of 1.5-2°C above pre-industrial, and protection of >10% of the world’s reefs is  
27 considered to require constraining warming to 1.5°C above pre-industrial<sup>113</sup>. This assessment is  
28 conservative as it does not consider the combined impacts of warming and acidification.

#### 29 **3.3. Transition to Very High risk at 2.6 °C above preindustrial.**

##### 30 *3.3.1. Impacts on unique and threatened species and ecosystems around or above 2° C warming*

31 Ref. 29 found that 15-37% of the thousands of species studied, which included a large proportion of  
32 species found in unique and threatened systems, would be at an increasing risk of extinction for a global  
33 temperature rise of 2.1-2.3°C above pre-industrial. There are a number of important examples of risks to  
34 specific unique and threatened species and ecosystems. A large fraction (48-57%) of the plants of the  
35 Cerrado are projected to be at increased risk of extinction<sup>114</sup>. In South Africa, 51-65% of the area of the  
36 Fynbos and 80% of the area of the unique Succulent Karoo floral kingdom<sup>115</sup>, both of which are unique  
37 and threatened biodiversity hotspots and of global significance, are projected to be lost<sup>15</sup>. A 24% loss of  
38 freshwater fish habitat in N America is also projected for this level of temperature rise<sup>116</sup>. In the

1 Queensland World Heritage rainforest, another unique ecosystem, which comprises the oldest  
 2 rainforest in the world, a local warming of 1°C above recent (equivalent to a global warming of  
 3 approximately 2°C above pre-industrial; ref. 75, Table 12.2; ref. 117, Figure A1.69) is projected to cause  
 4 a loss of approximately 40% of the rainforest habitat of endemic rainforest vertebrates, with the  
 5 regionally iconic Golden Bowerbird losing 40-60% of its range, one other species losing its entire  
 6 range<sup>106, 118</sup> and 5% of western Australian Banksia plant species projected to become extinct<sup>119</sup>. A meta-  
 7 analysis of projections of increased extinction risk found that the global extinction risk nearly doubled  
 8 from current rates with an increase of 2°C, and then trebled again between 2°C and 4°C<sup>32</sup>. While the  
 9 proportions at risk in ref. 32 for endemics were lower than those originally calculated in ref. 29, the  
 10 differences come from a more restricted definition of increased extinction risk. This most recent meta-  
 11 analysis supported the other findings in that it found that endemic species were more at risk than  
 12 widespread ones, and that species in South America, Australia/New Zealand and Africa were at higher  
 13 risk than those in North America or Europe.

### 14 3.3.2. *Risks to Arctic, coral reef, and mountain systems*

15 For example, a large proportion of CMIP5 climate models project a nearly ice-free Arctic Ocean in the  
 16 summer between 1.6°C and around 2.6°C of warming (ref. 75, Fig. 12-30), while a much smaller  
 17 proportion project that this occurs between 2.6°C and 3.6°C. Regarding coral reefs, ref. 113 find that  
 18 even under optimistic assumptions about ability to adapt to warming, only 6% of reefs would avoid long-  
 19 term degradation with 2.0°C warming above pre-industrial. Loss or degradation of corals would  
 20 endanger the livelihoods of resource dependent human communities and cause substantial economic  
 21 damage in East Africa, Asia, Australia, and parts of N. and S. America. For example, revenue loss of  
 22 between US\$95-140 million annually is expected in the Caribbean basin in the next decade, while in Fiji,  
 23 impacts of between US\$0.1 million to 2 million in subsistence fisheries, and US\$0.05-0.8 million in  
 24 commercial coastal fisheries by 2050<sup>120</sup> are estimated. Globally, 20-25% of the fish caught by developing  
 25 nations come from coral reefs<sup>121</sup> and 6 million people fish in coral reef systems<sup>122</sup>. Glacier meltwater  
 26 supports a number of unique mountain ecosystems and social systems, in particular in Central Asia and  
 27 S America. As glaciers melt, initially streamflow may increase as a result of accelerated melting, after  
 28 which flow declines again, as seen in the Cordillera Blanca of Peru<sup>123</sup>, while at the same time the melting  
 29 creates an associated risk of potentially damaging Glacial Lake Outburst Floods as meltwater collects  
 30 behind barriers and suddenly breaches them. In central Asia there is potential for loss of large (but  
 31 uncertain) fractions of central Asian glacier cover by 2100 if temperatures rise by approximately 2-4°C  
 32 above pre-industrial levels in the second half of the 21<sup>st</sup> century, potentially creating water security  
 33 issues in downstream populations dependent on glacier melt<sup>124, 125, 126, 127</sup>. In S. America, glacier melt-  
 34 dependent areas are at risk of water resource stress under 2°C warming above pre-industrial levels in  
 35 the second half of the 21<sup>st</sup> century (and even higher risks under 4°C warming), with several studies  
 36 projecting river flow declines of between 20 and 40% by the end of the century<sup>123</sup>.

### 37 3.3.3. *Limitations to risk judgments*

38 Studies often do not consider the benefits of adaptation designed to reduce species and ecosystem  
 39 impacts, such as interventions to increase ecosystem resilience or design of protected areas to facilitate  
 40 movement in response to warming. In addition, studies frequently do not consider the potential

1 negative effects on species and ecosystems of adaptation to other impacts. For example if an area  
 2 warms and dries, biodiversity and agriculture will both tend to move into the same cooler, wetter areas,  
 3 and the ability of natural systems to track climate change will be impeded by human efforts to adapt.

4 **4. RFC2: Risks from extreme events**

5 **4.1. Transition from Undetected to Moderate risk at recent temperatures.**

6 AR5 WGI found it *very likely* that human influence has contributed to changes in the global scale  
 7 frequency and intensity of daily temperature extremes (ref. 17, 10.6.1.1) and also increased the  
 8 probability of occurrence of heat waves in some locations (ref. 17, Section 10.6.2). AR5 assessed  
 9 *medium confidence* in attribution of intensification of heavy precipitation over Northern Hemisphere  
 10 land areas with sufficient data (ref. 17, Section 10.6.1.2). In developing countries where urban  
 11 infrastructure is expanding and population is growing, the intensification of the urban heat island effect  
 12 is increasing the risk of heat stress<sup>43</sup>. In other areas, such as Germany and Japan, the susceptibility of  
 13 people exposed to heat stress is generally increasing due to the aging population<sup>43</sup>. Vulnerability,  
 14 exposure, and resulting mortality from extreme heat is currently widespread<sup>128</sup>, as is vulnerability and  
 15 exposure of property and people to heavy precipitation. Accordingly, such risks were judged to be  
 16 sufficiently high currently to provide additional support for the assessment of a transition to Moderate  
 17 risk.

18 **4.2. Transition to High risk at 1.6 °C above preindustrial**

19 On average, the frequency and intensity of heavy precipitation events over land will *likely* increase over  
 20 much of the world, particularly the Northern Hemisphere and East Africa with exceptions for Central  
 21 America, the Mediterranean Basin, Australia and South Africa<sup>129</sup> (ref. 42, Section 11.3.2.5.2; ref. 77,  
 22 Table SPM.1).

23

24 **5. RFC3: Distributional Impacts**

25 **5.1. Transition from Undetected to Moderate risk at recent temperatures**

26 Climate change has also been shown to have played a major role in the decline of fruit-bearing trees in  
 27 the Sahel<sup>14, 130</sup> and a minor role in the spread of bluetongue virus in European sheep<sup>131</sup>.

28 **5.2. Transition to High risk between 1.6 and 2.6°C above preindustrial**

29 Many studies project significant reductions in yields of wheat, maize, millet and sorghum in Africa and S.  
 30 Asia by the 2050s<sup>55, 57</sup>, corresponding to global temperature of approximately 1.5-2C above pre-  
 31 industrial levels (based on temperature response to SRES scenarios<sup>132</sup>). Many studies project significant  
 32 negative impacts of climate change on crops and livestock in Central and S America<sup>123</sup>. For example with  
 33 around 2 °C warming above pre-industrial (SRES A2 scenario, 2050s), ref. 133 projects that 80% of all  
 34 crops grown in Colombia will be negatively affected in 60% of the current cultivated area. Ref. 134  
 35 projects reductions in yields of crops important to large food-insecure populations in Africa (e.g., maize  
 36 yield change of  $-21.4 \pm 8.6\%$ ) and S Asia (e.g., rice yield change of  $-4 \pm 2\%$ ) by the 2030s with  
 37 approximately 1.6°C warming relative to preindustrial (a somewhat lower level of global warming).

1 Regarding water resources, projected decreases for warming of 2.3°C<sup>58, 59</sup> include annual mean  
 2 discharge losses of between 10-30% in European Mediterranean areas, parts of the southern USA, and  
 3 central S America, and losses of 30-50% in African Mediterranean areas, parts of Western Australia and  
 4 Southern Africa. “Severe” impacts, defined as a reduction in water resources of more than 20% and/or  
 5 more than one standard deviation of the current natural variability, would affect about 8% and 14% of  
 6 the global population at 1.7°C and 2.7°C respectively above pre-industrial levels<sup>59</sup> with areas  
 7 surrounding the Mediterranean (Middle East, N. Africa, S. and E. Europe) most consistently affected.  
 8 Ref. 135 reports that for warming of 2.0°C above pre-industrial around 8% of the population is projected  
 9 to experience new or increased water stress compared to the present day. Ref. 58 find that water  
 10 scarcity (defined as a water crowding index below 1000 m<sup>3</sup>/capita/yr) increases rapidly in N., W., and E.  
 11 Africa, Central Asia, Central America and Mashriq as temperatures rise by about 1°C above pre-  
 12 industrial, and continues to increase as temperatures rise to about 2.0°C above pre-industrial. Use of  
 13 additional groundwater resources to compensate for projected reduced surface water availability may  
 14 be problematic in many areas since it is projected that for around 2.0°C warming, by the end of the  
 15 century 24% of the projected population will incur >10% decrease in groundwater resources<sup>136</sup>. Using  
 16 water storage to adapt may be physically infeasible in some areas, and is likely to be financially  
 17 infeasible in many others: a global study found that it would cost approximately \$12bn/yr to increase  
 18 global water storage by ~35% by 2050 in order to maintain water supplies<sup>137</sup>.

### 19 **5.3. Transition to Very High Risk at 4.6° C above preindustrial**

20 Risks of flooding, water scarcity and human health impacts contribute to this judgment. The proportion  
 21 of the global population exposed to a 20<sup>th</sup> century 100 year fluvial flood is projected to be three times  
 22 higher for warming of around 4°C above pre-industrial levels (RCP8.5, 2100<sup>75</sup>) than for warming of  
 23 approximately 1.6°C above pre-industrial (RCP2.6, 2100<sup>75</sup>)<sup>138</sup>, or 14 times higher than present day  
 24 exposure; whilst a multi-model study projects increases in flood frequency over half the land surface,  
 25 and decreases in one third<sup>139</sup>. Runoff in the Nile and Ganges basins are projected to increase by up to  
 26 150% and 80% respectively under warming of around 4°C above pre-industrial levels. Ref. 59 project  
 27 that the population exposed to a severe reduction in water resources would increase to 15% with 4°C  
 28 warming above pre-industrial. Regionally, water stress is projected to increase further, particularly in  
 29 West Africa, Central America, Brazil, USA, Eastern Europe and East Asia, as global temperatures increase  
 30 from around 2°C above pre-industrial to around 4°C above pre-industrial, associated with projected  
 31 runoff reductions of up to 90% in the Amazon and 75% in the Danube & Mississippi<sup>140</sup>.

32 With 4°C warming important tipping points for health impacts related to heat stress, such as heat stroke  
 33 and fatal occupational heat stroke, may be exceeded in many parts of the world. In S.E. Asia, for  
 34 example, such impacts may occur as regional temperatures increase by 4-7°C<sup>39, 141, 142, 143, 144</sup>, with  
 35 associated decreases in labor productivity both indoors and outdoors<sup>145</sup>.

### 36 **5.4. Limitations to risk judgments**

37 Civil conflict is an example of a type of impact beyond those in the food and water sectors for which  
 38 literature is sparse and difficult to relate to specific levels of GMT. Empirical studies do suggest that  
 39 extremes of temperature and precipitation increase the relative risk of civil conflict<sup>146</sup>; since the absolute

1 risk of civil conflict is highest in poorer countries, and since civil conflict reduces economic growth<sup>147</sup>,  
 2 this effect will most severely impact the poor.

3 Agricultural impacts are sensitive to assumptions regarding CO<sub>2</sub> fertilization. Food prices, which strongly  
 4 influence food security, are very likely to increase by 2050 in response to climate change if CO<sub>2</sub>  
 5 fertilization effects do not occur, but are only about as likely as not to increase if they do (ref. 53, 7.4.4).  
 6 In addition, a concurrent effect of this process is a reduction in the protein and nutrient content of  
 7 wheat, rice, maize, barley, potato, soybean and peas, meaning the contribution that yield makes to food  
 8 security per ton of crop harvested may decrease at higher CO<sub>2</sub> levels<sup>53, 148</sup>. Regarding the role of heat  
 9 extremes in crop yield impacts, warming is projected to triple the exposure of crops to drought<sup>149</sup>, for 4°  
 10 C relative to preindustrial) and extreme heat<sup>150</sup>, for wheat and maize at 3.5° C warming. A recent  
 11 study<sup>151</sup> found that extreme heat at anthesis is projected to double climate change-induced global losses  
 12 of maize yield, and to halve projected increases in wheat yield (due to an assumption of CO<sub>2</sub> fertilisation  
 13 effects in this particular model) for a warming of 4.3°C above pre-industrial levels (RCP8.5, 2080s; ref.  
 14 75, Table 12.2). The extreme 2003 and 2010 summers in Europe and Russia respectively are known to  
 15 have reduced grain yields by 20-30%<sup>152, 153</sup>.

16 As noted in the main text, the limitations to adaptation to agricultural impacts that play a role in the  
 17 judgment of Very High Risk do not consider a number of adaptation options. The adaptation limits refer  
 18 only to incremental agronomic adaptations such as changes in cultivars and planting dates, which are  
 19 projected to fall short of offsetting the negative effects on yield for local increases in temperature  
 20 exceeding 3°C in tropical areas (ref. 53, 7.5.1.1.1).

21

## 22 **6. RFC4: Global Aggregate Impacts**

### 23 **6.1. Transition from Undetected to Moderate risk**

24 While detection and attribution of globally aggregated impacts on socio-ecological systems has not been  
 25 achieved, detection and attribution *has* been accomplished for global biophysical impacts such as  
 26 changes in the cryosphere or global crop yields. It has also been established for widespread species  
 27 impacts such as earlier onset of spring events (flowering, breeding, etc.) and poleward and upward  
 28 movement of the geographic ranges of many species<sup>86, 154</sup>; changes in phenology in plants, birds,  
 29 amphibians, mammals, and freshwater plankton<sup>14, 155, 156, 157, 158, 159</sup>; changes in distribution<sup>14, 89, 158, 160, 161,</sup>  
 30 <sup>162</sup>; and increases in the prevalence of disease<sup>163</sup>. More recently, endemic species turnover in Mexico has  
 31 been detected and attributed to climate change<sup>164</sup>. It is not surprising that species extinctions have not  
 32 yet been attributed to climate change<sup>86</sup>, since it generally takes decades for a species to become  
 33 completely extinct, and extinction commonly occurs as a result of a combination of factors<sup>165</sup>.

34 In addition, individual studies have begun to find relationships between climate trends and economic  
 35 growth rates in developing countries<sup>166, 167</sup>, even if global aggregate economic damages are not yet  
 36 attributable to climate change.

### 37 **6.2. Moderate risk at 1.6-2.6°C above preindustrial (and transition from Undetected to Moderate risk** 38 **between current temperatures and 1.6 °C)**

#### 39 **6.2.1. Definition of “increasing risk of extinction”**

1 In the IPCC Fourth and Fifth Assessment Reports and in this paper, ‘increased risk of extinction’ reflects  
2 that the literature indicates a future increased risk of extinction, but does not go as far as projecting  
3 actual extinction rates. The proportions of species that are thought to be *exposed* to this increased risk  
4 are quantified in much of this literature, but this does not imply actual extinction within a defined  
5 period, such as the date at which they become exposed. This literature referring to increased risk of  
6 extinction includes (but is not completely limited to) studies which show that species are projected to  
7 lose large proportions of their historic geographic range once climate has changed (in other words, their  
8 climatic niche is largely lost). Once a species is confined to a small area it becomes more vulnerable to  
9 extinction – for example as a result of a disease outbreak, an extreme weather event such as a  
10 prolonged drought, or due to local land use change caused by humans fragmenting or completely  
11 removing their habitat. Thus the International Union for the Conservation of Nature (IUCN) criteria for  
12 classifying a species as critically endangered includes geographic range as one of the key criteria,  
13 alongside factors related to population size and population dynamics  
14 ([http://www.iucnredlist.org/static/categories\\_criteria](http://www.iucnredlist.org/static/categories_criteria)). Having a limited geographic range does not  
15 necessarily mean that a species will in fact become extinct within a period shorter than it would  
16 otherwise have lasted, as extinction depends on many other factors, such as population dynamics, other  
17 effects of climatic change such as the impacts of extreme weather events, and climate effects mediated  
18 through interactions with other species (eg its predators, prey or pollinators) as well as non-climate  
19 drivers. The combined outcome of these factors might act to exacerbate or reduce the potential for  
20 extinction, and are not included in most of the quantifications of increased extinction risk found in the  
21 literature.

#### 22 *6.2.2. Impacts on biodiversity at 1.6-2.6° C warming relative to preindustrial*

23 There have been many studies on the potential impacts of climate change on the ranges of a number of  
24 species with varying levels of warming. AR4 concluded that approximately 20-30% of plant and animal  
25 species assessed are likely to be at increased risk of extinction for warming of 2-3°C above  
26 preindustrial<sup>29</sup>. A recent meta-analysis of 131 published projections found that extinction risk increased  
27 from 2.8% (current) to 5.2% at 2°C, 8.5% at 3°C and 16% at ~4°C across both widespread and endemic  
28 species<sup>32</sup>. However, endemic species were found to have a 6% higher extinction risk in this study.  
29 Differences between the levels of increases in extinction risk between studies stem almost entirely from  
30 assumptions about how much range loss equates to increased risk of extinction. Ref. 69, in an  
31 assessment of 48,786 widespread animal and plant species across the globe, projected that with a  
32 warming of 2°C above pre-industrial levels by the 2080s, more than half of the potential range would be  
33 lost for 23+/-4% of the plants and 13+/-3% of animals studied globally, allowing for realistic dispersal  
34 rates. The species distribution models on which the global assessments in refs. 69 and 32 were largely  
35 based cannot be applied to narrowly-restricted species, which in Sub-Saharan amphibians, for example,  
36 make up 54% of species and 92% of threatened species<sup>168</sup>. Therefore these estimates may under-  
37 represent the total risk.

38 Other studies that inform this risk judgment have found similar risks to Western Hemisphere birds,  
39 substantial impacts on specific animal and plant species ranges, and projected turnovers of large  
40 fractions of marine species assemblages. To date a minimum of 66,000 species have been examined

1 (exact number larger but unobtainable as there are overlaps in species between different studies). The  
 2 types of studies include species distribution models (also called ecological niche models), mechanistic  
 3 models, expert judgment with or without traits-based analysis, and species area curves. These models  
 4 include almost all birds, amphibians, and warm-water corals, large numbers of plants as well as  
 5 mammals, reptiles and butterflies<sup>27, 32, 69</sup>. For example, 23-25% of Western Hemisphere birds are  
 6 projected to be at increased risk of extinction under approximately 1.6°C of warming above  
 7 preindustrial<sup>169</sup>. The numbers of species at increasing risk of extinction varies widely in the literature  
 8 owing to differences in modeling technique, assumptions about dispersal, endemic (RFC 1) versus  
 9 widespread (RFC 4) species and, especially, thresholds used to establish increasing risk of extinction<sup>32</sup>.  
 10 Globally, even with a more restricted definition of threshold of change to classify a species as under  
 11 increased extinction risk, this translates to a near doubling of current extinction risk with 2°C above pre-  
 12 industrial. However, increased extinction risk is not the only consideration when looking at biodiversity  
 13 and, in particular, its provision of ecosystem services. Large-scale losses of widespread and common  
 14 species<sup>69</sup> implies significant erosion of ecosystem services operating through predation, pollination,  
 15 carbon removal, subsistence, etc. There are also projected large turnovers of up to 60% in marine  
 16 species assemblages (for 1.8 – 2.4°C of warming above pre-industrial levels; corresponding to SRES A1B,  
 17 B1 and A2 in the 2050s), combined with shrinkage of fish body weight of 14–24%<sup>170, 171</sup>.

### 18 *6.2.3. Economic damages between 0 and 3°C of warming above pre-industrial levels*

19 Economic impacts are not an important factor at low levels of warming for global aggregate impacts  
 20 because studies point to the combination of winners and losers from limited levels of warming. Winners  
 21 include countries with presently cool climates that may experience increases in agricultural or forest  
 22 production, reductions in the number of cold-related deaths, and decreases in energy expenditures for  
 23 heating. Losers include countries with presently hot climates, or with high vulnerability to moderate  
 24 changes in hazard distribution (e.g. low-lying areas and small islands), and people living in extreme  
 25 poverty who are vulnerable to any kind of environmental change. Some assessments indicate that, on  
 26 average, these impacts cancel out for low levels of warming. As warming increases, negative impacts  
 27 are driven by, for example, increased heat-related mortality and morbidity, decreased labor  
 28 productivity, increased energy demand for cooling, coastal flooding, crop loss, ecosystem damage, and  
 29 the potential for catastrophic impacts.

30 Few studies assess global aggregate damages due to climate change at levels of warming close to the  
 31 present level. Estimates of global economic damages generally project small negative impacts around  
 32 1°C warming above pre-industrial levels<sup>70, 172, 173</sup> (although two older studies project small positive  
 33 impacts). One study<sup>173</sup> found that 1°C of warming caused ~2% ± 1% (1σ) increase in global GDP due to  
 34 impacts on agriculture, forestry, biodiversity, sea level rise, human health, energy demand, water  
 35 resources. The benefits arise from the benefits of assumed CO<sub>2</sub> fertilization to agriculture and forestry  
 36 after allowing for adaptation, from the reduction in the number of cold deaths, and from reduction in  
 37 heating demand. Note that CO<sub>2</sub> fertilization is itself uncertain in magnitude, and that its effects might  
 38 be negated by the effects of climate change upon the frequency and intensity of extreme weather such  
 39 as heatwaves<sup>151</sup> and/or by increases in agricultural pests and diseases. That study also found that costs  
 40 of climate change at 1°C fall disproportionately on poorer regions, especially Africa and South/Southeast

1 Asia. Another study<sup>172</sup> examined the effects of 1°C warming using a quite different methodology,  
 2 assessing a cross-country panel of self-reported happiness, and finding a decrease in happiness  
 3 equivalent to a 0.4% loss of GDP from 1°C of warming. Based on these studies, we conclude that the  
 4 globally aggregated economic effect of 1°C of warming is small.

5 The estimate cited in the main text of 0-3% impact on GDP for warming of 1.9-3° C relative to  
 6 preindustrial is based on studies that either enumerate and add different impacts or examine multiple  
 7 impacts in a model of the full economy<sup>45, 174, 175, 176, 177, 178, 179</sup>. (Note that the limits of the 0-3% GDP range  
 8 do not coincide with the limits of the 1.9-3°C range; most estimates in the Tol (2009, 2014) analysis are  
 9 for either 2.5 or 3.0°C, and these span the range.). Many estimates depend on a large number of  
 10 disputable assumptions (see section 6.4 below).

### 11 **6.3. Transition to High risk around 3.6°C above preindustrial**

12 *Impacts on biodiversity at 3.6 C warming relative to preindustrial.* The judgment for this risk transition is  
 13 supported by many of the same studies used to support the judgment of Moderate risk at lower  
 14 temperatures, as well as an assessment of 48,786 species which showed potential range losses of >50%  
 15 for 57+/-6% of plants and 34+/-7% of animals studied. Results from the global assessment of 16,857  
 16 species<sup>69</sup> indicated even higher risks at a global temperature rise of 2.9-3.4°C above pre-industrial levels  
 17 (A2, 2090), with extinction risks applying to approximately 26 - 62% of the birds, 30-58% of the  
 18 amphibians and 42-65% of corals studied. Extinction risks are estimated to increase to 32-34% for  
 19 Western Hemisphere birds<sup>169</sup>. Furthermore, 10-20% of natural vegetation has been projected to be at  
 20 severe risk of ecosystem transformation (and hence disruption of ecosystem services) under a global  
 21 temperature rise of approximately 3°C above pre-industrial levels<sup>174</sup>.

22 A species does not need to be considered globally extinct for its decline to have an impact on ecosystem  
 23 services. The local loss of one or more species will also impact a range of ecosystem services. Where  
 24 many species potentially lose large fractions of their range, but are not considered at risk of extinction,  
 25 this may ultimately have an even greater impact on ecosystem services at a global level. For example,  
 26 for a temperature rise of 3.6°C above preindustrial, it was estimated that >50% of the potential range  
 27 would be lost for 57+/-6% of plants and 34+/-7% of animals studied, allowing for realistic dispersal  
 28 rates<sup>69</sup>. For a temperature rise of 3.6°C above preindustrial the figures increase to 57+/-5% of plants  
 29 and 34+/-7% of animals studied. Such biodiversity loss amongst widespread and common species  
 30 implies a major erosion of ecosystem functioning and services<sup>180, 181</sup>.

31 A recent global analysis showed that species that are widespread geographically, not only endemics  
 32 (which have tended to be the focus of many previous studies), are at risk of high levels of range loss  
 33 meaning that they would disappear from many of the areas they currently inhabit<sup>69</sup>. 32-34% of Western  
 34 Hemisphere birds are projected to be at risk of extinction under approximately 3°C of warming above  
 35 preindustrial<sup>169</sup>. Furthermore, 10-20% of natural vegetation has been projected to be at severe risk of  
 36 ecosystem transformation (and hence disruption of ecosystem services) under a global temperature rise  
 37 of approximately 3°C<sup>182</sup>. While all of these studies use a range of taxa-specific dispersal rates (also, rate  
 38 of climate change above), paleoecological evidence shows that dispersal in response to climate change  
 39 is species-specific and that climate changes in the past led to what are known as non-analogue



1 communities; that is, assemblages of species that are different than currently occurring assemblages.  
 2 This ultimately means that the interactions with ecosystem services (e.g., seed dispersal, pollination and  
 3 predation, especially of insect pests) between natural and agricultural systems will be different than  
 4 current.

5 Together this evidence results in a transition to high risk for aggregate global biodiversity at 3°C above  
 6 pre-industrial levels (Figure S2).

7 *Economic damages.* There are too few studies of aggregate economic damages to provide support for  
 8 the judgment of risks above 3°C relative to pre-industrial. Ref. 183, as aggregated by ref. 184, found  
 9 losses of happiness at 3.2°C equivalent to 12.4% of global GDP (or 11.5% as aggregated in ref. 70), while  
 10 ref. 178 found losses of 6.1% of GDP at 5.4°C warming relative to pre-industrial, driven primarily by labor  
 11 productivity decline (or 4.6% of GDP as aggregated by ref. 70).

12 The combination of the evidence about impacts on aggregate global biodiversity and the economy is  
 13 taken to result in a transition to high risk at 3.6°C, given that risks to the economy in the literature do  
 14 not reach the ‘High’ level for any level of warming assessed, and hence the ‘High’ risk point for the  
 15 combination of economic and biodiversity-related impacts is placed at a larger temperature change than  
 16 it would have been were it based upon a biodiversity assessment alone (see Supplemental Figure S2).

17

#### 18 **6.4. Limitations of estimates of global aggregate economic impacts**

19 Economists have long used benefit-cost integrated assessment models (IAMs) to estimate the global  
 20 aggregate economic impacts of climate change, employing simplified representations of the physical  
 21 impacts of climate change upon a range of sectors and regions<sup>177, 178, 185, 186, 187, 188</sup>. For example, some  
 22 IAMs include simple representations of the global aggregate impacts of climate-change induced sea  
 23 level rise, the effects of climate change upon crop yields and the agricultural economy, and effects of  
 24 climate change upon water resources.

25 There are significant weaknesses in aggregate economic damage estimates, as has been noted by  
 26 numerous authors<sup>71, 189, 190, 191</sup>. IAMs may not capture the full range of uncertainties in the projection of  
 27 climate change and its impacts. For example FUND<sup>188</sup> assumes positive effects of CO<sub>2</sub> fertilisation on  
 28 crops, whereas debate continues about whether such effects are likely to manifest in the field<sup>53</sup>, and  
 29 meanwhile observations have already detected negative impacts of global climate change on yields of  
 30 maize and wheat<sup>54, 192</sup>. No benefit-cost IAM has the temporal resolution to explicitly include the impacts  
 31 of individual extreme weather events, meaning that they are likely to underestimate the magnitude of  
 32 damages from climate change<sup>189, 193</sup>.

33 Further, IAMs omit several key processes<sup>190, 194</sup>, including impacts upon ecosystem services, which then  
 34 impact upon the economy. Non-market interactions between impacts are excluded<sup>30</sup>, and there is an  
 35 assumption that loss of ecosystem services (including for example water purification, watershed  
 36 preservation, flood prevention, carbon sequestration, crop pollination, coastal protection, regulation of  
 37 pests and diseases, recycling of waste nutrients<sup>195</sup>) can be fully compensated for by market services<sup>196,</sup>  
 38 <sup>197, 198</sup>. Another key issue omitted from many integrated models is a representation of large-scale

1 singular events and their economic consequences, which a small number of studies are beginning to  
2 explore<sup>74, 186, 199</sup>.

3 New approaches to assessing aggregate impacts use risk assessment frameworks, aggregating risks  
4 across sectors globally using physical metrics such as numbers of people at increased risk of various  
5 kinds of impacts, instead of using economic methods of aggregation. These are based on much more  
6 detailed physical process-based or econometric modelling of the impacts of climate change in multiple  
7 sectors globally<sup>200, 201</sup> or nationally (e.g., in the USA<sup>202, 203</sup>). These studies overcome some of these  
8 limitations and can explicitly include effects of extreme weather events such as storm surges and heat  
9 waves, but also do not provide comprehensive estimates of the total aggregate economic risk across all  
10 key impact categories.

11

## 12 **7. RFC 5: Risks associated with large-scale singular events**

### 13 **7.1. Transition from Moderate to High risk between 1.6 and 3.6°C above preindustrial**

14 Drawing on AR5 WGI chapter 13 (ref. 76, 13.4, 13.5), AR5 WGII Chapter 19 placed the transition from  
15 Moderate to High risk of a multi-meter sea level rise between 1.6-4.6 C above preindustrial based on  
16 modeling of the sensitivity of the Greenland ice sheet to future warming and paleoclimatic evidence of  
17 large sea level rise due to contributions from both Greenland and Antarctic ice sheets during the Last  
18 Interglacial (LIG). Modeling studies indicate a threshold for complete loss of GIS in the 1.0 (low  
19 confidence)-4.0 (medium confidence) °C range while paleoclimatic evidence supports a partial loss of  
20 GIS during the LIG when global mean temperature was less than 2 °C above preindustrial levels.

21 However, WGI had low confidence in the lower end of the temperature range which was determined by  
22 one study<sup>204</sup> which placed the threshold for complete loss of GIS around 1C compared to preindustrial.  
23 Furthermore, the relevance of the LIG analog is limited by differences in the paleo orbital forcing  
24 compared to future greenhouse gas forcing. In this paper, the transition from moderate to high risk is  
25 slightly modified to 1.6-4.0C to take the differences in confidence levels into account.

26 The role of the West Antarctic Ice Sheet (WAIS) entered into the Chapter 19 judgment in assessing the  
27 gradient of risk within the 1.6-4.6C range, with particular reference to studies supporting LIG sea level  
28 rise in the range of 6-9m<sup>189, 205, 206</sup>, a range which would require substantial contribution of both ice  
29 sheets. One comprehensive post-AR5 review of evidence from paleo-sea levels supports such a large  
30 LIG sea level rise<sup>207</sup>. Other post-AR5 studies provide additional evidence of an accelerating contribution  
31 from WAIS currently<sup>84</sup> with the potential for a loss equivalent to a 3m or larger sea level rise in a multi-  
32 century timescale<sup>83, 208</sup>. Refs. 84 and 208 are not specific as to the relation of possible rapid loss to  
33 future temperatures but additional evidence in this regard (for example, see ref. 209) could support  
34 revision of the RFC 5 risk to include the Very High risk category.

35 The judgment that there is a more rapid increase in risk between 1.6°C and 2.6°C reflects additional risk  
36 of a very large sea level rise based on the LIG ice loss from both ice sheets<sup>196, 205, 206, 207</sup> (ref. 57, Section  
37 5.6.2), when GMT was no more than 2°C warmer than preindustrial levels. This assessment of risk is  
38 based primarily on the *magnitude* and *irreversibility* of such sea level rise and the widespread exposure

1 and vulnerability of coastal settlements and ecosystems to such a rise. However, the slower the rate of  
2 rise, the more feasible becomes adaptation to reduce vulnerability and exposure.

### 3 **7.2. Rate of ice sheet loss and associated risk**

4 Particularly with regard to rate of change, the assessment of risk associated with RFC5 is not well-  
5 characterized. “Abrupt” is a relative term: Disintegration of an ice sheet may be abrupt in a geological  
6 sense but, if stretched over a millennium or more, sufficiently slow to allow human and societal  
7 adjustment on a timescale as occurs routinely for large scale social reorganizations for economic  
8 reasons. On the other hand, there is no evident way to quantify cultural losses from permanent loss of  
9 (current and future) antiquities to the sea no matter how slowly they occur. In addition, multi-meter sea  
10 level rise of the magnitude associated with ice sheet loss stretching over a few centuries, while very  
11 slow compared to many other climate system changes, could seriously challenge adaptive capacity and,  
12 as a result, be viewed as abrupt.

13

## 14 **8. Additional Metrics**

### 15 **8.1. Rate of climate change: limits to adaptation**

16 Successful (and in some cases unsuccessful) natural adaptation of organisms to past changes in global  
17 mean temperature of similar magnitudes to those projected over the next century resulted mostly from  
18 their movement. The unassisted rate of movement of populations of organisms across the landscape –  
19 whether by migration in the case of mobile organisms, or by successive cycles of growth, propagule  
20 production, dispersal, and establishment for sessile organisms – has theoretical limits, borne out by  
21 observations. Paleoclimatic data show that on average, the rate of temperature rise during the  
22 emergence from the last glacial period was much slower than the rate currently observed and projected  
23 for this century (there may have been periods in the Younger Dryas where rates approached the current  
24 rate<sup>210, 211</sup>). Paleo-ecological data show that the realized migration rates of plants, vertebrates and  
25 invertebrates vary greatly, and extinctions occurred in many groups during past climate changes<sup>86</sup>.

### 26 **8.2. Rate of climate change: dependence of rate of change on topography**

27 For example, for the RCP8.5 scenario, GMT change over the period 2050-2090 averages about  
28 0.065°C/yr. This translates to a mean climate velocity in flat landscapes of about 70 km/decade, while in  
29 mountainous landscapes the rate of horizontal movement is less – averaging about 2.5 km/decade. The  
30 global average for all landscapes for the RCP8.5 scenario is about 20 km/decade.

### 31 **8.3. Sea level rise: Transition to high risk**

32 Additional examples include: For 20 cm of sea level rise, with local subsidence and without upgrade in  
33 coastal defenses, more than \$1 trillion in assets could be lost annually in large cities alone<sup>212</sup>. For sea  
34 level rise between 20 and 60 cm, many ecosystems such as wetlands, coral reefs, estuaries and lagoons,  
35 and deltas would be at risk of widespread losses. Moreover, some vulnerability hotspots such as small  
36 islands would reach adaptation limits.

## 37 **9. RFCs and the vulnerability of socio-ecological systems**

1 *Examples of dependence of impacts on societal conditions.* For example, the population exposed to  
 2 future water scarcity is sensitive to population growth assumptions<sup>58</sup>, and sea-level rise impacts depend  
 3 on future coastal development and on capacity to invest in protection<sup>212, 213</sup>. The ability of species to  
 4 move can be impacted by barriers (e.g., cities, agriculturally transformed landscapes) that are linked to  
 5 differing socio-economic assumptions. More recently, projections of global numbers of people at risk of  
 6 hunger<sup>101, 102</sup> and water scarcity<sup>214</sup>, as well as the health burden attributable to childhood  
 7 undernutrition<sup>215</sup>, have been found to depend more on changes in societal conditions than on climate  
 8 change. Additionally, a projection of US population exposed to extreme heat found that demographic  
 9 change was as important as climate change to outcomes<sup>47</sup>.

10

## 11 **10. Figure 3, main text**

### 12 **10.1. Methodology**

13 Figure 3 is based on the numbers of people at risk of hunger reported in the scenarios described further  
 14 in the extension to the Figure 3 Caption below and in Table S1. To normalize for the uncertainty in  
 15 numbers of people at risk for a given climate outcome and level of exposure and vulnerability (E&V), we  
 16 express all results as the difference in the number at risk relative to a benchmark case, defined as a  
 17 medium E&V scenario with about 2.6 C warming. This normalization is applied within each subset of  
 18 scenarios carried out with the same crop model and economic (or integrated assessment) model,  
 19 climate model output, assumption about CO<sub>2</sub> fertilization, and assumption about adaptation. This  
 20 normalization controls for principal factors that lead to uncertainty in risk outcomes for the benchmark  
 21 case. However, results can be counter-intuitive in some cases. For example, subsets of scenarios that  
 22 include explicit adaptation lead to smaller apparent benefits of lower climate change or E&V, because  
 23 the main benefit of adaptation appears in the benchmark case to which other scenarios are normalized.  
 24 In addition, subsets of scenarios that include the positive effects of CO<sub>2</sub> fertilization can lead to  
 25 outcomes in which less climate change results in higher risks, because CO<sub>2</sub> levels are also lower and  
 26 therefore its positive effects are lessened. Finally, some low E&V scenarios could lead to apparently  
 27 reduced positive benefits of climate change (and associated CO<sub>2</sub> levels) simply because there are few  
 28 people at risk of hunger even before climate change (and CO<sub>2</sub>) effects are considered.

### 29 **10.2. Figure 3 caption**

30 The figure incorporates 24 scenarios from ref. 101: three different societal development pathways  
 31 differentiated by exposure and vulnerability of socio-ecological systems (SSPs 1, 2 and 3), four different  
 32 climate change outcomes for 2050 (RCP8.5, 6.0, 4.5 and 2.6, median of 8 GCMs each), and two  
 33 adaptation assumptions (with and without). Two scenarios are included from ref. 102: a medium  
 34 vulnerability societal future (SSP2) with two climate outcomes for 2050 (RCP8.5, RCP2.6, median of 12  
 35 different GCMs). These two scenarios are from the same author team as for ref. 91 but produce  
 36 outcomes with substantially lower numbers of people at risk of hunger. The primary reason is that in ref.  
 37 92, changes in the within-country distribution of per capita calorie consumption are assumed, while the  
 38 distribution is held fixed in ref. 91. This change leads to a substantial reduction in projected risk.  
 39 Fourteen scenarios are included from ref. 103: four different development pathways (SRES A1FI, A2, B1,  
 40 B2) with their associated climate outcomes in 2080 based on the HadCM3 model<sup>216</sup> with different crop

1 yield models and assumptions about CO<sub>2</sub> fertilization (AEZ and DSSAT models with CO<sub>2</sub> fertilization, 4  
2 scenarios each; AEZ and DSSAT models without CO<sub>2</sub> fertilization, 2 and 4 scenarios, respectively). A  
3 summary of quantitative outcomes for each scenario that is plotted in Figure 3 is provided in  
4 Supplementary Table 1.

5 For the results from ref. 101, all values are for 2050. Global Mean Temperature is from a corrected  
6 version of their Figure S7 provided by the author and represents the median of 8 GCMs. Global  
7 population is from Figure S6, and the change in hunger due to climate change is taken from Figure 5 as  
8 the median percent change in numbers of people at risk of hunger relative to a scenario with no climate  
9 change. The total population at risk of hunger is from Figure 3, summed from separate estimates for  
10 Developing and Transition countries.

11 For the results from ref. 102, all values are for 2050. Global Mean Temperature is from a corrected  
12 version of their Figure S8 provided by the author and represents the median of 12 GCMs. Global  
13 Population is from Figure S9 (the same as in ref. 101). The change in hunger due to climate change is  
14 taken from Figure 2 as the median change in numbers of people at risk of hunger relative to scenario  
15 with no climate change.

16 For the results from ref. 103, all values are for 2080. Global Mean Temperature is from HadCM3 results  
17 for these scenarios available from ref. 216. Population is taken from the SRES database at CIESIN.  
18 Numbers of people at risk of hunger are from Table 1 for 2080. Results for scenarios A1 and B1 with the  
19 AEZ-BLS impact model including CO<sub>2</sub> fertilization were not provided in that Table.

20 In general Fig. 3 and Supplementary Table 1 present results from different studies that have been  
21 harmonized to common categories of vulnerability and to a single scale of GMT. The vulnerability and  
22 GMT levels reported are approximate, as each study and scenario has its own approach to characterizing  
23 these factors.

24

25

26

1 **Supplementary Table 1:** Summary of scenario results supporting Figure 3 (a and b).

2


	Vulnerability	GMT	Total population at risk of hunger	Total pop at risk relative to SSP2-8.5	Additional pop at risk due to CC	Additional pop at risk compared to add'l pop at risk in SSP2-8.5
		C, rel. to 1981-2000	Millions	Millions	Millions	Millions
<b>Hasegawa et al, 2014</b>						
Current (2005)			830			
No climate change						
SSP1-Ref	L		400			
SSP2-Ref	M		530			
SSP3-Ref	H		940			
With climate change, without adaptation						
SSP1-8.5	L	2.1	452	-152	52	-22
SSP1-6.0	L	1.5	426	-178	26	-48
SSP1-4.5	L	1.6	435	-169	35	-39
SSP1-2.6	L	1.4	420	-185	20	-55
SSP2-8.5	M	2.1	604	0	74	0
SSP2-6.0	M	1.5	569	-36	39	-36
SSP2-4.5	M	1.6	578	-26	48	-26
SSP2-2.6	M	1.4	558	-47	28	-47
SSP3-8.5	H	2.1	1055	450	115	40
SSP3-6.0	H	1.5	1010	405	70	-5
SSP3-4.5	H	1.6	1026	421	86	11
SSP3-2.6	H	1.4	990	386	50	-24
With climate change, with adaptation						
SSP1-8.5	L	2.1	414	-136	14	-6
SSP1-6.0	L	1.5	409	-141	9	-11
SSP1-4.5	L	1.6	409	-141	9	-11
SSP1-2.6	L	1.4	404	-145	4	-15
SSP2-8.5	M	2.1	550	0	20	0
SSP2-6.0	M	1.5	544	-5	14	-5
SSP2-4.5	M	1.6	543	-7	13	-7
SSP2-2.6	M	1.4	536	-13	6	-13
SSP3-8.5	H	2.1	976	426	36	16
SSP3-6.0	H	1.5	968	419	28	9
SSP3-4.5	H	1.6	964	415	24	5
SSP3-2.6	H	1.4	952	403	12	-7
<b>Hasegawa et al, 2015</b>						
No climate change						
SSP2-Ref			90			
With climate change, without adaptation						
SSP2-8.5	M	2.1	92.2	0	2.2	0
SSP2-2.6	M	1.5	90.5	-2	0.5	-2

3

	Vulnerability	GMT	Total population at risk of hunger	Total pop at risk relative to SSP2-8.5	Additional pop at risk due to CC	Additional pop at risk compared to add'l pop at risk in SSP2-8.5
		C, rel. to 1981-2000	Millions	Millions	Millions	Millions
<b>Schmidhuber &amp; Tubiello, 2007</b>						
No climate change						
A1	L		108			
A2	H		768			
B1	L		91			
B2	M		233			
Climate change, 2080, $\Delta$ EZ-BLS, with CO2 fertilization						
A1	L	3.7	136	-108	28	17
A2	H	3	885	641	117	106
B1	L	1.8	99	-145	8	-3
B2	M	2.1	244	0	11	0
Climate change, 2080, $\Delta$ EZ-BLS, without CO2 fertilization						
A1	L	3.7				
A2	H	3	950	693	182	158
B1	L	1.8				
B2	M	2.1	257	0	24	0
Climate change, 2080, $\Delta$ SSAT-BLS, with CO2 fertilization						
A1	L	3.7	136	-85	28	40
A2	H	3	742	521	-26	-14
B1	L	1.8	102	-119	11	23
B2	M	2.1	221	0	-12	0
Climate change, 2080, $\Delta$ SSAT-BLS, without CO2 fertilization						
A1	L	3.7	370	-14	262	111
A2	H	3	1320	936	552	401
B1	L	1.8	125	-259	34	-117
B2	M	2.1	384	0	151	0

1


1 **Supplementary Figure 1: Process for assessing risks associated with Reasons for Concern in the IPCC**  
 2 **Fifth Assessment Report.** After criteria for key risks were defined, two branches of assessment were  
 3 carried out: Working Group 2 chapters identified key risks in their domains, which were then  
 4 synthesized into a set of eight overarching key risks (left branch in figure); Chapter 19 authors defined  
 5 specific risk levels and used selected studies to make a preliminary assessment in terms of GMT (right  
 6 branch). The two assessments were then merged to produce the final risk judgments.



7  
 8  
 9



1 **Supplementary Figure 2: Decomposition of burning ember for RFC4 into two primary key risks.**  
 2



3  
 4  
 5

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39

## References

1. United Nations Framework Convention on Climate Change. Report on the structured expert dialogue on the 2013–2015 review; 4 May 2015.
2. Tschakert P. 1.5 C or 2 C: a conduit’s view from the science-policy interface at COP20 in Lima, Peru. *Climate Change Responses* 2015, **2**(1): 1.
3. United Nations Framework Convention on Climate Change. Adoption of the Paris Agreement; 12 December 2015.
4. Yohe G. “Reasons for concern”(about climate change) in the United States. *Climatic Change* 2010, **99**(1-2): 295-302.
5. Gattuso J-P, Magnan A, Bille R, Cheung W, Howes E, Joos F, *et al.* Contrasting futures for ocean and society from different anthropogenic CO2 emissions scenarios. *Science* 2015, **349**(6243): aac4722-4721 - aac4722-4710.
6. Oppenheimer M, Campos M, Warren R, Birkmann J, Luber G, O’Neill B, *et al.* Emergent risks and key vulnerabilities. In: Field CB, Barros VR, Dokken DJ, Mach KJ, Mastrandrea MD, Bilir TE, *et al.* (eds). *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press: Cambridge, United Kingdom and New York, NY, USA, 2014, pp 1039-1099.
7. Renn O. Concepts of Risk: An Interdisciplinary Review Part 1: Disciplinary Risk Concepts. *Gaia-Ecological Perspectives for Science and Society* 2008, **17**(1): 50-66.
8. Cardona O-D, van Aalst MK, Birkmann J, Fordham M, McGregor G, Perez R, *et al.* Determinants of risk: exposure and vulnerability. In: Field CB, Barros V, Stocker TF, Qin D, Dokken DJ, Ebi KL, *et al.* (eds). *Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation. A Special Report of Working Groups I and II of the Intergovernmental Panel on Climate Change (IPCC)*. Cambridge University Press: Cambridge University Press, Cambridge, UK, and New York, NY, USA, 2012, pp 65-108.
9. Cutter SL. Societal responses to environmental hazards. *International Social Science Journal* 1996, **48**(150): 525-536.

- 1 10. Agard J, Schipper E, Birkmann J, Campos M, Dubeux C, Nojiri Y, *et al.* Annex II: Glossary. In:  
2 Barros VR, Field CB, Dokken DJ, Mastrandrea MD, Mach KJ, Bilir TE, *et al.* (eds). *Climate Change*  
3 *2014: Impacts, Adaptation, and Vulnerability. Part B: Regional Aspects. Contribution of Working*  
4 *Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change.*  
5 Cambridge University Press: Cambridge, United Kingdom and New York, NY, USA, 2014, pp  
6 1757-1776.
- 7
- 8 11. Lavell A, Oppenheimer M, Diop C, Hess J, Lempert R, Li J, *et al.* Climate change: new dimensions  
9 in disaster risk, exposure, vulnerability, and resilience. In: Field CB, Barros V, Stocker TF, Qin D,  
10 Dokken DJ, Ebi KL, *et al.* (eds). *Managing the Risks of Extreme Events and Disasters to Advance*  
11 *Climate Change Adaptation. A Special Report of Working Groups I and II of the*  
12 *Intergovernmental Panel on Climate Change (IPCC).* Cambridge University Press: Cambridge  
13 University Press, Cambridge, UK, and New York, NY, USA, 2012, pp 25-64.
- 14
- 15 12. Patwardhan A, Semenov S, Schnieder S, Burton I, Magadza C, Oppenheimer M, *et al.* Assessing  
16 key vulnerabilities and the risk from climate change. In: Parry ML, Canziani OF, Palutikof JP, van  
17 der Linden PJ, Hanson CE (eds). *Change 2007: Impacts, Adaptation and Vulnerability.*  
18 *Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental*  
19 *Panel on Climate Change.* Cambridge University Press: Cambridge, UK, 2007, pp 779-810.
- 20
- 21 13. Smith JB, Schneider SH, Oppenheimer M, Yohe GW, Hare W, Mastrandrea MD, *et al.* Assessing  
22 dangerous climate change through an update of the Intergovernmental Panel on Climate  
23 Change (IPCC) "reasons for concern". *Proceedings of the National Academy of Sciences* 2009,  
24 **106**(11): 4133-4137.
- 25
- 26 14. Cramer W, Yohe G, Auffhammer M, Huggel C, Molau U, Dias MdS, *et al.* Detection and  
27 attribution of observed impacts. In: Field CB, Barros VR, Dokken DJ, Mach KJ, Mastrandrea MD,  
28 Bilir TE, *et al.* (eds). *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global*  
29 *and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the*  
30 *Intergovernmental Panel on Climate Change.* Cambridge University Press: Cambridge, United  
31 Kingdom and New York, NY, USA, 2014, pp 979-1038.
- 32
- 33 15. Myers N, Mittermeier RA, Mittermeier CG, Da Fonseca GA, Kent J. Biodiversity hotspots for  
34 conservation priorities. *Nature* 2000, **403**(6772): 853-858.
- 35
- 36 16. Smith JB, Schellnhuber H-J, Mirza MMQ, Fankhauser S, Leemans R, Lin E, *et al.* Vulnerability to  
37 climate change and reasons for concern: a synthesis. In: McCarthy JJ, Canziani OF, Leary NA,  
38 Dokken DJ, White KS (eds). *Climate Change 2001: Impacts, Adaptation, and Vulnerability.*  
39 Cambridge University Press: Cambridge, UK, 2001, pp 913-967.
- 40
- 41 17. Bindoff NL, Stott PA, AchutaRao M, Allen MR, Gillett N, Gutzler D, *et al.* Detection and  
42 attribution of climate change: from global to regional. In: Stocker TF, Qin D, Plattner G-K, Tignor  
43 M, Allen D, Boschung J, *et al.* (eds). *Climate Change 2013: The Physical Science Basis.*

- 1            *Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel*  
2            *on Climate Change*. Cambridge University Press: Cambridge, United Kingdom and New York, NY,  
3            USA, 2013, pp 867-952.
- 4
- 5    18.    Gooseff MN, Balsler A, Bowden WB, Jones JB. Effects of hillslope thermokarst in northern Alaska.  
6            *Eos, Transactions American Geophysical Union* 2009, **90**(4): 29-30.
- 7
- 8    19.    Karl TR, Melillo JM, Peterson TC (eds). *Global Climate Change Impacts in the United States*.  
9            Cambridge University Press: New York, NY, 2009.
- 10
- 11   20.    Jia GJ, Epstein HE, Walker DA. Vegetation greening in the Canadian Arctic related to decadal  
12            warming. *Journal of Environmental Monitoring* 2009, **11**(12): 2231-2238.
- 13
- 14   21.    Myers-Smith IH, Forbes BC, Wilmking M, Hallinger M, Lantz T, Blok D, *et al.* Shrub expansion in  
15            tundra ecosystems: dynamics, impacts and research priorities. *Environmental Research Letters*  
16            2011, **6**(4): 045509.
- 17
- 18   22.    Post E, Forchhammer MC, Bret-Harte MS, Callaghan TV, Christensen TR, Elberling B, *et al.*  
19            Ecological dynamics across the Arctic associated with recent climate change. *Science* 2009,  
20            **325**(5946): 1355-1358.
- 21
- 22   23.    Hoegh-Guldberg O, Mumby PJ, Hooten AJ, Steneck RS, Greenfield P, Gomez E, *et al.* Coral reefs  
23            under rapid climate change and ocean acidification. *Science* 2007, **318**(5857): 1737-1742.
- 24
- 25   24.    Baker AC, Glynn PW, Riegl B. Climate change and coral reef bleaching: An ecological assessment  
26            of long-term impacts, recovery trends and future outlook. *Estuarine, Coastal and Shelf Science*  
27            2008, **80**(4): 435-471.
- 28
- 29   25.    Veron J, Hoegh-Guldberg O, Lenton T, Lough J, Obura D, Pearce-Kelly P, *et al.* The coral reef  
30            crisis: The critical importance of <350ppm CO<sub>2</sub>. *Marine Pollution Bulletin* 2009, **58**(10): 1428-  
31            1436.
- 32
- 33   26.    Gattuso JP, Hoegh-Guldberg O, Portner HO. Cross Chapter Box on Coral Reefs. In: Field CB,  
34            Barros VR, Dokken DJ, Mach KJ, Mastrandrea MD, Bilir TE, *et al.* (eds). *Climate Change 2014:*  
35            *Impacts, Adaptation, And Vulnerability: Part A: Global And Sectoral Aspects. Contribution Of*  
36            *Working Group II To The Fifth Assessment Report Of The Intergovernmental Panel On Climate*  
37            *Change*. Cambridge University Press: Cambridge, United Kingdom and New York, NY, USA, 2014,  
38            pp 97-100.

39

- 1 27. Foden WB, Butchart SH, Stuart SN, Vié J-C, Akçakaya HR, Angulo A, *et al.* Identifying the world's  
2 most climate change vulnerable species: a systematic trait-based assessment of all birds,  
3 amphibians and corals. *Plos One* 2013, **8**(6): e65427.
- 4
- 5 28. Rode KD, Robbins CT, Nelson L, Amstrup SC. Can polar bears use terrestrial foods to offset lost  
6 ice - based hunting opportunities? *Frontiers in Ecology and the Environment* 2015, **13**(3): 138-  
7 145.
- 8
- 9 29. Fischlin A, Midgley GF, Price J, Leemans R, Gopal B, Turley C, *et al.* Ecosystems, their properties,  
10 goods and services. In: Parry ML, Canziani OF, Palutikof JP, van der Linden PJ, Hanson CE (eds).  
11 *Change 2007: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the*  
12 *Fourth Assessment Report of the Intergovernmental Panel on Climate Change.* Cambridge  
13 University Press: Cambridge, UK, 2007, pp 211-272.
- 14
- 15 30. Warren R, Price J, Fischlin A, de la Nava Santos S, Midgley G. Increasing impacts of climate  
16 change upon ecosystems with increasing global mean temperature rise. *Climatic Change* 2011,  
17 **106**(2): 141-177.
- 18
- 19 31. Malcolm JR, Liu C, Neilson RP, Hansen L, Hannah L. Global warming and extinctions of endemic  
20 species from biodiversity hotspots. *Conservation Biology* 2006, **20**(2): 538-548.
- 21
- 22 32. Urban MC. Accelerating extinction risk from climate change. *Science* 2015, **348**(6234): 571-573.
- 23
- 24 33. Wong PP, Losada IJ, Gattuso J, Hinkel J, Khattabi A, McInnes K, *et al.* Coastal systems and low-  
25 lying areas. In: Field CB, Barros VR, Dokken DJ, Mach KJ, Mastrandrea MD, Bilir TE, *et al.* (eds).  
26 *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral*  
27 *Aspects. Contribution of Working Group II to the Fifth Assessment Report of the*  
28 *Intergovernmental Panel on Climate Change.* Cambridge University Press: Cambridge, United  
29 Kingdom and New York, NY, USA, 2014, pp 361-409.
- 30
- 31 34. van Vuuren DP, Carter TR. Climate and socio-economic scenarios for climate change research  
32 and assessment: reconciling the new with the old. *Climatic Change* 2014, **122**(3): 415-429.
- 33
- 34 35. Segan DB, Hole DG, Donatti CI, Zganjar C, Martin S, Butchart SH, *et al.* Considering the impact of  
35 climate change on human communities significantly alters the outcome of species and site -  
36 based vulnerability assessments. *Diversity and Distributions* 2015, **21**(9): 1101-1111.
- 37
- 38 36. Visser ME. Keeping up with a warming world; assessing the rate of adaptation to climate  
39 change. *Proceedings of the Royal Society of London B: Biological Sciences* 2008, **275**(1635): 649-  
40 659.

- 1  
2 37. Hoffmann AA, Sgrò CM. Climate change and evolutionary adaptation. *Nature* 2011, **470**(7335):  
3 479-485.
- 4  
5 38. Strong AE, Liu G, Skirving W, Eakin CM. NOAA's Coral Reef Watch program from satellite  
6 observations. *Annals of GIS* 2011, **17**(2): 83-92.
- 7  
8 39. Smith K, Woodward A, Campell-Lendrum D, Tobago) DDCTa, Honda Y, Liu Q, *et al.* Human  
9 health: impacts adaptation and co-benefits. In: Field CB, Barros VR, Dokken DJ, Mach KJ,  
10 Mastrandrea MD, Bilir TE, *et al.* (eds). *Climate Change 2014: Impacts, Adaptation, and*  
11 *Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth*  
12 *Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University  
13 Press: Cambridge, United Kingdom and New York, NY, USA, 2014, pp 709-754.
- 14  
15 40. Fischer EM, Knutti R. Anthropogenic contribution to global occurrence of heavy-precipitation  
16 and high-temperature extremes. *Nature Climate Change* 2015, **5**(6): 560-564.
- 17  
18 41. Christidis N, Stott PA, Jones GS, Shiogama H, Nozawa T, Luterbacher J. Human activity and  
19 anomalously warm seasons in Europe. *International Journal of Climatology* 2012, **32**(2): 225-  
20 239.
- 21  
22 42. Kirtman B, Power S, Adedoyin J, Boer G, Bojariu R, Camilloni I, *et al.* Near-term climate change:  
23 projections and predictability. In: Stocker TF, Qin D, Plattner G-K, Tignor M, Allen D, Boschung J,  
24 *et al.* (eds). *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to*  
25 *the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge  
26 University Press: Cambridge, United Kingdom and New York, NY, USA, 2013, pp 953-1028.
- 27  
28 43. Field CB, Barros V, Stocker TF, Qin D, Dokken DJ, Ebi KL, *et al.* (eds). *Managing the Risks of*  
29 *Extreme Events and Disasters to Advance Climate Change Adaptation. A Special Report of*  
30 *Working Groups I and II of the Intergovernmental Panel on Climate Change*. Cambridge  
31 University Press: Cambridge, UK, and New York, NY, USA, 2012.
- 32  
33 44. Peduzzi P, Chatenoux B, Dao H, De Bono A, Herold C, Kossin J, *et al.* Global trends in tropical  
34 cyclone risk. *Nature Climate Change* 2012, **2**(4): 289-294.
- 35  
36 45. Mendelsohn R, Emanuel K, Chonabayashi S, Bakkensen L. The impact of climate change on  
37 global tropical cyclone damage. *Nature Climate Change* 2012, **2**(3): 205-209.
- 38  
39 46. Dong W, Liu Z, Liao H, Tang Q, Li Xe. New climate and socio-economic scenarios for assessing  
40 global human health challenges due to heat risk. *Climatic Change* 2015, **130**(4): 505-518.

- 1  
2 47. Jones B, O'Neill BC, McDaniel L, McGinnis S, Mearns LO, Tebaldi C. Future population exposure  
3 to US heat extremes. *Nature Climate Change* 2015, **5**(7): 652-655.
- 4  
5 48. Birkmann J, Cutter SL, Rothman DS, Welle T, Garschagen M, van Ruijven B, *et al.* Scenarios for  
6 vulnerability: opportunities and constraints in the context of climate change and disaster risk.  
7 *Climatic Change* 2015, **133**(1): 53-68.
- 8  
9 49. Visser H, Petersen AC, Ligtoet W. On the relation between weather-related disaster impacts,  
10 vulnerability and climate change. *Climatic Change* 2014, **125**(3-4): 461-477.
- 11  
12 50. Fouillet A, Rey G, Wagner V, Laaidi K, Empereur-Bissonnet P, Le Tertre A, *et al.* Has the impact of  
13 heat waves on mortality changed in France since the European heat wave of summer 2003? A  
14 study of the 2006 heat wave. *International Journal of Epidemiology* 2008, **37**(2): 309-317.
- 15  
16 51. Garschagen M, Romero-Lankao P. Exploring the relationships between urbanization trends and  
17 climate change vulnerability. *Climatic Change* 2015, **133**(1): 37-52.
- 18  
19 52. Welle T, Birkmann J. The World Risk Index—An Approach to Assess Risk and Vulnerability on a  
20 Global Scale. *Journal of Extreme Events* 2015, **2**(01): 1550003.
- 21  
22 53. Porter JR, Xie L, Challinor AJ, Cochrane K, Howden SM, Iqbal MM, *et al.* Food Security and Food  
23 Production Systems. In: Field CB, Barros VR, Dokken DJ, Mach KJ, Mastrandrea MD, Bilir TE, *et al.*  
24 (eds). *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral*  
25 *Aspects. Contribution of Working Group II to the Fifth Assessment Report of the*  
26 *Intergovernmental Panel on Climate Change*. Cambridge University Press: Cambridge, United  
27 Kingdom and New York, NY, USA, 2014, pp 485-533.
- 28  
29 54. Lobell DB, Schlenker W, Costa-Roberts J. Climate trends and global crop production since 1980.  
30 *Science* 2011, **333**(6042): 616-620.
- 31  
32 55. Challinor A, Watson J, Lobell D, Howden S, Smith D, Chhetri N. A meta-analysis of crop yield  
33 under climate change and adaptation. *Nature Climate Change* 2014, **4**: 287-291.
- 34  
35 56. Knox J, Hess T, Daccache A, Wheeler T. Climate change impacts on crop productivity in Africa  
36 and South Asia. *Environmental Research Letters* 2012, **7**(3): 034032.
- 37  
38 57. Rosenzweig C, Elliott J, Deryng D, Ruane AC, Müller C, Arneth A, *et al.* Assessing agricultural risks  
39 of climate change in the 21st century in a global gridded crop model intercomparison.  
40 *Proceedings of the National Academy of Sciences* 2014, **111**(9): 3268-3273.

- 1  
2 58. Gosling SN, Arnell NW. A global assessment of the impact of climate change on water scarcity.  
3 *Climatic Change* 2013: 1-15.
- 4  
5 59. Schewe J, Heinke J, Gerten D, Haddeland I, Arnell NW, Clark DB, *et al.* Multimodel assessment of  
6 water scarcity under climate change. *Proceedings of the National Academy of Sciences* 2014,  
7 **111**(9): 3245-3250.
- 8  
9 60. Thornton PK, Jones PG, Ericksen PJ, Challinor AJ. Agriculture and food systems in sub-Saharan  
10 Africa in a 4 C+ world. *Philosophical Transactions of the Royal Society of London A:*  
11 *Mathematical, Physical and Engineering Sciences* 2011, **369**(1934): 117-136.
- 12  
13 61. Niang I, Ruppel OC, Abdrabo MA, Essel A, Lennard C, Padgham J, *et al.* Africa. In: Barros VR, Field  
14 CB, Dokken DJ, Mastrandrea MD, Mach KJ, Bilir TE, *et al.* (eds). *Climate Change 2014: Impacts,*  
15 *Adaptation, and Vulnerability. Part B: Regional Aspects. Contribution of Working Group II to the*  
16 *Fifth Assessment Report of the Intergovernmental Panel on Climate Change.* Cambridge  
17 University Press: Cambridge, United Kingdom and New York, NY, USA, 2014, pp 1199-1265.
- 18  
19 62. Carleton TA, Hsiang SM. Social and economic impacts of climate. *Science* 2016, **353**(6304).
- 20  
21 63. Dell M, Jones BF, Olken BA. What Do We Learn from the Weather? The New Climate-Economy  
22 Literature. *Journal of Economic Literature* 2014, **52**(3): 740-798.
- 23  
24 64. Hallegatte S, Bangalore M, Bonzanigo L, Fay M, Kane T, Narloch U, *et al.* *Shock Waves: Managing*  
25 *the impacts of climate change on Poverty.* World Bank Publications, 2015.
- 26  
27 65. Nelson GC, Valin H, Sands RD, Havlík P, Ahammad H, Deryng D, *et al.* Climate change effects on  
28 agriculture: Economic responses to biophysical shocks. *Proceedings of the National Academy of*  
29 *Sciences* 2014, **111**(9): 3274-3279.
- 30  
31 66. Parfitt J, Barthel M, Macnaughton S. Food waste within food supply chains: quantification and  
32 potential for change to 2050. *Philosophical Transactions of the Royal Society of London B:*  
33 *Biological Sciences* 2010, **365**(1554): 3065-3081.
- 34  
35 67. Kummu M, De Moel H, Porkka M, Siebert S, Varis O, Ward P. Lost food, wasted resources: Global  
36 food supply chain losses and their impacts on freshwater, cropland, and fertiliser use. *Science of*  
37 *the Total Environment* 2012, **438**: 477-489.
- 38  
39 68. Cassidy ES, West PC, Gerber JS, Foley JA. Redefining agricultural yields: from tonnes to people  
40 nourished per hectare. *Environmental Research Letters* 2013, **8**(3): 034015.



- 1  
2 69. Warren R, VanDerWal J, Price J, Welbergen J, Atkinson I, Ramirez-Villegas J, *et al.* Quantifying  
3 the benefit of early climate change mitigation in avoiding biodiversity loss. *Nature Climate*  
4 *Change* 2013, **3**(7): 678-682.
- 5  
6 70. Tol RS. Correction and update: The economic effects of climate change. *The Journal of Economic*  
7 *Perspectives* 2014, **28**(2): 221-225.
- 8  
9 71. Revesz RL, Howard PH, Arrow K, Goulder LH, Kopp RE, Livermore MA, *et al.* Global warming:  
10 Improve economic models of climate change. *Nature* 2014, **508**(7495): 173-175.
- 11  
12 72. Ockendon N, Baker DJ, Carr JA, White EC, Almond RE, Amano T, *et al.* Mechanisms underpinning  
13 climatic impacts on natural populations: altered species interactions are more important than  
14 direct effects. *Global Change Biol* 2014, **20**(7): 2221-2229.
- 15  
16 73. Lenton TM, Held H, Kriegler E, Hall JW, Lucht W, Rahmstorf S, *et al.* Tipping elements in the  
17 Earth's climate system. *Proceedings of the National Academy of Sciences* 2008, **105**(6): 1786-  
18 1793.
- 19  
20 74. Kopp RE, Shwom R, Wagner G, Yuan J. Tipping elements and climate-economic shocks: Pathways  
21 toward integrated assessment. *Earth's Future* 2016: 346-372.
- 22  
23 75. Collins M, Knutti R, Arblaster J, Dufresne J-L, Fichet T, Friedlingstein P, *et al.* Long-term climate  
24 change: projections, commitments and irreversibility. In: Stocker TF, Qin D, Plattner G-K, Tignor  
25 M, Allen D, Boschung J, *et al.* (eds). *Climate Change 2013: The Physical Science Basis.*  
26 *Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel*  
27 *on Climate Change.* Cambridge University Press: Cambridge, United Kingdom and New York, NY,  
28 USA, 2013, pp 1029-1136.
- 29  
30 76. Church JA, Clark PU, Cazenave A, Gregory JM, Jevrejeva S, Levermann A, *et al.* Sea level change.  
31 In: Stocker TF, Qin D, Plattner G-K, Tignor M, Allen D, Boschung J, *et al.* (eds). *Climate Change*  
32 *2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report*  
33 *of the Intergovernmental Panel on Climate Change.* Cambridge University Press: Cambridge,  
34 United Kingdom and New York, NY, USA, 2013, pp 1137-1216.
- 35  
36 77. Intergovernmental Panel on Climate Change. Summary for Policy Makers. In: Stocker TF, Qin D,  
37 Plattner G-K, Tignor M, Allen D, Boschung J, *et al.* (eds). *Climate Change 2013: The Physical*  
38 *Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the*  
39 *Intergovernmental Panel on Climate Change.* Cambridge University Press: Cambridge, United  
40 Kingdom and New York, NY, USA, 2013, pp 3-29.

41

- 1 78. Masson-Delmotte V, Schulz M, Abe-Ouchi A, Beer J, Ganopolski A, González Rouco J, *et al.*  
2 Information from paleoclimate archives. In: Stocker TF, Qin D, Plattner G-K, Tignor M, Allen D,  
3 Boschung J, *et al.* (eds). *Climate Change 2013: The Physical Science Basis. Contribution of*  
4 *Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate*  
5 *Change*. Cambridge University Press: Cambridge, United Kingdom and New York, NY, USA, 2013,  
6 pp 383-464.
- 7
- 8 79. Turney CS, Jones RT. Does the Agulhas Current amplify global temperatures during super -  
9 interglacials? *Journal of Quaternary Science* 2010, **25**(6): 839-843.
- 10
- 11 80. Cornford SL, Martin D, Payne A, Ng E, Le Brocq A, Gladstone R, *et al.* Century-scale simulations  
12 of the response of the West Antarctic Ice Sheet to a warming climate. *Cryosphere Discussions*  
13 *(The)* 2015, **9**: 1579-1600.
- 14
- 15 81. Intergovernmental Panel on Climate Change. Climate Change 2014: Synthesis Report.  
16 Contribution of Working Groups I, II and III to the Fifth Assessment Report of the  
17 Intergovernmental Panel on Climate Change. Geneva, Switzerland: Intergovernmental Panel on  
18 Climate Change; 2014.
- 19
- 20 82. Pollard D, Chang W, Haran M, Applegate P, DeConto R. Large ensemble modeling of last  
21 deglacial retreat of the West Antarctic Ice Sheet: comparison of simple and advanced statistical  
22 techniques. *Geoscientific Model Development Discussions* 2015, **8**: 9925-9963.
- 23
- 24 83. Winkelmann R, Levermann A, Ridgwell A, Caldeira K. Combustion of available fossil fuel  
25 resources sufficient to eliminate the Antarctic Ice Sheet. *Science Advances* 2015, **1**(8): e1500589.
- 26
- 27 84. Rignot E, Mouginot J, Morlighem M, Seroussi H, Scheuchl B. Widespread, rapid grounding line  
28 retreat of Pine Island, Thwaites, Smith, and Kohler glaciers, West Antarctica, from 1992 to 2011.  
29 *Geophysical Research Letters* 2014, **41**(10): 3502-3509.
- 30
- 31 85. Oppenheimer M, Little CM, Cooke RM. Expert judgement and uncertainty quantification for  
32 climate change. *Nature Climate Change* 2016, **6**(5): 445-451.
- 33
- 34 86. Settele J, Scholes R, Betts RA, Bunn S, Leadley P, Nepstad D, *et al.* Terrestrial and inland water  
35 systems. In: Field CB, Barros VR, Dokken DJ, Mach KJ, Mastrandrea MD, Bilir TE, *et al.* (eds).  
36 *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral*  
37 *Aspects. Contribution of Working Group II to the Fifth Assessment Report of the*  
38 *Intergovernmental Panel on Climate Change*. Cambridge University Press: Cambridge, United  
39 Kingdom and New York, NY, USA, 2014, pp 271-359.

40

- 1 87. Burrows MT, Schoeman DS, Richardson AJ, Molinos JG, Hoffmann A, Buckley LB, *et al.*  
2 Geographical limits to species-range shifts are suggested by climate velocity. *Nature* 2014,  
3 **507**(7493): 492-495.
- 4
- 5 88. Burrows MT, Schoeman DS, Buckley LB, Moore P, Poloczanska ES, Brander KM, *et al.* The pace of  
6 shifting climate in marine and terrestrial ecosystems. *Science* 2011, **334**(6056): 652-655.
- 7
- 8 89. Chen I-C, Hill JK, Ohlemüller R, Roy DB, Thomas CD. Rapid range shifts of species associated with  
9 high levels of climate warming. *Science* 2011, **333**(6045): 1024-1026.
- 10
- 11 90. Dobrowski SZ, Abatzoglou J, Swanson AK, Greenberg JA, Mynsberge AR, Holden ZA, *et al.* The  
12 climate velocity of the contiguous United States during the 20th century. *Global Change Biol*  
13 2013, **19**(1): 241-251.
- 14
- 15 91. Feeley KJ, Rehm EM. Amazon's vulnerability to climate change heightened by deforestation and  
16 man - made dispersal barriers. *Global Change Biol* 2012, **18**(12): 3606-3614.
- 17
- 18 92. Loarie SR, Duffy PB, Hamilton H, Asner GP, Field CB, Ackerly DD. The velocity of climate change.  
19 *Nature* 2009, **462**(7276): 1052-1055.
- 20
- 21 93. Sandel B, Arge L, Dalsgaard B, Davies RG, Gaston KJ, Sutherland WJ, *et al.* The influence of Late  
22 Quaternary climate-change velocity on species endemism. *Science* 2011, **334**(6056): 660-664.
- 23
- 24 94. Bopp L, Resplandy L, Orr JC, Doney SC, Dunne JP, Gehlen M, *et al.* Multiple stressors of ocean  
25 ecosystems in the 21st century: projections with CMIP5 models. *Biogeosciences* 2013, **10**: 6225–  
26 6245.
- 27
- 28 95. Pörtner H-O, Karl DM, Boyd PW, Cheung W, Lluch-Cota S, Nojiri Y, *et al.* Ocean systems. In: Field  
29 CB, Barros VR, Dokken DJ, Mach KJ, Mastrandrea MD, Bilir TE, *et al.* (eds). *Climate Change 2014:*  
30 *Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of*  
31 *Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate*  
32 *Change*. Cambridge University Press: Cambridge, United Kingdom and New York, NY, USA, 2014,  
33 pp 411-484.
- 34
- 35 96. Bednaršek N, Feely R, Reum J, Peterson B, Menkel J, Alin S, *et al.* *Limacina helicina* shell  
36 dissolution as an indicator of declining habitat suitability owing to ocean acidification in the  
37 California Current Ecosystem. *Proceedings of the Royal Society of London B: Biological Sciences*  
38 2014, **281**(1785): 20140123.

39

- 1 97. Wittmann AC, Pörtner H-O. Sensitivities of extant animal taxa to ocean acidification. *Nature*  
2 *Climate Change* 2013, **3**(11): 995-1001.
- 3
- 4 98. Gattuso JP, Brewer PG, Hoegh-Guldberg O, Kleypas JA, Pörtner H-O. Cross Chapter Box on Ocean  
5 Acidification. In: Field CB, Barros VR, Dokken DJ, Mach KJ, Mastrandrea MD, Bilir TE, *et al.* (eds).  
6 *Climate Change 2014: Impacts, Adaptation, And Vulnerability. Part A: Global And Sectoral*  
7 *Aspects. Contribution Of Working Group II To The Fifth Assessment Report Of The*  
8 *Intergovernmental Panel On Climate Change*. Cambridge University Press: Cambridge, United  
9 Kingdom and New York, NY, USA, 2014, pp 129-131.
- 10
- 11 99. Hinkel J, van Vuuren DP, Nicholls RJ, Klein RJ. The effects of adaptation and mitigation on coastal  
12 flood impacts during the 21st century. An application of the DIVA and IMAGE models. *Climatic*  
13 *Change* 2013, **117**(4): 783-794.
- 14
- 15 100. Birkmann J, Welle T. Assessing the risk of loss and damage: exposure, vulnerability and risk to  
16 climate-related hazards for different country classifications. *International Journal of Global*  
17 *Warming* 2015, **8**(2): 191-212.
- 18
- 19 101. Hasegawa T, Fujimori S, Shin Y, Takahashi K, Masui T, Tanaka A. Climate change impact and  
20 adaptation assessment on food consumption utilizing a new scenario framework. *Environmental*  
21 *Science & Technology* 2013, **48**(1): 438-445.
- 22
- 23 102. Hasegawa T, Fujimori S, Shin Y, Tanaka A, Takahashi K, Masui T. Consequence of Climate  
24 Mitigation on the Risk of Hunger. *Environmental Science & Technology* 2015, **49**(12): 7245-7253.
- 25
- 26 103. Schmidhuber J, Tubiello FN. Global food security under climate change. *Proceedings of the*  
27 *National Academy of Sciences* 2007, **104**(50): 19703-19708.
- 28
- 29 104. Intergovernmental Panel on Climate Change. Summary for Policymakers. In: Field CB, Barros VR,  
30 Dokken DJ, Mach KJ, Mastrandrea MD, Bilir TE, *et al.* (eds). *Climate Change 2014: Impacts,*  
31 *Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working*  
32 *Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*.  
33 Cambridge University Press: Cambridge, United Kingdom and New York, NY, USA, 2014, pp 1-32.
- 34
- 35 105. Hartmann DL, Klein Tank AMG, Rusticucci M, Alexander LV, Brönnimann S, Charabi Y, *et al.*  
36 Observations: Atmosphere and Surface. In: Stocker TF, Qin D, Plattner G-K, Tignor M, Allen D,  
37 Boschung J, *et al.* (eds). *Climate Change 2013: The Physical Science Basis. Contribution of*  
38 *Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate*  
39 *Change*, vol. 159-254. Cambridge University Press: Cambridge, United Kingdom and New York,  
40 NY, USA, 2013.

41

- 1 106. Williams SE, Bolitho EE, Fox S. Climate change in Australian tropical rainforests: an impending  
2 environmental catastrophe. *Proceedings of the Royal Society of London B: Biological Sciences*  
3 2003, **270**(1527): 1887-1892.
- 4
- 5 107. Allen D, Whitfield P, Werner A. Groundwater level responses in temperate mountainous terrain:  
6 regime classification, and linkages to climate and streamflow. *Hydrological Processes* 2010,  
7 **24**(23): 3392-3412.
- 8
- 9 108. Gottfried M, Pauli H, Futschik A, Akhalkatsi M, Barančok P, Alonso JLB, *et al.* Continent-wide  
10 response of mountain vegetation to climate change. *Nature Climate Change* 2012, **2**(2): 111-  
11 115.
- 12
- 13 109. Pauli H, Gottfried M, Dullinger S, Abdaladze O, Akhalkatsi M, Alonso JLB, *et al.* Recent plant  
14 diversity changes on Europe's mountain summits. *Science* 2012, **336**(6079): 353-355.
- 15
- 16 110. Hungtington H, Fox S. The changing Arctic: indigenous perspectives. In: Symon C, Arris L, Hill B  
17 (eds). *Arctic Climate Impact Assessment*. Cambridge University Press: New York, NY, 2005, pp  
18 61-98.
- 19
- 20 111. Wiig ØA, S.; Atwood, T.; Laidre, K.; Lun, K.; *et al.* *Ursus maritimus*. The IUCN Red List of  
21 Threatened Species 2015 2015 [cited 2016] Available from:  
22 <http://dx.doi.org/10.2305/IUCN.UK.2015-4.RLTS.T22823A14871490.en>
- 23
- 24 112. Silverman J, Lazar B, Cao L, Caldeira K, Erez J. Coral reefs may start dissolving when atmospheric  
25 CO<sub>2</sub> doubles. *Geophysical Research Letters* 2009, **36**(5).
- 26
- 27 113. Frieler K, Meinshausen M, Golly A, Mengel M, Lebek K, Donner S, *et al.* Limiting global warming  
28 to 2 C is unlikely to save most coral reefs. *Nature Climate Change* 2013, **3**(2): 165-170.
- 29
- 30 114. Siqueira MFd, Peterson AT. Consequences of global climate change for geographic distributions  
31 of cerrado tree species. *Biota Neotropica* 2003, **3**(2): 1-14.
- 32
- 33 115. Midgley G, Hannah L, Millar D, Rutherford M, Powrie L. Assessing the vulnerability of species  
34 richness to anthropogenic climate change in a biodiversity hotspot. *Global Ecology and*  
35 *Biogeography* 2002, **11**(6): 445-451.
- 36
- 37 116. Preston BL. Risk-based reanalysis of the effects of climate change on US cold-water habitat.  
38 *Climatic Change* 2006, **76**(1-2): 91-119.
- 39

- 1 117. Van Oldenborgh G, Collins M, Arblaster J, Christensen J, Marotzke J, Power S, *et al.* Annex I: Atlas  
2 of global and regional climate projections. In: Stocker TF, Qin D, Plattner G-K, Tignor M, Allen D,  
3 Boschung J, *et al.* (eds). *Climate Change 2013: The Physical Science Basis. Contribution of*  
4 *Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate*  
5 *Change*. Cambridge University Press: Cambridge, United Kingdom and New York, NY, USA, 2013,  
6 pp 1311-1393.
- 7
- 8 118. Hilbert DW, Bradford M, Parker T, Westcott DA. Golden bowerbird (*Prionodura newtonia*)  
9 habitat in past, present and future climates: predicted extinction of a vertebrate in tropical  
10 highlands due to global warming. *Biological Conservation* 2004, **116**(3): 367-377.
- 11
- 12 119. Fitzpatrick MC, Gove AD, Sanders NJ, Dunn RR. Climate change, plant migration, and range  
13 collapse in a global biodiversity hotspot: the *Banksia* (Proteaceae) of Western Australia. *Global*  
14 *Change Biol* 2008, **14**(6): 1337-1352.
- 15
- 16 120. Sumaila UR, Cheung WW, Lam VW, Pauly D, Herrick S. Climate change impacts on the biophysics  
17 and economics of world fisheries. *Nature Climate Change* 2011, **1**(9): 449-456.
- 18
- 19 121. Garcia SM, de Leiva Moreno I. Global overview of marine fisheries. In: Sinclair M, Valdimarsson G  
20 (eds). *Responsible Fisheries in the Marine Ecosystem*. CABI Publishing: Oxon, UK and Cambridge,  
21 MA, USA, 2003, pp 1-24.
- 22
- 23 122. Teh LS, Teh LC, Sumaila UR. A global estimate of the number of coral reef fishers. *Plos One* 2013,  
24 **8**(6): e65397.
- 25
- 26 123. Magrin GO, Marengo JA, Boulanger MS, Buckeridge E, Castellanos E, Poveda G, *et al.* Central and  
27 South America. In: Barros VR, Field CB, Dokken DJ, Mastrandrea MD, Mach KJ, Bilir TE, *et al.*  
28 (eds). *Climate Change 2014: Impacts, Adaptation, and Vulnerability: Part B: Regional Aspects.*  
29 *Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel*  
30 *on Climate Change*. Cambridge University Press: Cambridge, United Kingdom and New York, NY,  
31 USA, 2014, pp 1499-1566.
- 32
- 33 124. Kaser G, Großhauser M, Marzeion B. Contribution potential of glaciers to water availability in  
34 different climate regimes. *Proceedings of the National Academy of Sciences* 2010, **107**(47):  
35 20223-20227.
- 36
- 37 125. Immerzeel WW, Van Beek LP, Bierkens MF. Climate change will affect the Asian water towers.  
38 *Science* 2010, **328**(5984): 1382-1385.
- 39
- 40 126. Bolch T, Kulkarni A, Kääb A, Huggel C, Paul F, Cogley J, *et al.* The state and fate of Himalayan  
41 glaciers. *Science* 2012, **336**(6079): 310-314.

- 1  
2 127. Immerzeel W, Pellicciotti F, Bierkens M. Rising river flows throughout the twenty-first century in  
3 two Himalayan glacierized watersheds. *Nature Geoscience* 2013, **6**(9): 742-745.
- 4  
5 128. Honda Y, Kondo M, McGregor G, Kim H, Guo Y-L, Hijioka Y, *et al.* Heat-related mortality risk  
6 model for climate change impact projection. *Environmental Health and Preventive Medicine*  
7 2014, **19**(1): 56-63.
- 8  
9 129. Sillmann J, Kharin V, Zwiers F, Zhang X, Bronaugh D. Climate extremes indices in the CMIP5  
10 multimodel ensemble: Part 2. Future climate projections. *Journal of Geophysical Research:*  
11 *Atmospheres* 2013, **118**(6): 2473-2493.
- 12  
13 130. Maranz S. Tree mortality in the African Sahel indicates an anthropogenic ecosystem displaced by  
14 climate change. *Journal of Biogeography* 2009, **36**(6): 1181-1193.
- 15  
16 131. Guis H, Caminade C, Calvete C, Morse AP, Tran A, Baylis M. Modelling the effects of past and  
17 future climate on the risk of bluetongue emergence in Europe. *Journal of the Royal Society*  
18 *Interface* 2012, **9**(67): 339-350.
- 19  
20 132. Intergovernmental Panel on Climate Change. Summary for Policymakers. In: Solomon S, Qin D,  
21 Manning M, Chen Z, Marqui M, Averyt KB, *et al.* (eds). *Climate Change 2007: The Physical*  
22 *Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the*  
23 *Intergovernmental Panel on Climate Change*. Cambridge University Press: Cambridge, United  
24 Kingdom and New York, NY, USA, 2007.
- 25  
26 133. Ramirez-Villegas J, Salazar M, Jarvis A, Navarro-Racines CE. A way forward on adaptation to  
27 climate change in Colombian agriculture: perspectives towards 2050. *Climatic Change* 2012,  
28 **115**(3-4): 611-628.
- 29  
30 134. Lobell DB, Burke MB, Tebaldi C, Mastrandrea MD, Falcon WP, Naylor RL. Prioritizing climate  
31 change adaptation needs for food security in 2030. *Science* 2008, **319**(5863): 607-610.
- 32  
33 135. Gerten D, Lucht W, Ostberg S, Heinke J, Kowarsch M, Kreft H, *et al.* Asynchronous exposure to  
34 global warming: freshwater resources and terrestrial ecosystems. *Environmental Research*  
35 *Letters* 2013, **8**(3): 034032.
- 36  
37 136. Portmann FT, Döll P, Eisner S, Flörke M. Impact of climate change on renewable groundwater  
38 resources: assessing the benefits of avoided greenhouse gas emissions using selected CMIP5  
39 climate projections. *Environmental Research Letters* 2013, **8**(2): 024023.

40

- 1 137. Ward PJ, Strzepek KM, Pauw WP, Brander LM, Hughes GA, Aerts JC. Partial costs of global  
2 climate change adaptation for the supply of raw industrial and municipal water: a methodology  
3 and application. *Environmental Research Letters* 2010, **5**(4): 044011.
- 4
- 5 138. Hirabayashi Y, Mahendran R, Koirala S, Konoshima L, Yamazaki D, Watanabe S, *et al.* Global  
6 flood risk under climate change. *Nature Climate Change* 2013, **3**(9): 816-821.
- 7
- 8 139. Dankers R, Arnell NW, Clark DB, Falloon PD, Fekete BM, Gosling SN, *et al.* First look at changes in  
9 flood hazard in the Inter-Sectoral Impact Model Intercomparison Project ensemble. *Proceedings*  
10 *of the National Academy of Sciences* 2014, **111**(9): 3257-3261.
- 11
- 12 140. Fung F, Lopez A, New M. Water availability in +2 C and +4 C worlds. *Philosophical Transactions of*  
13 *the Royal Society of London A: Mathematical, Physical and Engineering Sciences* 2011,  
14 **369**(1934): 99-116.
- 15
- 16 141. Kjellstrom T, Lemke B, Otto M. Mapping occupational heat exposure and effects in South-East  
17 Asia: ongoing time trends 1980–2011 and future estimates to 2050. *Industrial Health* 2013,  
18 **51**(1): 56-67.
- 19
- 20 142. Kjellstrom T, Holmer I, Lemke B. Workplace heat stress, health and productivity - an increasing  
21 challenge for low and middle-income countries during climate change. *Global Health Action*  
22 2009, **2**.
- 23
- 24 143. Sherwood SC, Huber M. An adaptability limit to climate change due to heat stress. *Proceedings*  
25 *of the National Academy of Sciences* 2010, **107**(21): 9552-9555.
- 26
- 27 144. Hijioka Y, Lin E, Pereira JJ, Corlett RT, Cui X, Insarov GE, *et al.* Asia. In: Barros VR, Field CB,  
28 Dokken DJ, Mastrandrea MD, Mach KJ, Bilir TE, *et al.* (eds). *Climate Change 2014: Impacts,*  
29 *Adaptation, and Vulnerability: part B: Regional Aspects. Contribution of Working Group II to the*  
30 *Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge  
31 University Press: Cambridge, United Kingdom and New York, NY, USA, 2014, pp 1327-1370.
- 32
- 33 145. Kjellstrom T, Kovats RS, Lloyd SJ, Holt T, Tol RS. The direct impact of climate change on regional  
34 labor productivity. *Archives of Environmental & Occupational Health* 2009, **64**(4): 217-227.
- 35
- 36 146. Hsiang SM, Burke M, Miguel E. Quantifying the influence of climate on human conflict. *Science*  
37 2013, **341**(6151): 1235367.
- 38
- 39 147. Blattman C, Miguel E. Civil war. *Journal of Economic Literature* 2010, **48**(1): 3-57.
- 40



- 1 148. Myers SS, Zanobetti A, Kloog I, Huybers P, Leakey AD, Bloom A, *et al.* Rising CO2 threatens  
2 human nutrition. *Nature* 2014, **510**(7503): 139-142.
- 3
- 4 149. Li Y, Ye W, Wang M, Yan X. Climate change and drought: a risk assessment of crop-yield impacts.  
5 *Clim Res* 2009, **39**(1): 31.
- 6
- 7 150. Tebaldi C, Lobell D. Estimated impacts of emission reductions on wheat and maize crops.  
8 *Climatic Change* 2015: 1-13.
- 9
- 10 151. Deryng D, Conway D, Ramankutty N, Price J, Warren R. Global crop yield response to extreme  
11 heat stress under multiple climate change futures. *Environmental Research Letters* 2014, **9**(3):  
12 034011.
- 13
- 14 152. Barriopedro D, Fischer EM, Luterbacher J, Trigo RM, García-Herrera R. The hot summer of 2010:  
15 redrawing the temperature record map of Europe. *Science* 2011, **332**(6026): 220-224.
- 16
- 17 153. Easterling W, Aggarwal P, Batima P, Brander K, Erda L, Howden, Mark, *et al.* Food, fibre and  
18 forest products. In: Parry ML, Canziani OF, Palutikof JP, van der Linden PJ, Hanson CE (eds).  
19 *Change 2007: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the*  
20 *Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge  
21 University Press: Cambridge, UK, 2007, pp 273-313.
- 22
- 23 154. Poloczanska E, Hoegh-Guldberg O, Cheung W, Pörtner H-O, Burrows MT. Crosss Chapter Box on  
24 Observed global responses of marine biogeography, abundance, and phenology to climate  
25 change. In: Field CB, Barros VR, Dokken DJ, Mach KJ, Mastrandrea MD, Bilir TE, *et al.* (eds).  
26 *Climate Change 2014: Impacts, Adaptation, And Vulnerability: Part A: Global And Sectoral*  
27 *Aspects. Contribution Of Working Group II To The Fifth Assessment Report Of The*  
28 *Intergovernmental Panel On Climate Change*. Cambridge University Press: Cambridge, United  
29 Kingdom and New York, NY, USA, 2014, pp 123-127.
- 30
- 31 155. Adamík P, Král M. Climate - and resource - driven long - term changes in dormice populations  
32 negatively affect hole - nesting songbirds. *Journal of Zoology* 2008, **275**(3): 209-215.
- 33
- 34 156. Adrian R, O'Reilly CM, Zagarese H, Baines SB, Hessen DO, Keller W, *et al.* Lakes as sentinels of  
35 climate change. *Limnology and Oceanography* 2009, **54**(6): 2283.
- 36
- 37 157. Cook BI, Wolkovich EM, Davies TJ, Ault TR, Betancourt JL, Allen JM, *et al.* Sensitivity of spring  
38 phenology to warming across temporal and spatial climate gradients in two independent  
39 databases. *Ecosystems* 2012, **15**(8): 1283-1294.
- 40

- 1 158. Devictor V, van Swaay C, Brereton T, Chamberlain D, Heliölä J, Herrando S, *et al.* Differences in  
2 the climatic debts of birds and butterflies at a continental scale. *Nature Climate Change* 2012,  
3 **2**(2): 121-124.
- 4
- 5 159. Phillimore AB, Hadfield JD, Jones OR, Smithers RJ. Differences in spawning date between  
6 populations of common frog reveal local adaptation. *Proceedings of the National Academy of*  
7 *Sciences* 2010, **107**(18): 8292-8297.
- 8
- 9 160. Gehrig - Fasel J, Guisan A, Zimmermann NE. Tree line shifts in the Swiss Alps: Climate change or  
10 land abandonment? *Journal of Vegetation Science* 2007, **18**(4): 571-582.
- 11
- 12 161. Kusano T, Inoue M. Long-term trends toward earlier breeding of Japanese amphibians. *Journal*  
13 *of Herpetology* 2008, **42**(4): 608-614.
- 14
- 15 162. Lenoir J, Gégout J-C, Marquet P, De Ruffray P, Brisse H. A significant upward shift in plant species  
16 optimum elevation during the 20th century. *Science* 2008, **320**(5884): 1768-1771.
- 17
- 18 163. Garamszegi LZ. Climate change increases the risk of malaria in birds. *Global Change Biol* 2011,  
19 **17**(5): 1751-1759.
- 20
- 21 164. Peterson AT, Navarro-Sigüenza AG, Martínez-Meyer E, Cuervo-Robayo AP, Berlanga H, Soberón  
22 J. Twentieth century turnover of Mexican endemic avifaunas: Landscape change versus climate  
23 drivers. *Science Advances* 2015, **1**(4): e1400071.
- 24
- 25 165. Brook BW, Sodhi NS, Bradshaw CJ. Synergies among extinction drivers under global change.  
26 *Trends in Ecology & Evolution* 2008, **23**(8): 453-460.
- 27
- 28 166. Burke M, Hsiang SM, Miguel E. Global non-linear effect of temperature on economic production.  
29 *Nature* 2015.
- 30
- 31 167. Dell M, Jones BF, Olken BA. Temperature shocks and economic growth: Evidence from the last  
32 half century. *American Economic Journal: Macroeconomics* 2012: 66-95.
- 33
- 34 168. Platts PJ, Garcia RA, Hof C, Foden W, Hansen LA, Rahbek C, *et al.* Conservation implications of  
35 omitting narrow - ranging taxa from species distribution models, now and in the future.  
36 *Diversity and Distributions* 2014, **20**(11): 1307-1320.
- 37
- 38 169. Sekercioglu CHS, S. H.; Fay, J. P. Climate Change, Elevational Range Shifts, and Bird Extinctions.  
39 *Conservation Biology* 2008, **22**: 140-150.

- 1  
2 170. Cheung WW, Lam VW, Sarmiento JL, Kearney K, Watson R, Pauly D. Projecting global marine  
3 biodiversity impacts under climate change scenarios. *Fish and Fisheries* 2009, **10**(3): 235-251.
- 4  
5 171. Cheung WW, Sarmiento JL, Dunne J, Frölicher TL, Lam VW, Palomares MD, *et al.* Shrinking of  
6 fishes exacerbates impacts of global ocean changes on marine ecosystems. *Nature Climate*  
7 *Change* 2013, **3**(3): 254-258.
- 8  
9 172. Rehdanz K, Maddison D. Climate and happiness. *Ecological Economics* 2005, **52**(1): 111-125.
- 10  
11 173. Tol RSJ. Estimates of the damage costs of climate change. Part 1: Benchmark estimates.  
12 *Environmental and Resource Economics* 2002, **21**(1): 47-73.
- 13  
14 174. Bosello F, Eboli F, Pierfederici R. Assessing the economic impacts of climate change. *Review of*  
15 *Environment, Energy and Economics* 2012: 1-9.
- 16  
17 175. Fankhauser S. *Valuing climate change: the economics of the greenhouse*. Routledge, 2013.
- 18  
19 176. Hope C. The marginal impact of CO2 from PAGE2002: An integrated assessment model  
20 incorporating the IPCC's five reasons for concern. *Integrated Assessment* 2006, **6**(1).
- 21  
22 177. Nordhaus WD. *A question of balance: Weighing the options on global warming policies*. Yale  
23 University Press, 2014.
- 24  
25 178. Roson R, Van der Mensbrugghe D. Climate change and economic growth: impacts and  
26 interactions. *International Journal of Sustainable Economy* 2012, **4**(3): 270-285.
- 27  
28 179. Tol RSJ. The damage costs of climate change toward more comprehensive calculations.  
29 *Environmental and Resource Economics* 1995, **5**(4): 353-374.
- 30  
31 180. Cardinale BJ, Duffy JE, Gonzalez A, Hooper DU, Perrings C, Venail P, *et al.* Biodiversity loss and its  
32 impact on humanity. *Nature* 2012, **486**(7401): 59-67.
- 33  
34 181. Midgley GF. Biodiversity and ecosystem function. *Science* 2012, **335**(6065): 174-175.
- 35  
36 182. Warszawski L, Friend A, Ostberg S, Frieler K, Lucht W, Schaphoff S, *et al.* A multi-model analysis  
37 of risk of ecosystem shifts under climate change. *Environmental Research Letters* 2013, **8**(4):  
38 044018.

- 1  
2 183. Maddison D, Rehdanz K. The impact of climate on life satisfaction. *Ecological Economics* 2011,  
3 **70**(12): 2437-2445.
- 4  
5 184. Arent D, Tol R, Faust E, Hella J, Kumar S, Strzepek KM, *et al.* Key economic sectors and services.  
6 In: Field CB, Barros VR, Dokken DJ, Mach KJ, Mastrandrea MD, Bilir TE, *et al.* (eds). *Climate*  
7 *Change 2014: Impacts, Adaptation, And Vulnerability: Part A: Global And Sectoral Aspects.*  
8 *Contribution Of Working Group II To The Fifth Assessment Report Of The Intergovernmental*  
9 *Panel On Climate Change.* Cambridge University Press: Cambridge, United Kingdom and New  
10 York, NY, USA, 2014, pp 659-708.
- 11  
12 185. Anthoff D, Tol RS. The climate framework for uncertainty, negotiation and distribution (FUND),  
13 Technical description, Version 3.5. 2010.
- 14  
15 186. Hope C. Critical issues for the calculation of the social cost of CO2: why the estimates from  
16 PAGE09 are higher than those from PAGE2002. *Climatic Change* 2013, **117**(3): 531-543.
- 17  
18 187. Nordhaus WD. An optimal transition path for controlling greenhouse gases. *Science* 1992,  
19 **258**(5086): 1315-1319.
- 20  
21 188. Tol RSJ. On the optimal control of carbon dioxide emissions: an application of FUND.  
22 *Environmental Modeling & Assessment* 1997, **2**(3): 151-163.
- 23  
24 189. Kopp RE, Hsiang SM, Oppenheimer M. Empirically calibrating damage functions and considering  
25 stochasticity when integrated assessment models are used as decision tools. *Impacts World*  
26 *2013, International Conference on Climate Change Effects.* Potsdam, Germany; 2013.
- 27  
28 190. Kopp RE, Mignone BK. The US government's social cost of carbon estimates after their first two  
29 years: Pathways for improvement. *Economics: The Open-Access, Open-Assessment E-Journal*  
30 2012, **6**.
- 31  
32 191. Pindyck RS. Climate change policy: What do the models tell us? *Journal of Economic Literature*  
33 2013, **51**(3): 860-872.
- 34  
35 192. Lobell DB, Field CB. Global scale climate–crop yield relationships and the impacts of recent  
36 warming. *Environmental Research Letters* 2007, **2**(1): 014002.
- 37  
38 193. Mastrandrea MD. Representation of Climate Impacts in Integrated Assessment Models. In:  
39 Gullege J, Richardson LJ, Adkins L, Seidel S, editors. *Assessing the Benefits of Avoided Climate*

- 1 Change: Cost Benefit Analysis and Beyond; 2010; Arlington, VA: Pew Center on Global Climate  
2 Change; 2010. p. 85-99.
- 3
- 4 194. Yohe GW, Tirpak D. A research agenda to improve economic estimates of the benefits of climate  
5 change policies. *Integrated Assessment Journal* 2008, **8**(1): 1-17.
- 6
- 7 195. Chivian E, Bernstein A. *Sustaining life: how human health depends on biodiversity*. Oxford  
8 University Press, 2008.
- 9
- 10 196. Kopp RE, Golub A, Keohane NO, Onda C. The influence of the specification of climate change  
11 damages on the social cost of carbon. *Economics: The Open-Access, Open-Assessment E-Journal*  
12 2012, **6**.
- 13
- 14 197. Sterner T, Persson UM. An even sterner review: Introducing relative prices into the discounting  
15 debate. *Review of Environmental Economics and Policy* 2008, **2**(1): 61-76.
- 16
- 17 198. Weitzman ML. What Is The "Damages Function" For Global Warming—And What Difference  
18 Might It Make? *Climate Change Economics* 2010, **1**(01): 57-69.
- 19
- 20 199. Lontzek TS, Cai Y, Judd KL, Lenton TM. Stochastic integrated assessment of climate tipping  
21 points indicates the need for strict climate policy. *Nature Climate Change* 2015, **5**(5): 441-444.
- 22
- 23 200. Arnell NW, Lowe J, Brown S, Gosling S, Gottschalk P, Hinkel J, *et al*. A global assessment of the  
24 effects of climate policy on the impacts of climate change. *Nature Climate Change* 2013, **3**(5):  
25 512-519.
- 26
- 27 201. Warren R, Lowe JA, Arnell N, Hope C, Berry P, Brown S, *et al*. The AVOID programme's new  
28 simulations of the global benefits of stringent climate change mitigation. *Climatic Change* 2013,  
29 **120**(1-2): 55-70.
- 30
- 31 202. Houser T, Hsiang S, Kopp R, Larsen K. *Economic risks of climate change: an American prospectus*.  
32 Columbia University Press, 2015.
- 33
- 34 203. Waldhoff S, Martinich J, Sarofim M, DeAngelo B, McFarland J, Jantarasami L, *et al*. Overview of  
35 the special issue: a multi-model framework to achieve consistent evaluation of climate change  
36 impacts in the United States. *Climatic Change* 2014: 1-20.
- 37
- 38 204. Robinson A, Calov R, Ganopolski A. Multistability and critical thresholds of the Greenland ice  
39 sheet. *Nature Climate Change* 2012, **2**(6): 429-432.

- 1  
2 205. Dutton A, Lambeck K. Ice volume and sea level during the last interglacial. *Science* 2012,  
3 **337**(6091): 216-219.
- 4  
5 206. Kopp RE, Simons FJ, Mitrovica JX, Maloof AC, Oppenheimer M. Probabilistic assessment of sea  
6 level during the last interglacial stage. *Nature* 2009, **462**(7275): 863-867.
- 7  
8 207. Dutton A, Webster JM, Zwartz D, Lambeck K, Wohlfarth B. Tropical tales of polar ice: evidence of  
9 Last Interglacial polar ice sheet retreat recorded by fossil reefs of the granitic Seychelles islands.  
10 *Quaternary Science Reviews* 2015, **107**: 182-196.
- 11  
12 208. Joughin I, Smith BE, Medley B. Marine ice sheet collapse potentially under way for the Thwaites  
13 Glacier Basin, West Antarctica. *Science* 2014, **344**(6185): 735-738.
- 14  
15 209. DeConto RM, Pollard D. Contribution of Antarctica to past and future sea-level rise. *Nature*  
16 2016, **531**(7596): 591-597.
- 17  
18 210. Alley RB, Marotzke J, Nordhaus WD, Overpeck JT, Peteet DM, Pielke RA, *et al.* Abrupt climate  
19 change. *Science* 2003, **299**(5615): 2005-2010.
- 20  
21 211. Steffensen JP, Andersen KK, Bigler M, Clausen HB, Dahl-Jensen D, Fischer H, *et al.* High-  
22 resolution Greenland ice core data show abrupt climate change happens in few years. *Science*  
23 2008, **321**(5889): 680-684.
- 24  
25 212. Hallegatte S, Green C, Nicholls RJ, Corfee-Morlot J. Future flood losses in major coastal cities.  
26 *Nature Climate Change* 2013, **3**(9): 802-806.
- 27  
28 213. Anthoff D, Nicholls RJ, Tol RS. The economic impact of substantial sea-level rise. *Mitigation and*  
29 *Adaptation Strategies for Global Change* 2010, **15**(4): 321-335.
- 30  
31 214. Hanasaki N, Fujimori S, Yamamoto T, Yoshikawa S, Masaki Y, Hijioka Y, *et al.* A global water  
32 scarcity assessment under Shared Socio-economic Pathways—Part 2: Water availability and  
33 scarcity. *Hydrol Earth Syst Sci* 2013, **17**(7): 2393-2413.
- 34  
35 215. Ishida H, Kobayashi S, Kanae S, Hasegawa T, Fujimori S, Shin Y, *et al.* Global-scale projection and  
36 its sensitivity analysis of the health burden attributable to childhood undernutrition under the  
37 latest scenario framework for climate change research. *Environmental Research Letters* 2014,  
38 **9**(6): 064014.
- 39

- 1 216. Johns T, Gregory J, Ingram W, Johnson C, Jones A, Lowe J, *et al.* Anthropogenic climate change
- 2 for 1860 to 2100 simulated with the HadCM3 model under updated emissions scenarios.
- 3 *Climate Dynamics* 2003, **20**(6): 583-612.

4

5