

Late Quaternary uplift and earthquake potential of the San Joaquin Hills, southern Los Angeles basin, California

Lisa B. Grant Department of Environmental Analysis, University of California, Irvine, California 92697-7070, USA
Karl J. Mueller Department of Geological Sciences, University of Colorado, Boulder, Colorado 80309, USA
Eldon M. Gath Department of Environmental Analysis, University of California, Irvine, California 92697-7070, USA
Hai Cheng
R. Lawrence Edwards } Department of Geology and Geophysics, University of Minnesota, Minneapolis, Minnesota 55455, USA
Rosalind Munro Leighton and Associates, 17781 Cowan, Irvine, California 92614, USA
George L. Kennedy Department of Geological Sciences, San Diego State University, San Diego, California 92182-1020, USA

ABSTRACT

Analysis of emergent marine terraces in the San Joaquin Hills, California, and ^{230}Th dating of solitary corals from the lowest terraces reveal that the San Joaquin Hills have risen at a rate of 0.21–0.27 m/k.y. during the past 122 k.y. Movement on a blind thrust fault in the southern Los Angeles basin has uplifted the San Joaquin Hills and has the potential to generate an M_w 7.3 earthquake within this densely populated area. Our structural modeling suggests that the fault dips to the southwest and slips at ~0.42–0.79 m/k.y., yielding an estimated minimum average recurrence interval of ~1650–3100 yr for moderate-sized earthquakes. Recognition of this blind thrust extends the known area of active blind thrusts and fault-related folding southward from Los Angeles into coastal Orange County.

INTRODUCTION AND GEOLOGIC SETTING

The Whittier Narrows and Northridge blind thrust faults in the Los Angeles, California, area were recognized only after they generated damaging, moderate-sized earthquakes (M_w 6.0 and 6.7) in 1987 and 1994 (Bullard and Lettis, 1993; Scientists of USGS and the Southern California Earthquake Center, 1994). Blind thrust faults are often associated with active folds (Stein and Yeats, 1989; Ward and Valensise, 1994; Lettis et al., 1997). Indications of late Quaternary folding are present in the San Joaquin Hills at the southern margin of the Los Angeles basin (Fig. 1). We present stratigraphic and geomorphic analyses in combination with ^{230}Th coral dating to calculate rates of uplift and evaluate the potential for blind thrust earthquakes beneath the San Joaquin Hills.

The Los Angeles basin is a northwest-trending structural trough bounded by active reverse and left-lateral faults of the Transverse Ranges to the north and by active right-lateral faults of the Peninsular Ranges to the northeast and southwest (Wright, 1991; see inset map in Fig. 1). The San Joaquin Hills are the topographic expression of a northwest-trending anticline between San Juan Capistrano and Huntington Mesa (Vedder, 1975; Lajoie et al., 1991) (Fig. 1). The Newport-Inglewood fault zone and its offshore extension (Wright, 1991; Fig. 1) bound the San Joaquin Hills on the southwest. The Newport-Inglewood fault zone generated the 1933 M_w 6.4 Long Beach earthquake near the mouth of the Santa Ana River (Hauksson and Gross, 1991).

Uplift of the San Joaquin Hills began in the early Pleistocene (Barrie et al., 1992). A suite of emergent marine terraces is present along the coastal San Joaquin Hills (Vedder et al., 1957). We extended previous mapping of terrace deposits (Barrie et al., 1992) and measured the eleva-

tions of shorelines and terrace platforms using geotechnical excavations, borings, natural exposures, and topography (Fig. 2). We correlated coastal terraces with inland terraces by comparing elevation, stratigraphic relationships, degree of dissection, soil profiles, and faunal assemblages (Kanakoff and Emerson, 1959; Peska, 1984).

MEASUREMENT OF UPLIFT RATE

Fossil localities in the northern San Joaquin Hills have been used for zoogeographic correlation and aminostratigraphic dating of upper Quaternary marine deposits in the Los Angeles basin, but there are significant uncertainties in amino acid dating due to temperature sensitivity (Wehmiller et al., 1977; Kennedy et al., 1982; Lajoie et al., 1991). The ^{230}Th dating of solitary corals is a more reliable method of dating marine platforms. It can calibrate amino acid dates and ages of faunal assemblages. Because corals are uncommon in southern California, ^{230}Th ages from the Los Angeles basin have not previously been reported.

We calculated the average late Quaternary uplift rate of the San Joaquin Hills using coral ages from three localities and measurements of paleoshoreline (shoreline angle) elevations (Fig. 2). Coral U and Th concentrations, isotopic ratios, and ^{230}Th ages (Table 1) were determined by thermal ionization mass spectrometry using the methods of Edwards et al. (1987) and Cheng et al. (1999).

Three subsamples of a single *Paracyathus pedroensis* skeleton (sample FP-28) from terrace 2, on the east side of upper Newport Bay, yielded similar ages and initial $\delta^{234}\text{U}$ values only slightly higher than the modern marine value of $146\text{‰} \pm 2\text{‰}$ (Cheng et al., 1999). Near-primary uranium isotopic composition and agreement among the radiometric ages imply minimal diagenetic alter-

ation. The ages (Table 1) correlate well with the last interglacial period (Edwards et al., 1987), marine oxygen isotope substage 5e.

Three solitary corals (*Balanophyllia elegans* samples 12654-0, 12654-1, and 12654-2; Table 1) from the second terrace along the coast of the San Joaquin Hills (Fig. 2) had initial $\delta^{234}\text{U}$ values significantly higher than the modern marine value, suggesting diagenetic alteration. The range of $^{230}\text{Th}/^{238}\text{U}$ values for a set of California solitary corals correlated to the last interglacial period (Stein et al., 1991) is a guide to the magnitude of diagenetic shifts in $^{230}\text{Th}/^{238}\text{U}$ values (and ^{230}Th ages). We assume that the range in $^{230}\text{Th}/^{238}\text{U}$ activity ratios (0.7516–0.8516) for the Stein et al. (1991) samples is due solely to diagenesis, that the same percentage spread affected our samples at the 12654 locality, and we then center this spread on the mean $^{230}\text{Th}/^{238}\text{U}$ ratio of the 12654 samples. The lowest $^{230}\text{Th}/^{238}\text{U}$ ratio corresponds to an age of about 120 ka, and the highest ratio corresponds to an age of about 240 ka.

A specimen of *Paracyathus pedroensis* from the first terrace (locality GK-90-14, Fig. 2) yielded an age of 106 ka (Table 1). The sample has an initial $\delta^{234}\text{U}$ value close to the modern marine value, suggesting that the radiometric age is reliable. Barrie et al. (1992) correlated the lowest San Joaquin Hills terrace with substage 5a on the basis of the lack of extralimital southern (i.e., warm water) fossils, the extent of amino acid racemization, and elevation of the shoreline angle. Elevation of the substage 5a highstand was well constrained as -1 m for the southern California coast by Muhs et al. (1994), but they reported -20 m to -1 m elevation of the substage 5c highstand. On the basis of terrace and eustatic sea-level correlations (Fig. 3), we conclude that the substage 5c coral was reworked onto the substage 5a terrace, or the substage 5c and 5a sea stands occupied the same platform.

The elevation difference between the terrace 2 shoreline angle (32–35 m) and substage 5e sea level (6 m; Lajoie et al., 1991) divided by the average age of sample FP-28 (122 ka) yields an uplift rate of 0.21–0.24 m/k.y. Analogous calculations for the 12654 locality yield uplift rates of 0.24–0.25 m/k.y. for a substage 5e correlation. Using 0.24 m/k.y. as our best estimate of uplift rate during the Quaternary, the age of the highest shoreline (305 m) is ca. 1.3 Ma, and several inter-




Figure 1. Regional map of San Joaquin Hills (SJH) showing approximate location of fold crest of San Joaquin Hills anticline. Other features: AC—Aliso Canyon, DP—Dana Point, HM—Huntington Mesa, LC—Laguna Canyon, NIF—Newport-Inglewood fault, NM—Newport Mesa, SDC—San Diego Creek, and SJC—San Juan Capistrano. Epicenters of 1933, 1987, and 1994 earthquakes are marked by stars. Modified from Vedder et al. (1957). Inset map shows greater Los Angeles region.

mediate terraces were probably occupied by multiple highstands (Fig. 3).

REGENCY OF UPLIFT

The location and thickness of Holocene sediments in the San Joaquin Hills suggest that tectonic uplift continued during the middle to late Holocene. Isopach and structure contour maps of the Holocene Talbert aquifer beneath the Santa Ana River (Sprotte et al., 1980) suggest that the aquifer has been deformed. The distribution of fluvial and estuarine sediments southeast of the Santa Ana River also suggests that Holocene uplift occurred. Newport Bay was incised at least 36 m below present sea level during the last glacial maximum, but rapid sea-level rise at the close of the Pleistocene inundated coastal drainages and induced sedimentary infilling with Holocene sediments (Stevenson, 1954). Stevenson concluded that an elevated bench of former marsh deposits in Newport Bay was created during the late Holocene by emergence. Stevenson speculated that the emergence was due to tectonic uplift and, based on elevation profiles, the uplift reflected anticlinal folding along a northwest-trending fold axis.

DISCUSSION

The late Quaternary uplift rate, anticlinal structure, and indications of Holocene uplift imply that the San Joaquin Hills are the surface expression of an active contractile fold (see Fig. 1), formed above a potentially seismogenic thrust fault (Stein and Yeats, 1989; Lettis et al., 1997; Shaw and Shearer, 1999). Geomorphic analysis of the San Joaquin Hills provides some constraints on the geometry of the proposed blind fault. A fault-bend fold model with movement on a northeast-vergent thrust fault best explains the elevation of marine terraces on the northeast limb of the San Joaquin Hills anticline, as shown schematically in Figure 4. Anticlinal structure of the San Joaquin Hills and northeast vergence of the underlying fault are supported by structural data of Vedder (1975) and our terrace mapping.

The maximum-magnitude earthquake that could occur on the San Joaquin Hills thrust can be estimated from empirical relationships between magnitude and subsurface fault-rupture length, as defined by the areal distribution of Quaternary uplift. This method does not require assumptions about fault geometry. The area of uplift extends at least ~38 km, from northwestern Huntington Mesa southeast to Dana Point, and therefore the




Figure 2. Map of marine-terrace platforms, location of measured shoreline elevations (open circles), and fossil localities (stars). See Figure 1 for location.

Table 1. URANIUM AND THORIUM ISOTOPIC COMPOSITIONS AND CONCENTRATIONS OF SOLITARY CORALS

Locality*/ sample number	Species	Sample/shoreline elevation (m)	²³⁸ U (ppm)	²³² Th (ppb)	²³⁰ Th/ ²³⁸ U (activity ratio)	²³⁰ Th age † (ka)	δ ²³⁴ U ‡ initial	Stage or substage
SDSU 3812/FP-28a	<i>Paracyathus pedroensis</i>	24/33-34	3.471 (7)	37.9 (3)	0.7616 (26)	122.6 (1.0)	157 (4)	5e
SDSU 3812/FP-28b	<i>Paracyathus pedroensis</i>	24/33-34	3.678 (4)	61.8 (6)	0.7717 (68)	124.1 (2.1)	165 (3)	5e
SDSU 3812/FP-28c	<i>Paracyathus pedroensis</i>	24/33-34	3.327 (3)	74.5 (5)	0.7563 (48)	120.4 (1.6)	159 (3)	5e
GK-90-14	<i>Paracyathus pedroensis</i>	16 – 17/21	4.681 (6)	129.6 (1.1)	0.7131 (24)	105.6 (0.7)	173 (3)	5c
LACMNH 12654/12654-0	<i>Balanophyllia elegans</i>	29-30.5/35-36	3.702 (7)	283 (4)	0.9448 (47)	172.8 (3.1)	247 (6)	5e or 7
LACMNH 12654/12654-1	<i>Balanophyllia elegans</i>	29-30.5/35-36	3.691 (4)	419.6 (2.3)	0.9430 (36)	167.7 (2.2)	258 (4)	5e or 7
LACMNH 12654/12654-2	<i>Balanophyllia elegans</i>	29-30.5/35-36	3.618 (22)	179.6 (1.1)	0.8509 (57)	137.6 (5.0)	230 (26)	5e or 7


Note: The first three analyses are different fragments of the same individual coral skeleton. All other analyses are on individual corals. Decay constants for ²³⁴U and ²³⁰Th are new values from Cheng et al. (1999): λ₂₃₀ = 9.1577 × 10⁻⁶/y and λ₂₃₄ = 2.8263 × 10⁻⁶/y. The decay constant for ²³⁸U is referenced in Cheng et al. (1999): λ₂₃₈ = 1.551 × 10⁻¹⁰/y. 2σ errors are in parentheses and represent errors in the last significant figures.

*Locality data available from E. C. Allison Center, Department of Geological Sciences, San Diego State University (SDSU), San Diego, California; Invertebrate Paleontology Section, Los Angeles County Museum of Natural History (LACMNH), Los Angeles, California.

† Calculated from isotope ratios using the standard age equation (e.g., Edwards et al., 1987), with correction for initial ²³⁰Th assuming an initial ²³⁰Th/²³²Th value of (4.4 ± 4.4) × 10⁻⁶, the bulk Earth thorium isotopic value with an arbitrarily assumed error of ±100%. For all samples, the correction is small and error introduced by the correction is small compared to analytical error.

‡ δ²³⁴U = [(²³⁴U/²³⁸U) - 1] × 1000; brackets indicate an activity ratio. Initial δ²³⁴U value is calculated from the measured value by using the ²³⁰Th age.

Figure 3. Correlation of shoreline-angle elevations of coastal marine terraces with oxygen isotope record of eustatic sea-level highstands using methodology of Merritts and Bull (1989). Ages of terraces are estimated by their correlation with sea-level highstands, assuming constant 0.24 m/k.y. rate of uplift (slope of line) based on substage 5e terrace. Sea-level curve compiled from Chappell and Shackleton (1986) with modifications from Muhs et al. (1994) and Gallup et al. (1994). Oxygen isotope stages are numbered. Maximum and minimum reported sea-level elevations for oxygen isotope substages 5a and 5c are both shown. Width of correlation line corresponds to range in mapped shoreline angle elevations.



fault could generate a M_w 6.8 earthquake, using the length regression of Wells and Coppersmith (1994). Alternatively, the magnitude of a maximum credible earthquake is estimated by assuming that the San Joaquin Hills thrust extends to the base of the seismogenic crust at 17 km, dips between 20° and 30°, and extends upward to within 2 km of the surface. In this interpretation the San Joaquin Hills thrust is a backthrust that soles into the Oceanside detachment (Bohannon and Geist, 1998) as part of a wedge-thrust structure. The dip of the San Joaquin Hills fault is modeled after the offshore segment of the San Joaquin Hills anticline, south of Dana Point (e.g., Fisher and Mills, 1991). In this model, rupture of the entire fault area could generate a M_w 7.2–7.3 earthquake, using the regression of Dolan et al. (1995). The late




Figure 4. Block diagram of marine-terraces on northern San Joaquin Hills (viewed toward southwest). Heavy dashed lines indicate active axial surfaces where terrace platforms are folded. Large arrows denote sense of folding. Direction of movement (out of or into page, respectively) is shown by small arrows and plus or minus sign. NIFZ is Newport-Inglewood fault zone.

Pleistocene slip rate of 0.42–0.79 m/k.y. is estimated from inferred fault dip (20°–30°) and the measured uplift rate of 0.21–0.27 m/k.y.

The recurrence time for earthquakes is estimated from the equation $T = D/V$, where T is the average recurrence time, V is the slip rate, and D is the average displacement per earthquake. We use $D = 1.3$ m of average uplift (Grant and Ballenger, 1999) for the most recent event and the slip rates (0.42–0.79 m/k.y.) to calculate average recurrence times of 1650–3100 yr for moderate-magnitude earthquakes.

We prefer to interpret movement of the San Joaquin Hills blind thrust to be the product of partitioned strike slip and compressive shortening across the southern Newport-Inglewood fault zone, similar to partitioned slip reported by Hauksson (1990) along the western Los Angeles basin. Grant et al. (1997) reported multiple surface ruptures on the southern Newport-Inglewood fault zone during the Holocene, but other than observations of an elevated marsh bench in Newport Bay and sparse microseismicity, we do not have direct evidence for Holocene activity of the San Joaquin Hills thrust.

However, on July 28, 1769, Gaspar de Portola, the first Spanish explorer into southern California, camped on the eastern bank of the Santa Ana River, ~15–20 km north of the San Joaquin Hills. A violent earthquake occurred at noon, followed 10 minutes later by a severe aftershock and two other strong aftershocks later that day. Aftershocks continued for the next five days (Smith and Teggart, 1909). Although there are many active faults in southern California that could have generated the 1769 earthquake, Portola's location makes a blind San Joaquin Hills source worthy of serious consideration.

ACKNOWLEDGMENTS

This research was supported by the Southern California Earthquake Center (SCEC). SCEC is funded by the National Science Foundation (NSF) Cooperative Agreement EAR-8920136 and U.S. Geological Survey Cooperative Agreements 14-08-0001-A0899 and 1434-HQ-97AG01718. This is SCEC contribution 433. Partial support was provided by Chapman University and University of California Irvine awards to Grant, NSF grants EAR-9406183, EAR-9512334, and EAR-9712037 to Edwards, and University of Colorado Boulder CRCW grant to Mueller. We thank F. Peska Jr., P. Hughes, M. C. Wallis, J. A. Minch, C. L. Powell, S. W. Conklin, and reviewers C. Prentice and J. Shaw.

REFERENCES CITED

Barrie, D. S., Tatnall, T. S., and Gath, E. M., 1992, Neotectonic uplift and ages of Pleistocene marine terraces, San Joaquin Hills, Orange County, California, *in* Heath, E., and Lewis, L., eds., *The regressive Pleistocene shoreline in southern California*: Santa Ana, California, South Coast Geological Society Annual Field Trip Guidebook 20, p. 115–121.

Bohannon, R. G., and Geist, E., 1998, Upper crustal structure and Neogene tectonic development of the California continental borderland: *Geological Society of America Bulletin*, v. 110, p. 779–900.

Bullard, T. F., and Lettis, W. R., 1993, Quaternary fold deformation associated with blind thrust faulting, Los Angeles basin: California: *Journal of Geophysical Research*, v. 98, p. 8349–8369.

Chappell, J. M., and Shackleton, N. J., 1986, Oxygen isotopes and sea level: *Nature*, v. 324, p. 137–140.

Cheng, H., Edwards, R. L., Hoff, J., Gallup, C. D., Richards, D. A., and Asmerom, Y., 1999, The half-lives of uranium-234 and thorium-230: *Chemical Geology* (in press).

Dolan, J. F., Sieh, K., Rockwell, T. K., Yeats, R. S., Shaw, J., Suppe, J., Huftile, G. J., and Gath, E. M., 1995, Prospects for larger or more frequent earthquakes in the Los Angeles metropolitan region: *Science*, v. 267, p. 199–205.

Edwards, R. L., Chen, J. H., and Wasserburg, G. J., 1987, U-238–U-234–Th-230–Th-232 systematics and the precise measurement of time over the past 500,000 years: *Earth and Planetary Science Letters*, v. 81, p. 175–192.

Fischer, P. J., and Mills, G. I., 1991, The offshore Newport-Inglewood–Rose Canyon fault zone, California: Structure, segmentation and tectonics, *in* Abbott, P. L., and Elliott, W. J., eds., *Environmental perils of the San Diego region*: San Diego, California, San Diego Association of Geologists, p. 17–36.

Gallup, C. D., Edwards, R. L., and Johnson, R. G., 1994, The timing of high sea levels over the past 200,000 years: *Science*, v. 263, p. 796–800.

Grant, L. B., and Ballenger, L., 1999, Holocene uplift and paleoseismology of the San Joaquin Hills, Orange County, California: *Seismological Research Letters*, v. 70, p. 266–267.

Grant, L. B., Waggoner, J. T., Rockwell, T. K., and von Stein, C., 1997, Paleoseismicity of the north branch of the Newport-Inglewood fault zone in Huntington Beach, California, from cone penetrometer test data: *Seismological Society of America Bulletin*, v. 87, p. 277–293.

Hauksson, E., 1990, Earthquakes, faulting and stress in the Los Angeles basin, southern California: *Journal of Geophysical Research*, v. 95, p. 15,365–15,394.

Hauksson, E., and Gross, S., 1991, Source parameters of the 1933 Long Beach earthquake: *Seismological Society of America Bulletin*, v. 81, p. 81–98.

Kanakoff, G. P., and Emerson, W. K., 1959, Late Pleistocene invertebrates of the Newport Bay area, California: *Los Angeles County Museum Contributions in Science*, no. 31, 47 p.

Kennedy, G. L., Lajoie, K. R., and Wehmiller, J. F., 1982, Aminostratigraphy and faunal correlations of late Quaternary marine terraces, Pacific Coast, USA: *Nature*, v. 299, p. 545–547.

Lajoie, K. R., Ponti, D. J., Powell, S. A., Mathieson, S. A., and Sarna-Wojcicki, A. M., 1991, Emergent marine strandlines and associated sediments, coastal California: A record of Quaternary sea-level fluctuations, vertical tectonic movements, climatic changes, and coastal processes, *in* Morrison, R. B., ed., *Quaternary nonglacial geology: Conterminous U.S.*: Boulder, Colorado, Geological Society of America, *Geology of North America*, v. K-2, p. 190–214.

Lettis, W. R., Wells, D. L., and Baldwin, J. N., 1997, Empirical observations regarding reverse earthquakes, blind thrust faults, and Quaternary deformation; are blind thrust faults truly blind?: *Seismological Society of America Bulletin*, v. 87, p. 1171–1198.

Merritts, D., and Bull, W. B., 1989, Interpreting Quaternary uplift rates at the Mendocino triple junction, northern California, from uplifted marine terraces: *Geology*, v. 17, p. 1020–1024.

Muhs, D. R., Kennedy, G. L., and Rockwell, T. K., 1994, Uranium-series ages of marine terrace corals from the Pacific coast of North America and implications for last-interglacial sea level history: *Quaternary Research*, v. 42, p. 72–87.

Peska, F., Jr., 1984, Late Pleistocene fossils from upper Newport Bay, California, *in* Butler, B., et al., eds., *The natural science of Orange County: Natural History Foundation of Orange County Memoir*, v. 1, p. 55–60.

Scientists of USGS and the Southern California Earthquake Center, 1994, The magnitude 6.7 Northridge, California, earthquake of 17 January 1994: *Science*, v. 266, p. 389–397.

Shaw, J. H., and Shearer, P. M., 1999, An elusive blind-thrust fault beneath metropolitan Los Angeles: *Science*, v. 283, p. 1516–1518.

Smith, D. E., and Teggart, F. J., eds., 1909, *Diary of Gaspar de Portola during the California expedition of 1769–1770*: Berkeley, University of California Press, Publications of the Academy of Pacific Coast History, v. 1, no. 3, 59 p.

Sprotte, E. C., Fuller, D. R., Greenwood, R. B., and Mumm, H. A., 1980, Classification and mapping of Quaternary sedimentary deposits for purposes of seismic zonation, south coastal Los Angeles basin, Orange County, California: California Division of Mines and Geology Open File Report 80-19, 250 p.

Stein, M., Wasserburg, G. J., Lajoie, K. R., and Chen, J. H., 1991, TIMS U-series ages of solitary corals from the California coast by mass spectrometry: *Geochimica et Cosmochimica Acta*, v. 55, p. 3709–3722.

Stein, R. S., and Yeats, R. S., 1989, Hidden earthquakes: *Scientific American*, v. 260, p. 48–57.

Stevenson, R. E., 1954, The marshlands at Newport Bay, California [Ph.D. thesis]: Los Angeles, University of Southern California, 199 p.

Vedder, J. G., 1975, Revised geologic map, structure sections and well table, San Joaquin Hills–San Juan Capistrano area, California: U.S. Geological Survey Open-File Report 75-552, scale 1:24000, 3 p.

Vedder, J. G., Yerkes, R. F., and Schoelhamer, J. E., 1957, Geologic map of the San Joaquin Hills–San Juan Capistrano area, Orange County, California: U.S. Geological Survey Oil and Gas Investigations Map OM-193, scale 1:24000.

Ward, S. N., and Valensise, G., 1994, The Palos Verdes terraces, California: Bathub rings from a buried reverse fault: *Journal of Geophysical Research*, v. 99, no. B3, p. 4485–4494.

Wehmiller, J. F., Lajoie, K. R., Kvenvolden, K. A., Peterson, E., Belknap, D. F., Kennedy, G. L., Addicott, W. O., Vedder, J. G., and Wright, R. W., 1977, Correlation and chronology of Pacific Coast marine terrace deposits of continental United States by amino acid stereo-chemistry; technique evaluation, relative ages, kinetic model ages, and geologic implications: U.S. Geological Survey Open-File Report 77-680, 107 p.

Wells, D. L., and Coppersmith, K. J., 1994, New empirical relationships among magnitude, rupture length, rupture area, and surface displacement: *Seismological Society of America Bulletin*, v. 84, p. 974–1002.

Wright, T. L., 1991, Structural geology and tectonic evolution of the Los Angeles basin, *in* Biddle, K. T., ed., *Active margin basins: American Association of Petroleum Geologists Memoir* 52, p. 35–134.

Manuscript received February 8, 1999

Revised manuscript received July 23, 1999

Manuscript accepted August 3, 1999