

receiving PVI at ambulatory surgical centers and office-based labs (ASC/OBL) versus the outpatient hospital (hospital) site of service. **METHODS/STUDY POPULATION:** We performed a retrospective analysis using 100% Medicare fee-for-service claims data between January 1, 2017 and December 31, 2022. We used Current Procedural Terminology (CPT) codes to identify patients undergoing angioplasty, stenting, or atherectomy. Patient demographics were collected from the Medicare Master Beneficiary Summary File and associated comorbidities and PVI indications were identified using International Classification of Disease (ICD)-10 codes. We used patient ZIP codes to determine patients' residence densities and regions. We used site of service codes to determine whether PVI were performed in the ASC/OBL versus hospital. Results were analyzed with descriptive statistics. **RESULTS/ANTICIPATED RESULTS:** Of 817,241 patients undergoing PVI for PAD, 461,068 (56.4%) were treated in an ASC/OBL. Compared to patients treated in the hospital, patients receiving PVI at ASC/OBLs were more likely to be older, female, non-white race, with fewer comorbidities (end stage renal disease, diabetes, hypertension, and any history of tobacco use) (all,  $P < 0.001$ ). Patients treated in ASC/OBLs more frequently resided in urban (vs. rural) locations, and in the South and West (both,  $P < 0.001$ ). Indication for PVI was predominately chronic limb-threatening ischemia, and clinically similar between groups (77.1% vs. 76.2%). There was a significant change in site of service over time: a minority (47.6%) of PVIs were performed in the ASC/OBL in 2017, whereas the majority (64.7%) of PVIs were performed in the ASC/OBL in 2022 ( $P < 0.001$ ). **DISCUSSION/SIGNIFICANCE:** Patients treated in ASC/OBLs were less medically complex compared to those treated in the outpatient hospital setting. Further study is needed to examine whether differences in patient characteristics versus other factors (e.g. reimbursement) are driving the increase in PVIs performed in the ASC/OBL over time.

21

### The Microbial Antibigram as a Function of Testing Indication: Susceptibility Analysis of *Escherichia coli* from Symptomatic and Asymptomatic Bacteriuria Patients, 2020-2021

Allison Chan<sup>1</sup>, Maddie Spradley<sup>1</sup>, Tim Williams<sup>1</sup>, Grace Morales<sup>1</sup>, Gerald Van Horn<sup>1</sup> and Jonathan E. Schmitz<sup>1,2</sup>

<sup>1</sup>Vanderbilt University Medical Center, Department of Pathology, Microbiology, and Immunology; Nashville, Tennessee 37232 USA and <sup>2</sup>Department of Urology, Vanderbilt Institute for Infection, Immunology, and Inflammation; Nashville, Tennessee 37232 USA

**OBJECTIVES/GOALS:** Antibigrams are used to guide empiric antibiotic selection. However, it is unclear if antibiotic profiles differ between symptomatic urinary tract infections (UTIs) and asymptomatic bacteriuria (ASB). We aimed to compare antibiotic susceptibility profiles of urinary *E. coli* isolates from patients with a symptomatic UTI to those with ASB. **METHODS/STUDY POPULATION:** We conducted a cohort study of 1,140 urinary *E. coli* isolates from unique patients that received care through Vanderbilt University Medical Center (VUMC) from Nov 2020 – Jun 2021. We included any patient that was seen at VUMC as an inpatient, outpatient or at the emergency department with  $\geq 10^5$  colony forming units/mL *E. coli* detected from a clinical urine specimen. Chart abstractions were performed to capture reported UTI symptoms and demographic information. Descriptive statistics were

conducted to compare antibiotic susceptibility profiles (i.e., susceptible, intermediate, resistant) between symptomatic and ASB groups. The risk of detection of a multidrug-resistant organism (MDRO) (intermediate, or resistant to at least one antibiotic in three or more classes) was assessed between groups. **RESULTS/ANTICIPATED RESULTS:** Among 1,140, 1,018 (89%) and 122 (11%) were symptomatic and ASB, respectively. When comparing symptomatic and ASB, the median ages were 50 and 46. Groups had similar proportions of no indwelling catheter (94% v. 95%) and without diabetes (87% v. 88%). The collection setting between inpatient, emergency department, and outpatient were similar with most being outpatient (79% v. 83%). The proportion of patients who were pregnant, immuno compromised, or had a structural/functional urinary tract abnormality were higher in the symptomatic group. The proportion of isolates resistant and susceptible to tested antibiotics were similar between groups, with only ciprofloxacin showing slightly higher resistance among ASB (16% v. 25%). The risk of MDRO detection was similar between groups (RR: 0.858, 95% CI: 0.64, 1.15). **DISCUSSION/SIGNIFICANCE:** Antibiotic susceptibility comparison demonstrated similar profiles, which suggests antibiogram use as appropriate to guide ASB treatment. Results offer insight on whether traditional methods for assessing antibiotic susceptibility on population-levels could benefit from further refinement by patient-specific clinical parameters.

23

### AHA's Essential Eight: Opportunities for Preventive Care among Adults with Peripheral Arterial Disease

Edward Huang and Elizabeth A. Jackson

University of Alabama at Birmingham

**OBJECTIVES/GOALS:** The objective was to examine the American Heart Association's (AHA) Essential Eight metrics of cardiovascular (CV) health among Black and White adults with peripheral arterial disease (PAD) collected via validated surveys and medical records. Each metric was examined in association with available social determinants of health (SDoH) factors. **METHODS/STUDY POPULATION:** This observational study completed data collection through surveys and medical record review. Validated surveys were used to collect Essential Eight metrics of diet, physical activity, sleep, and smoking status. Medical records were used to collect data on body mass index, blood lipids, blood glucose, and blood pressure. Participants with a diagnosis of lower extremity PAD, ability to complete surveys, and provided informed written consent were eligible. Equal numbers of Black and White participants were enrolled. Essential Eight metrics were used to calculate CV health scores for each participant. Scores were examined for association with SDoH factors and by race using Student's T-test or ANOVA for continuous variables or Chi-Square tests for categorical variables. **RESULTS/ANTICIPATED RESULTS:** A total of 50 participants will be enrolled, with the expected majority being men and half self-reporting as Black individuals. Worse SDoH is expected to be associated with lower CV health metrics, including lower levels of physical activity and higher levels of saturated fatty food consumption. Higher levels of blood lipids, blood glucose, and blood pressure are expected to be associated with worse SDoH factors. We expect this association to be attenuated by rates of CV medications, such as statin therapy, antidiabetic medications, and antihypertensive medications. No effect modification by rurality is expected, although