



# OLDER ADULTS BEHAVIORAL HEALTH PROFILES

## REGION 8

COLORADO  
MONTANA  
NORTH DAKOTA  
SOUTH DAKOTA  
UTAH  
WYOMING



**A Behavioral Health Resource**  
SAMHSA's State Technical Assistance Contract  
August 2016

# OLDER ADULTS BEHAVIORAL HEALTH PROFILES

---

## OVERVIEW

The Substance Abuse and Mental Health Services Administration recognizes the importance of behavioral health for older adults and the key role states and communities play in addressing the needs of this vital and growing population. Administrators need clear and concise information to plan, implement, and evaluate effective prevention and treatment efforts in their states.

The Older Adults Behavioral Health Profiles help states and communities identify focus areas for behavioral health plans, select population-level goals, and coordinate and target services to address priority issues. The profiles compare state trends with those in the region and the nation. State and community administrators, planners, and providers can use the information in these profiles and their own data, knowledge, and experience to establish and implement policies that improve the health and future of our valued older citizens.

Disclaimer: Data were current at the time this document was prepared.

# OLDER ADULTS BEHAVIORAL HEALTH PROFILE

---

## Colorado

# Colorado

## OLDER ADULTS BEHAVIORAL HEALTH PROFILE

August 2016

### COLORADO'S POPULATION

#### Colorado Population by Age Group

Colorado is home to 5,355,866 people. Of these:

- 1,721,823 (32.1 percent) are over age 50.
- 991,802 (18.5 percent) are over age 60.
- 433,786 (8.1 percent) are over age 70.
- 159,760 (3.0 percent) are ages 80 and older.

The proportion of women rises fairly steadily in each age group, and women make up 61.3 percent of the 80+ group. The racial/ethnic composition of older Coloradans is as follows:

#### Race/Ethnicity of Coloradans Ages 50+


White	AI/AN	Black	Asian	NH/PI	Other	Hispanic
91.9%	1.1%	3.4%	2.4%	0.1%	1.1%	11.9%

Source: U.S. Census Bureau, 2015

AI/AN stands for American Indian and Alaska Native.

NH/PI stands for Native Hawaiian and Other Pacific Islander.


Exhibit 1. Colorado Population by Age Group, 2014



Source: U.S. Census Bureau, 2015

#### The Number of Older Coloradans Is Growing

Exhibit 2. Colorado Population by Age Group, 2000–2030



The proportion of Colorado's population that is 65 and older is growing while the proportion that is younger than 65 is shrinking. The U.S. Census Bureau estimates that 16.5 percent of Colorado's population will be 65 and older by the year 2030, an increase of 52.4 percent from 2015.

#### Projected Population in Colorado

Age Group	2015	2025	2030
Under 18	24.9%	25.2%	25.3%
18 to 44	37.3%	36.8%	37.0%
45 to 64	25.3%	22.2%	21.2%
65+	12.4%	15.7%	16.5%

Source: U.S. Census Bureau, 2005


## SUICIDE AMONG OLDER COLORADANS

### Colorado Suicide Rate Compared With Regional and National Rates

The suicide rate among Coloradans ages 50 and older is higher than the rate among younger age groups. In 2013, the total suicide rate among all people ages 50+ was 26.4 per 100,000 people (12.9 for women and 41.3 for men). The rate among those ages 50–64 was lower than the rate in the region (including Montana, North Dakota, South Dakota, Utah, and Wyoming) and higher than the rate in the United States.

States vary in their reporting of suicides. The suicide rate is influenced by these reporting practices.


**Exhibit 3. Suicide Rates in Colorado, Region 8, and the United States, 2013**



Source: Centers for Disease Control and Prevention (CDC), National Center for Health Statistics (NCHS), Underlying Cause of Death 1999-2013 on CDC WONDER Online Database, released 2015

### Trends in Suicide Rates in Colorado

**Exhibit 4. Trends in Suicide Rates in Colorado by Age Group, 2004-2013**



Source: CDC, NCHS, Underlying Cause of Death 1999-2013 on CDC WONDER Online Database, released 2015

The suicide rate among Coloradans ages 50+ fluctuated from a low of 19.1 per 100,000 in 2006 to a high of 26.8 per 100,000 in 2012. From 2004 to 2013, the rate was generally highest among those in the 50–64 age group.


How a state reports suicides can vary from year to year. The number of suicides is generally low, so even a small difference in reported numbers may make the rate fluctuate widely.

## SUBSTANCE USE DISORDER AND SUBSTANCE USE DISORDER TREATMENT AMONG OLDER COLORADANS

### 30-Day Binge Drinking Among Older Coloradans

Binge drinking, defined as five or more drinks for men and four or more drinks for women on a single occasion, can lead to serious health problems. Such problems include neurological damage, cardiovascular disease, liver disease, stroke, and poor control of diabetes. Binge drinkers are more likely to take risks such as driving while intoxicated and to experience falls and other accidents. Older people have a lower tolerance for alcohol. Binge drinking decreases with age and occurs more frequently among men than it does among women. As Exhibit 5 shows, 16.3 percent of Colorado men ages 50–64 reported binge drinking in the past 30 days, while 6.6 percent of those in the 65+ group reported similar behavior.


Exhibit 5. Binge Drinking Rates in Colorado by Age Group and Sex, 2013



Source: Behavioral Risk Factor Surveillance System (BRFSS), 2013

### Illicit Drug Use Among Older Americans

Exhibit 6. Illicit Drug Use Among Older Americans, 2002–2013



Source: National Survey on Drug Use and Health (NSDUH), 2013  
 Illicit drug use includes illegal drugs and prescription drugs used nonmedically. SAMHSA provides a list of drugs included in its survey in the NSDUH methodological summary.

Nationally, the rate of illicit drug use among older adults ages 50–64 more than doubled from 2002 to 2013. This development partially reflects the entrance into this age group of the baby boom cohort, whose rates of illicit drug use have been higher than those of older cohorts. Although state-specific data are not available, the *Colorado Behavioral Health Barometer* is available for download from the Substance Abuse and Mental Health Services Administration (SAMHSA) website ([www.samhsa.gov/data/population-data-nsduh/reports?tab=33](http://www.samhsa.gov/data/population-data-nsduh/reports?tab=33)).


## Admissions to Substance Use Disorder Treatment Among Older Coloradans

In 2012, there were 18,213 admissions of Coloradans ages 50 and older to substance use disorder (SUD) treatment in state-funded treatment programs, a rate of 1,057.8 per 100,000 people ages 50+. This rate was higher than the regional rate and the national average. Men made up 82.0 percent of these admissions. Of all admissions, 83.0 percent were White/Caucasian, 10.8 percent were Black/African American, and 24.6 percent were Hispanic.

The principal sources of referral to treatment among those ages 50 and older were:

Self-Referral	Criminal Justice	Other
18.4%	27.4%	54.2%


**Exhibit 7. SUD Treatment Admissions of Adults Ages 50+ in Colorado, Region 8, and the United States by Sex, 2012**



Source: Treatment Episode Data Set (TEDS), 2012<sup>1</sup>  
Data include only those clients reported to SAMHSA.

## SUD Treatment Admissions Among Coloradans Ages 50+ by Insurance Type

**Exhibit 8. SUD Treatment Admissions of Adults Ages 50+ in Colorado, Region 8, and the United States by Insurance Type, 2012**



Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

In Colorado, 67.3 percent of older adult admissions to SUD treatment were uninsured, 3.5 percent had Medicaid, 2.7 percent had Medicare, and 26.4 percent had private insurance.

### *SUD Treatment Admissions Among Coloradans Ages 50+ by Primary Sources of Payment*

Self-Pay	Medicaid	Other
32.8%	4.1%	63.1%


Self-pay data include private insurance. Exhibit 8 and the accompanying text reflect the type of health insurance (if any) clients had at admission; the table above represents primary sources of payment.

<sup>1</sup> TEDS data are collected by states as a condition of the Substance Abuse Prevention and Treatment Block Grant. Guidelines suggest that states report all clients admitted to publicly financed treatment; however, states are inconsistent in applying the guidelines. States may structure and implement different quality controls over the data. For example, states may collect different categories of information to answer TEDS questions. Information is then "walked over" to TEDS definitions.

## Alcohol Use Disorder Treatment Admissions Among Coloradans Ages 50+

Alcohol was the most frequently cited substance used by older Coloradans in publicly financed SUD treatment in 2012. It was mentioned as a substance of use in 88.8 percent of admissions among those ages 50+. This was higher than the regional and national rates.


**Exhibit 9. Alcohol Use Treatment Admissions of Adults Ages 50+ in Colorado, Region 8, and the United States, 2012**



Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

## SUD Treatment Admissions for Non-Alcohol Substance Use

**Exhibit 10. Non-Alcohol Substance Use Treatment Admissions of Adults Ages 50+ in Colorado, Region 8, and the United States, 2012**



Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

Substances other than alcohol were cited as the primary substances of use for 11.2 percent of older adult admissions to publicly funded treatment in Colorado.



## Drug-Related Emergency Department Visits Involving Pharmaceutical Misuse and Abuse by Older Adults

SAMHSA periodically releases reports from the Drug Abuse Warning Network (DAWN). DAWN, discontinued in 2011, consisted of a nationwide network of hospital emergency departments (EDs) primarily located in large metropolitan areas. DAWN data consist of professional reviews of ED records to determine the extent to which alcohol and other substance abuse was involved in ED visits. According to the November 25, 2010, *DAWN Report*:


- *In 2004, there were an estimated 115,803 ED visits involving pharmaceutical misuse and abuse by adults aged 50 or older; in 2008, there were 256,097 such visits, representing an increase of 121.1 percent*
- *One fifth (19.7 percent) of ED visits involving pharmaceutical misuse and abuse among older adults were made by persons aged 70 or older*
- *Among ED visits made by older adults, pain relievers were the type of pharmaceutical most commonly involved (43.5 percent), followed by drugs used to treat anxiety or insomnia (31.8 percent) and antidepressants (8.6 percent)*
- *Among patients aged 50 or older who visited the ED for pharmaceutical misuse or abuse, more than half (52.3 percent) were treated and released, and more than one third (37.5 percent) were admitted to the hospital*

Bulleted text in italics above is taken from SAMHSA's Center for Behavioral Health Statistics and Quality. (2010, November 25). *The DAWN Report: Drug-related emergency department visits involving pharmaceutical misuse and abuse by older adults*. Rockville, MD: SAMHSA.

## CO-OCCURRING SUBSTANCE USE AND MENTAL DISORDERS

### Older Coloradans in SUD Treatment With Co-Occurring Mental Disorders

**Exhibit 11. SUD Treatment Admissions of Adults Ages 50+ With a Co-Occurring Mental Disorder in Colorado, Region 8, and the United States, 2012**



Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

The national literature shows a strong relationship between substance use and mental disorders. Studies show that 30–80 percent of individuals with a substance use or mental disorder also have a co-occurring disorder.


Exhibit 11 shows the proportion of SUD treatment admissions of Coloradans ages 50+ with a co-occurring mental disorder. This rate is lower than the regional rate and the national average. However, state reporting practices are a factor in these results.

## MENTAL HEALTH

### Older Coloradans Reporting Frequent Mental Distress

BRFSS, a household survey conducted in all 50 states and several territories, asks the following question: "Now thinking about your mental health, which includes stress, depression, and problems with emotions, for how many days during the past 30 days was your mental health not good?" Those individuals reporting 14 or more "Yes" days in response to this question experience frequent mental distress (FMD). Exhibit 12 shows that older Coloradans experience FMD at a rate that is lower than the regional and national rates.


**Exhibit 12. Individuals Reporting Frequent Mental Distress in Colorado, Region 8, and the United States, 2013**



Source: BRFSS, 2013

### Older Coloradans Reporting Frequent Mental Distress by Age Group and Sex

**Exhibit 13. Coloradans Reporting Frequent Mental Distress by Age Group and Sex, 2013**



Source: BRFSS, 2013

Older men in Colorado were more likely to indulge in binge drinking, but older women were more likely to report that they had FMD (14 days or more per 30-day period). As Exhibit 13 shows, 11.3 percent of women in the 50–64 age group and 6.9 percent in the 65+ age group reported FMD, while 7.7 percent of men in the 50–64 age group and 3.9 percent in the 65+ age group reported FMD.

## Other Measures of Mental Health

BRFSS collected other measures showing risk factors for mental and/or physical illness. These included:

- Social and Emotional Support (2010). BRFSS asked, "How often do you get the social and emotional support you need?" The possible responses were always, usually, sometimes, rarely, or never.
- Life Satisfaction (2010). BRFSS asked, "In general, how satisfied are you with your life?" The possible responses were very satisfied, satisfied, dissatisfied, or very dissatisfied.

Exhibit 14 presents the results of these surveys among older Coloradans.

**Exhibit 14. BRFSS Measures, 2010**


Indicator	Ages 50+	Ages 50–64	Ages 65+
Rarely or never get social or emotional support	7.6%	6.5%	9.6%
Dissatisfied or very dissatisfied	4.3%	4.9%	3.3%

Source: BRFSS, 2010

## Individuals With Frequent Mental Distress Report High Rates of Poor Physical Health

Older Americans who experienced FMD were more likely to report that their physical health was poor (14 days or more in the past 30-day period when physical health was "not good"). As shown in Exhibit 15, although nearly 11 percent of older Americans with no mental distress reported poor physical health, more than 50 percent of those with FMD reported poor physical health.

**Exhibit 15. Individuals Ages 50+ in the United States Reporting Poor Physical Health by Level of Mental Distress, 2013**




Source: BRFSS, 2013

## Relationship Among Mental Distress, Diabetes, Stroke, Heart Attack, High Blood Pressure, and Coronary Disease

Older Americans who experience FMD are more likely to report that they have health problems. People with FMD were more than twice as likely to report having a stroke than those with some or no mental distress. They experienced coronary disease and heart attack at more than 1.6 times, diabetes at more than 1.4 times, and high blood pressure at nearly 1.2 times the rate of those with some or no mental distress.

**Exhibit 16. Individuals Ages 50+ in the United States With Mental Distress and Serious Health Problems, 2013**



Source: BRFSS, 2013

## Older Coloradans Admitted to State Mental Health Services

Approximately 2.9 percent of the people served by the Colorado mental health system were ages 65 and older. This represents more than 3,330 adults.

Source: Center for Mental Health Services (CMHS) Uniform Reporting System (URS) Output Tables, 2014

## DATA SOURCES

**BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM** ([www.cdc.gov/brfss](http://www.cdc.gov/brfss)). BRFSS, managed by CDC, is the largest ongoing telephone health survey system in the world, tracking health conditions and risk behaviors in the United States annually since 1984. Data are collected on a monthly basis in all 50 states, the District of Columbia, and participating territories. BRFSS data are collected by local jurisdictions and reported to CDC.

**CENTERS FOR DISEASE CONTROL AND PREVENTION, NATIONAL CENTER FOR HEALTH STATISTICS, UNDERLYING CAUSE OF DEATH 1999-2013 ON CDC WONDER ONLINE DATABASE, RELEASED 2015** (<http://wonder.cdc.gov/ucd-icd10.html>). The WONDER online database “contains mortality and population counts for all U.S. counties. Data are based on death certificates for U.S. residents. Each death certificate identifies a single underlying cause of death and demographic data. The number of deaths, crude death rates or age-adjusted death rates, and 95% confidence intervals and standard errors for death rates can be obtained by place of residence (total U.S., region, state and county), age group (single-year-of age, 5-year age groups, 10-year age groups and infant age groups), race, Hispanic ethnicity, gender, year, cause-of-death (4-digit ICD-10 code or group of codes), injury intent and injury mechanism, drug/alcohol induced causes and urbanization categories. Data are also available for place of death, month and week day of death, and whether an autopsy was performed.”

**CENTER FOR MENTAL HEALTH SERVICES UNIFORM REPORTING SYSTEM** ([www.samhsa.gov/data/us\\_map](http://www.samhsa.gov/data/us_map)). States that receive CMHS Block Grants are required to report aggregate data to URS. URS reports include information about utilization of mental health services and client demographic and outcome information.

**NATIONAL SURVEY ON DRUG USE AND HEALTH** (<https://nsduhweb.rti.org>). The NSDUH, managed by SAMHSA, is “an annual nationwide survey involving interviews with approximately 70,000 randomly selected individuals aged 12 and older.” NSDUH data are most frequently used by state planners to assess the need for SUD treatment. NSDUH data also include information about mental health conditions.

**TREATMENT EPISODE DATA SET** ([www.samhsa.gov/data/client-level-data-teds/reports?tab=19](http://www.samhsa.gov/data/client-level-data-teds/reports?tab=19)). States that receive Substance Abuse Prevention and Treatment Block Grant funds submit individual client data to TEDS. TEDS includes both admission and discharge data sets, and some 1.5 million admissions are reported annually. TEDS includes information about utilization of SUD treatment services and client demographic and outcome information. TEDS is an admission-based system, and TEDS admissions do not represent individuals. Thus, an individual admitted to treatment twice within a calendar year would be counted as two admissions.

**U.S. CENSUS BUREAU** ([www.census.gov/people](http://www.census.gov/people)). Two main sources of Census Bureau data were used in this report: (1) population estimates and (2) population projections.

# OLDER ADULTS BEHAVIORAL HEALTH PROFILE

---

## Montana

# Montana

## OLDER ADULTS BEHAVIORAL HEALTH PROFILE

August 2016

### MONTANA'S POPULATION

#### Montana Population by Age Group

Montana is home to 1,023,579 people. Of these:

- 393,962 (38.5 percent) are over age 50.
- 243,992 (23.8 percent) are over age 60.
- 113,167 (11.1 percent) are over age 70.
- 42,389 (4.1 percent) are ages 80 and older.


The proportion of women rises fairly steadily in each age group, and women make up 59.5 percent of the 80+ group. The racial/ethnic composition of older Montanans is as follows:

#### Race/Ethnicity of Montanans Ages 50+

White	AI/AN	Black	Asian	NH/PI	Other	Hispanic
94.2%	3.8%	0.2%	0.5%	0.1%	1.2%	1.5%

Source: U.S. Census Bureau, 2015  
AI/AN stands for American Indian and Alaska Native.  
NH/PI stands for Native Hawaiian and Other Pacific Islander.


Exhibit 1. Montana Population by Age Group, 2014



Source: U.S. Census Bureau, 2015

#### The Number of Older Montanans Is Growing

Exhibit 2. Montana Population by Age Group, 2000–2030



The proportion of Montana's population that is 65 and older is growing while the proportion that is younger than 65 is shrinking. The U.S. Census Bureau estimates that 25.8 percent of Montana's population will be 65 and older by the year 2030, an increase of 55.1 percent from 2015.

#### Projected Population in Montana

Age Group	2015	2025	2030
Under 18	21.6%	20.8%	20.1%
18 to 44	32.6%	30.2%	28.4%
45 to 64	28.4%	25.2%	25.7%
65+	17.4%	23.9%	25.8%

Source: U.S. Census Bureau, 2005


## SUICIDE AMONG OLDER MONTANANS

### Montana Suicide Rate Compared With Regional and National Rates

The suicide rate among Montanans ages 50 and older is higher than the rate among younger age groups. In 2013, the total suicide rate among all people ages 50+ was 26.5 per 100,000 people (10.0 for women and 44.0 for men). The rate among those ages 50–64 was higher than both the rate in the region (including Colorado, North Dakota, South Dakota, Utah, and Wyoming) and the rate in the United States.

States vary in their reporting of suicides. The suicide rate is influenced by these reporting practices.


Exhibit 3. Suicide Rates in Montana, Region 8, and the United States, 2013



Source: Centers for Disease Control and Prevention (CDC), National Center for Health Statistics (NCHS), Underlying Cause of Death 1999–2013 on CDC WONDER Online Database, released 2015

### Trends in Suicide Rates in Montana

Exhibit 4. Trends in Suicide Rates in Montana by Age Group, 2004–2013



Source: CDC, NCHS, Underlying Cause of Death 1999–2013 on CDC WONDER Online Database, released 2015

The suicide rate among Montanans ages 50+ fluctuated from a low of 20.7 per 100,000 in 2004 to a high of 29.9 per 100,000 in 2010. From 2004 to 2013, the rate was generally highest among those in the 50–64 age group.

How a state reports suicides can vary from year to year. The number of suicides is generally low, so even a small difference in reported numbers may make the rate fluctuate widely.




## SUBSTANCE USE DISORDER AND SUBSTANCE USE DISORDER TREATMENT AMONG OLDER MONTANANS

### 30-Day Binge Drinking Among Older Montanans

Binge drinking, defined as five or more drinks for men and four or more drinks for women on a single occasion, can lead to serious health problems. Such problems include neurological damage, cardiovascular disease, liver disease, stroke, and poor control of diabetes. Binge drinkers are more likely to take risks such as driving while intoxicated and to experience falls and other accidents. Older people have a lower tolerance for alcohol. Binge drinking decreases with age and occurs more frequently among men than it does among women. As Exhibit 5 shows, 23.7 percent of Montana men ages 50–64 reported binge drinking in the past 30 days, while 7.4 percent of those in the 65+ group reported similar behavior.


Exhibit 5. Binge Drinking Rates in Montana by Age Group and Sex, 2013



Source: Behavioral Risk Factor Surveillance System (BRFSS), 2013

### Illicit Drug Use Among Older Americans

Exhibit 6. Illicit Drug Use Among Older Americans, 2002–2013



Source: National Survey on Drug Use and Health (NSDUH), 2013  
 Illicit drug use includes illegal drugs and prescription drugs used nonmedically. SAMHSA provides a list of drugs included in its survey in the NSDUH methodological summary.

Nationally, the rate of illicit drug use among older adults ages 50–64 more than doubled from 2002 to 2013. This development partially reflects the entrance into this age group of the baby boom cohort, whose rates of illicit drug use have been higher than those of older cohorts. Although state-specific data are not available, the *Montana Behavioral Health Barometer* is available for download from the Substance Abuse and Mental Health Services Administration (SAMHSA) website ([www.samhsa.gov/data/population-data-nsduh/reports?tab=33](http://www.samhsa.gov/data/population-data-nsduh/reports?tab=33)).


## Admissions to Substance Use Disorder Treatment Among Older Montanans

In 2012, there were 1,160 admissions of Montanans ages 50 and older to substance use disorder (SUD) treatment in state-funded treatment programs, a rate of 294.4 per 100,000 people ages 50+. This rate was lower than the regional rate and the national average. Men made up 75.7 percent of these admissions. Of all admissions, 79.3 percent were White/Caucasian, 0.4 percent were Black/African American, and 2.3 percent were Hispanic.

The principal sources of referral to treatment among those ages 50 and older were:

Self-Referral	Criminal Justice	Other
19.8%	58.6%	21.6%


**Exhibit 7. SUD Treatment Admissions of Adults Ages 50+ in Montana, Region 8, and the United States by Sex, 2012**



Source: Treatment Episode Data Set (TEDS), 2012<sup>1</sup>  
Data include only those clients reported to SAMHSA.

## SUD Treatment Admissions Among Montanans Ages 50+ by Insurance Type

**Exhibit 8. SUD Treatment Admissions of Adults Ages 50+ in Montana, Region 8, and the United States by Insurance Type, 2012**



Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

In Montana, 70.5 percent of older adult admissions to SUD treatment were uninsured, 5.4 percent had Medicaid, 13.4 percent had Medicare, and 10.8 percent had private insurance.

### *SUD Treatment Admissions Among Montanans Ages 50+ by Primary Sources of Payment*

Self-Pay	Medicaid	Other
25.9%	4.7%	69.3%


Self-pay data include private insurance. Exhibit 8 and the accompanying text reflect the type of health insurance (if any) clients had at admission; the table above represents primary sources of payment.

<sup>1</sup> TEDS data are collected by states as a condition of the Substance Abuse Prevention and Treatment Block Grant. Guidelines suggest that states report all clients admitted to publicly financed treatment; however, states are inconsistent in applying the guidelines. States may structure and implement different quality controls over the data. For example, states may collect different categories of information to answer TEDS questions. Information is then "walked over" to TEDS definitions.

## Alcohol Use Disorder Treatment Admissions Among Montanans Ages 50+

Alcohol was the most frequently cited substance used by older Montanans in publicly financed SUD treatment in 2012. It was mentioned as a substance of use in 84.3 percent of admissions among those ages 50+. This was lower than the regional rate and higher than the national rate.

**Exhibit 9. Alcohol Use Treatment Admissions of Adults Ages 50+ in Montana, Region 8, and the United States, 2012**




Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

## SUD Treatment Admissions for Non-Alcohol Substance Use

**Exhibit 10. Non-Alcohol Substance Use Treatment Admissions of Adults Ages 50+ in Montana, Region 8, and the United States, 2012**

Substances other than alcohol were cited as the primary substances of use for 15.7 percent of older adult admissions to publicly funded treatment in Montana.



Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

## Drug-Related Emergency Department Visits Involving Pharmaceutical Misuse and Abuse by Older Adults

SAMHSA periodically releases reports from the Drug Abuse Warning Network (DAWN). DAWN, discontinued in 2011, consisted of a nationwide network of hospital emergency departments (EDs) primarily located in large metropolitan areas. DAWN data consist of professional reviews of ED records to determine the extent to which alcohol and other substance abuse was involved in ED visits. According to the November 25, 2010, *DAWN Report*:


- *In 2004, there were an estimated 115,803 ED visits involving pharmaceutical misuse and abuse by adults aged 50 or older; in 2008, there were 256,097 such visits, representing an increase of 121.1 percent*
- *One fifth (19.7 percent) of ED visits involving pharmaceutical misuse and abuse among older adults were made by persons aged 70 or older*
- *Among ED visits made by older adults, pain relievers were the type of pharmaceutical most commonly involved (43.5 percent), followed by drugs used to treat anxiety or insomnia (31.8 percent) and antidepressants (8.6 percent)*
- *Among patients aged 50 or older who visited the ED for pharmaceutical misuse or abuse, more than half (52.3 percent) were treated and released, and more than one third (37.5 percent) were admitted to the hospital*

Bulleted text in italics above is taken from SAMHSA's Center for Behavioral Health Statistics and Quality. (2010, November 25). *The DAWN Report: Drug-related emergency department visits involving pharmaceutical misuse and abuse by older adults*. Rockville, MD: SAMHSA.

## CO-OCCURRING SUBSTANCE USE AND MENTAL DISORDERS

### Older Montanans in SUD Treatment With Co-Occurring Mental Disorders

**Exhibit 11. SUD Treatment Admissions of Adults Ages 50+ With a Co-Occurring Mental Disorder in Montana, Region 8, and the United States, 2012**



Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

The national literature shows a strong relationship between substance use and mental disorders. Studies show that 30–80 percent of individuals with a substance use or mental disorder also have a co-occurring disorder.


Exhibit 11 shows the proportion of SUD treatment admissions of Montanans ages 50+ with a co-occurring mental disorder. This rate is higher than the regional rate and lower than the national average. However, state reporting practices are a factor in these results.

## MENTAL HEALTH

### Older Montanans Reporting Frequent Mental Distress

BRFSS, a household survey conducted in all 50 states and several territories, asks the following question: “Now thinking about your mental health, which includes stress, depression, and problems with emotions, for how many days during the past 30 days was your mental health not good?” Those individuals reporting 14 or more “Yes” days in response to this question experience frequent mental distress (FMD). Exhibit 12 shows that older Montanans experience FMD at a rate that is higher than the regional rate and lower than the national rate.


**Exhibit 12. Individuals Reporting Frequent Mental Distress in Montana, Region 8, and the United States, 2013**



Source: BRFSS, 2013

### Older Montanans Reporting Frequent Mental Distress by Age Group and Sex

**Exhibit 13. Montanans Reporting Frequent Mental Distress by Age Group and Sex, 2013**



Source: BRFSS, 2013

Older men in Montana were more likely to indulge in binge drinking, but older women were more likely to report that they had FMD (14 days or more per 30-day period). As Exhibit 13 shows, 14.1 percent of women in the 50–64 age group and 7.8 percent in the 65+ age group reported FMD, while 10.1 percent of men in the 50–64 age group and 6.1 percent in the 65+ age group reported FMD.

## Other Measures of Mental Health

BRFSS collected other measures showing risk factors for mental and/or physical illness. These included:

- Social and Emotional Support (2010). BRFSS asked, "How often do you get the social and emotional support you need?" The possible responses were always, usually, sometimes, rarely, or never.
- Life Satisfaction (2010). BRFSS asked, "In general, how satisfied are you with your life?" The possible responses were very satisfied, satisfied, dissatisfied, or very dissatisfied.

Exhibit 14 presents the results of these surveys among older Montanans.

**Exhibit 14. BRFSS Measures, 2010**


Indicator	Ages 50+	Ages 50–64	Ages 65+
Rarely or never get social or emotional support	9.7%	7.6%	12.9%
Dissatisfied or very dissatisfied	4.9%	5.6%	3.7%

Source: BRFSS, 2010

## Individuals With Frequent Mental Distress Report High Rates of Poor Physical Health

Older Americans who experienced FMD were more likely to report that their physical health was poor (14 days or more in the past 30-day period when physical health was "not good"). As shown in Exhibit 15, although nearly 11 percent of older Americans with no mental distress reported poor physical health, more than 50 percent of those with FMD reported poor physical health.

**Exhibit 15. Individuals Ages 50+ in the United States Reporting Poor Physical Health by Level of Mental Distress, 2013**




Source: BRFSS, 2013

## Relationship Among Mental Distress, Diabetes, Stroke, Heart Attack, High Blood Pressure, and Coronary Disease

Older Americans who experience FMD are more likely to report that they have health problems. People with FMD were more than twice as likely to report having a stroke than those with some or no mental distress. They experienced coronary disease and heart attack at more than 1.6 times, diabetes at more than 1.4 times, and high blood pressure at nearly 1.2 times the rate of those with some or no mental distress.

**Exhibit 16. Individuals Ages 50+ in the United States With Mental Distress and Serious Health Problems, 2013**



Source: BRFSS, 2013

## Older Montanans Admitted to State Mental Health Services

Approximately 6.3 percent of the people served by the Montana mental health system were ages 65 and older. This represents more than 2,500 adults.

Source: Center for Mental Health Services (CMHS) Uniform Reporting System (URS) Output Tables, 2014

## DATA SOURCES

**BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM** ([www.cdc.gov/brfss](http://www.cdc.gov/brfss)). BRFSS, managed by CDC, is the largest ongoing telephone health survey system in the world, tracking health conditions and risk behaviors in the United States annually since 1984. Data are collected on a monthly basis in all 50 states, the District of Columbia, and participating territories. BRFSS data are collected by local jurisdictions and reported to CDC.

**CENTERS FOR DISEASE CONTROL AND PREVENTION, NATIONAL CENTER FOR HEALTH STATISTICS, UNDERLYING CAUSE OF DEATH 1999-2013 ON CDC WONDER ONLINE DATABASE, RELEASED 2015** (<http://wonder.cdc.gov/ucd-icd10.html>). The WONDER online database "contains mortality and population counts for all U.S. counties. Data are based on death certificates for U.S. residents. Each death certificate identifies a single underlying cause of death and demographic data. The number of deaths, crude death rates or age-adjusted death rates, and 95% confidence intervals and standard errors for death rates can be obtained by place of residence (total U.S., region, state and county), age group (single-year-of age, 5-year age groups, 10-year age groups and infant age groups), race, Hispanic ethnicity, gender, year, cause-of-death (4-digit ICD-10 code or group of codes), injury intent and injury mechanism, drug/alcohol induced causes and urbanization categories. Data are also available for place of death, month and week day of death, and whether an autopsy was performed."

**CENTER FOR MENTAL HEALTH SERVICES UNIFORM REPORTING SYSTEM** ([www.samhsa.gov/data/us\\_map](http://www.samhsa.gov/data/us_map)). States that receive CMHS Block Grants are required to report aggregate data to URS. URS reports include information about utilization of mental health services and client demographic and outcome information.

**NATIONAL SURVEY ON DRUG USE AND HEALTH** (<https://nsduhweb.rti.org>). The NSDUH, managed by SAMHSA, is "an annual nationwide survey involving interviews with approximately 70,000 randomly selected individuals aged 12 and older." NSDUH data are most frequently used by state planners to assess the need for SUD treatment. NSDUH data also include information about mental health conditions.

**TREATMENT EPISODE DATA SET** ([www.samhsa.gov/data/client-level-data-teds/reports?tab=19](http://www.samhsa.gov/data/client-level-data-teds/reports?tab=19)). States that receive Substance Abuse Prevention and Treatment Block Grant funds submit individual client data to TEDS. TEDS includes both admission and discharge data sets, and some 1.5 million admissions are reported annually. TEDS includes information about utilization of SUD treatment services and client demographic and outcome information. TEDS is an admission-based system, and TEDS admissions do not represent individuals. Thus, an individual admitted to treatment twice within a calendar year would be counted as two admissions.

**U.S. CENSUS BUREAU** ([www.census.gov/people](http://www.census.gov/people)). Two main sources of Census Bureau data were used in this report: (1) population estimates and (2) population projections.



# OLDER ADULTS BEHAVIORAL HEALTH PROFILE

---

## North Dakota

# North Dakota

## OLDER ADULTS BEHAVIORAL HEALTH PROFILE

August 2016

### NORTH DAKOTA'S POPULATION

#### North Dakota Population by Age Group

North Dakota is home to 739,482 people. Of these:

- 246,452 (33.3 percent) are over age 50.
- 147,462 (19.9 percent) are over age 60.
- 73,416 (9.9 percent) are over age 70.
- 32,755 (4.4 percent) are ages 80 and older.


The proportion of women rises fairly steadily in each age group, and women make up 63.0 percent of the 80+ group. The racial/ethnic composition of older North Dakotans is as follows:

#### Race/Ethnicity of North Dakotans Ages 50+

White	AI/AN	Black	Asian	NH/PI	Other	Hispanic
95.1%	3.0%	0.7%	0.6%	0.0%	0.6%	0.9%

Source: U.S. Census Bureau, 2015  
AI/AN stands for American Indian and Alaska Native.  
NH/PI stands for Native Hawaiian and Other Pacific Islander.


Exhibit 1. North Dakota Population by Age Group, 2014



Source: U.S. Census Bureau, 2015

#### The Number of Older North Dakotans Is Growing

Exhibit 2. North Dakota Population by Age Group, 2000–2030



The proportion of North Dakota's population that is 65 and older is growing while the proportion that is younger than 65 is shrinking. The U.S. Census Bureau estimates that 25.1 percent of North Dakota's population will be 65 and older by the year 2030, an increase of 41.2 percent from 2015.

#### Projected Population in North Dakota

Age Group	2015	2025	2030
Under 18	21.8%	21.6%	21.2%
18 to 44	34.5%	32.1%	31.0%
45 to 64	26.7%	23.4%	22.7%
65+	17.0%	22.9%	25.1%

Source: U.S. Census Bureau, 2005

## SUICIDE AMONG OLDER ADULTS IN REGION 8


### Region 8 Suicide Rates Compared With National Rates

Suicide data for North Dakotans of various ages were unavailable for 2013. Therefore, the rates for Region 8 (including Colorado, Montana, South Dakota, Utah, and Wyoming) are used instead.

The suicide rate among individuals ages 50+ in Region 8 was higher than the rate among younger age groups. In 2013, the total suicide rate among all people ages 50+ was 25.7 per 100,000 people (10.8 for women and 41.9 for men). The rate among those ages 50–64 was higher than the rate in the United States.

States vary in their reporting of suicides. The suicide rate is influenced by these reporting practices.


Exhibit 3. Suicide Rates in Region 8 and the United States, 2013



Source: Centers for Disease Control and Prevention (CDC), National Center for Health Statistics (NCHS), Underlying Cause of Death 1999-2013 on CDC WONDER Online Database, released 2015

### Trends in Suicide Rates in Region 8

Exhibit 4. Trends in Suicide Rates in Region 8 by Age Group, 2004–2013



Source: CDC, NCHS, Underlying Cause of Death 1999-2013 on CDC WONDER Online Database, released 2015

Suicide data for North Dakotans of various ages were unavailable. Therefore, the rates for Region 8 are used instead.

The suicide rate among individuals in Region 8 ages 50+ fluctuated from a low of 20.5 per 100,000 in 2004 to a high of 25.8 per 100,000 in 2012. From 2004 to 2013, the rate was generally highest among those in the 50–64 age group.


How a state reports suicides can vary from year to year. The number of suicides is generally low, so even a small difference in reported numbers may make the rate fluctuate widely.

## SUBSTANCE USE DISORDER AND SUBSTANCE USE DISORDER TREATMENT AMONG OLDER NORTH DAKOTANS

### 30-Day Binge Drinking Among Older North Dakotans

Binge drinking, defined as five or more drinks for men and four or more drinks for women on a single occasion, can lead to serious health problems. Such problems include neurological damage, cardiovascular disease, liver disease, stroke, and poor control of diabetes. Binge drinkers are more likely to take risks such as driving while intoxicated and to experience falls and other accidents. Older people have a lower tolerance for alcohol. Binge drinking decreases with age and occurs more frequently among men than it does among women. As Exhibit 5 shows, 24.0 percent of North Dakota men ages 50–64 reported binge drinking in the past 30 days, while 8.5 percent of those in the 65+ group reported similar behavior.


Exhibit 5. Binge Drinking Rates in North Dakota by Age Group and Sex, 2013



Source: Behavioral Risk Factor Surveillance System (BRFSS), 2013

### Illicit Drug Use Among Older Americans

Exhibit 6. Illicit Drug Use Among Older Americans, 2002–2013



Source: National Survey on Drug Use and Health (NSDUH), 2013  
 Illicit drug use includes illegal drugs and prescription drugs used nonmedically. SAMHSA provides a list of drugs included in its survey in the NSDUH methodological summary.

Nationally, the rate of illicit drug use among older adults ages 50–64 more than doubled from 2002 to 2013. This development partially reflects the entrance into this age group of the baby boom cohort, whose rates of illicit drug use have been higher than those of older cohorts. Although state-specific data are not available, the *North Dakota Behavioral Health Barometer* is available for download from the Substance Abuse and Mental Health Services Administration (SAMHSA) website ([www.samhsa.gov/data/population-data-nsduh/reports?tab=33](http://www.samhsa.gov/data/population-data-nsduh/reports?tab=33)).

# North Dakota


## Admissions to Substance Use Disorder Treatment Among Older North Dakotans

In 2012, there were 286 admissions of North Dakotans ages 50 and older to substance use disorder (SUD) treatment in state-funded treatment programs, a rate of 116.0 per 100,000 people ages 50+. This rate was lower than the regional rate and the national average. Men made up 64.7 percent of these admissions. Of all admissions, 83.9 percent were White/Caucasian, 1.4 percent were Black/African American, and 1.4 percent were Hispanic.

The principal sources of referral to treatment among those ages 50 and older were:

Self-Referral	Criminal Justice	Other
25.9%	36.4%	37.8%


**Exhibit 7. SUD Treatment Admissions of Adults Ages 50+ in North Dakota, Region 8, and the United States by Sex, 2012**



Source: Treatment Episode Data Set (TEDS), 2012<sup>1</sup>  
Data include only those clients reported to SAMHSA.

## SUD Treatment Admissions Among North Dakotans Ages 50+ by Insurance Type

**Exhibit 8. SUD Treatment Admissions of Adults Ages 50+ in North Dakota, Region 8, and the United States by Insurance Type, 2012**



Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

In North Dakota, 57.2 percent of older adult admissions to SUD treatment were uninsured, 10.4 percent had Medicaid, 20.1 percent had Medicare, and 12.2 percent had private insurance.

### *SUD Treatment Admissions Among North Dakotans Ages 50+ by Primary Sources of Payment*

Self-Pay	Medicaid	Other
48.5%	10.6%	40.9%


Self-pay data include private insurance. Exhibit 8 and the accompanying text reflect the type of health insurance (if any) clients had at admission; the table above represents primary sources of payment.

<sup>1</sup> TEDS data are collected by states as a condition of the Substance Abuse Prevention and Treatment Block Grant. Guidelines suggest that states report all clients admitted to publicly financed treatment; however, states are inconsistent in applying the guidelines. States may structure and implement different quality controls over the data. For example, states may collect different categories of information to answer TEDS questions. Information is then "walked over" to TEDS definitions.

## Alcohol Use Disorder Treatment Admissions Among North Dakotans Ages 50+

Alcohol was the most frequently cited substance used by older North Dakotans in publicly financed SUD treatment in 2012. It was mentioned as a substance of use in 85.3 percent of admissions among those ages 50+. This was lower than the regional and national rates.

**Exhibit 9. Alcohol Use Treatment Admissions of Adults Ages 50+ in North Dakota, Region 8, and the United States, 2012**




Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

## SUD Treatment Admissions for Non-Alcohol Substance Use

**Exhibit 10. Non-Alcohol Substance Use Treatment Admissions of Adults Ages 50+ in North Dakota, Region 8, and the United States, 2012**

Substances other than alcohol were cited as the primary substances of use for 14.7 percent of older adult admissions to publicly funded treatment in North Dakota.



Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

## Drug-Related Emergency Department Visits Involving Pharmaceutical Misuse and Abuse by Older Adults

SAMHSA periodically releases reports from the Drug Abuse Warning Network (DAWN). DAWN, discontinued in 2011, consisted of a nationwide network of hospital emergency departments (EDs) primarily located in large metropolitan areas. DAWN data consist of professional reviews of ED records to determine the extent to which alcohol and other substance abuse was involved in ED visits. According to the November 25, 2010, *DAWN Report*:


- *In 2004, there were an estimated 115,803 ED visits involving pharmaceutical misuse and abuse by adults aged 50 or older; in 2008, there were 256,097 such visits, representing an increase of 121.1 percent*
- *One fifth (19.7 percent) of ED visits involving pharmaceutical misuse and abuse among older adults were made by persons aged 70 or older*
- *Among ED visits made by older adults, pain relievers were the type of pharmaceutical most commonly involved (43.5 percent), followed by drugs used to treat anxiety or insomnia (31.8 percent) and antidepressants (8.6 percent)*
- *Among patients aged 50 or older who visited the ED for pharmaceutical misuse or abuse, more than half (52.3 percent) were treated and released, and more than one third (37.5 percent) were admitted to the hospital*

Bulleted text in italics above is taken from SAMHSA's Center for Behavioral Health Statistics and Quality. (2010, November 25). *The DAWN Report: Drug-related emergency department visits involving pharmaceutical misuse and abuse by older adults*. Rockville, MD: SAMHSA.

## CO-OCCURRING SUBSTANCE USE AND MENTAL DISORDERS

### Older North Dakotans in SUD Treatment With Co-Occurring Mental Disorders

**Exhibit 11. SUD Treatment Admissions of Adults Ages 50+ With a Co-Occurring Mental Disorder in North Dakota, Region 8, and the United States, 2012**



Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

The national literature shows a strong relationship between substance use and mental disorders. Studies show that 30–80 percent of individuals with a substance use or mental disorder also have a co-occurring disorder.


Exhibit 11 shows the proportion of SUD treatment admissions of North Dakotans ages 50+ with a co-occurring mental disorder. This rate is higher than the regional rate and the national average. However, state reporting practices are a factor in these results.

## MENTAL HEALTH

### Older North Dakotans Reporting Frequent Mental Distress

BRFSS, a household survey conducted in all 50 states and several territories, asks the following question: "Now thinking about your mental health, which includes stress, depression, and problems with emotions, for how many days during the past 30 days was your mental health not good?" Those individuals reporting 14 or more "Yes" days in response to this question experience frequent mental distress (FMD). Exhibit 12 shows that older North Dakotans experience FMD at a rate that is lower than the regional and national rates.


**Exhibit 12. Individuals Reporting Frequent Mental Distress in North Dakota, Region 8, and the United States, 2013**



Source: BRFSS, 2013

### Older North Dakotans Reporting Frequent Mental Distress by Age Group and Sex

**Exhibit 13. North Dakotans Reporting Frequent Mental Distress by Age Group and Sex, 2013**



Source: BRFSS, 2013

Older men in North Dakota were more likely to indulge in binge drinking, but older women were more likely to report that they had FMD (14 days or more per 30-day period). As Exhibit 13 shows, 10.1 percent of women in the 50–64 age group and 4.3 percent in the 65+ age group reported FMD, while 9.8 percent of men in the 50–64 age group and 5.2 percent in the 65+ age group reported FMD.



## Other Measures of Mental Health

BRFSS collected other measures showing risk factors for mental and/or physical illness. These included:

- Social and Emotional Support (2010). BRFSS asked, "How often do you get the social and emotional support you need?" The possible responses were always, usually, sometimes, rarely, or never.
- Life Satisfaction (2010). BRFSS asked, "In general, how satisfied are you with your life?" The possible responses were very satisfied, satisfied, dissatisfied, or very dissatisfied.

Exhibit 14 presents the results of these surveys among older North Dakotans.

**Exhibit 14. BRFSS Measures, 2010**


Indicator	Ages 50+	Ages 50–64	Ages 65+
Rarely or never get social or emotional support	8.3%	5.3%	12.0%
Dissatisfied or very dissatisfied	3.6%	4.5%	2.5%

Source: BRFSS, 2010

## Individuals With Frequent Mental Distress Report High Rates of Poor Physical Health

Older Americans who experienced FMD were more likely to report that their physical health was poor (14 days or more in the past 30-day period when physical health was "not good"). As shown in Exhibit 15, although nearly 11 percent of older Americans with no mental distress reported poor physical health, more than 50 percent of those with FMD reported poor physical health.

**Exhibit 15. Individuals Ages 50+ in the United States Reporting Poor Physical Health by Level of Mental Distress, 2013**




Source: BRFSS, 2013

## Relationship Among Mental Distress, Diabetes, Stroke, Heart Attack, High Blood Pressure, and Coronary Disease

Older Americans who experience FMD are more likely to report that they have health problems. People with FMD were more than twice as likely to report having a stroke than those with some or no mental distress. They experienced coronary disease and heart attack at more than 1.6 times, diabetes at more than 1.4 times, and high blood pressure at nearly 1.2 times the rate of those with some or no mental distress.

**Exhibit 16. Individuals Ages 50+ in the United States With Mental Distress and Serious Health Problems, 2013**



Source: BRFSS, 2013

## Older North Dakotans Admitted to State Mental Health Services

Approximately 5.8 percent of the people served by the North Dakota mental health system were ages 65 and older. This represents more than 990 adults.

Source: Center for Mental Health Services (CMHS) Uniform Reporting System (URS) Output Tables, 2014

## DATA SOURCES

**BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM** ([www.cdc.gov/brfss](http://www.cdc.gov/brfss)). BRFSS, managed by CDC, is the largest ongoing telephone health survey system in the world, tracking health conditions and risk behaviors in the United States annually since 1984. Data are collected on a monthly basis in all 50 states, the District of Columbia, and participating territories. BRFSS data are collected by local jurisdictions and reported to CDC.

**CENTERS FOR DISEASE CONTROL AND PREVENTION, NATIONAL CENTER FOR HEALTH STATISTICS, UNDERLYING CAUSE OF DEATH 1999-2013 ON CDC WONDER ONLINE DATABASE, RELEASED 2015** (<http://wonder.cdc.gov/ucd-icd10.html>). The WONDER online database "contains mortality and population counts for all U.S. counties. Data are based on death certificates for U.S. residents. Each death certificate identifies a single underlying cause of death and demographic data. The number of deaths, crude death rates or age-adjusted death rates, and 95% confidence intervals and standard errors for death rates can be obtained by place of residence (total U.S., region, state and county), age group (single-year-of age, 5-year age groups, 10-year age groups and infant age groups), race, Hispanic ethnicity, gender, year, cause-of-death (4-digit ICD-10 code or group of codes), injury intent and injury mechanism, drug/alcohol induced causes and urbanization categories. Data are also available for place of death, month and week day of death, and whether an autopsy was performed."

**CENTER FOR MENTAL HEALTH SERVICES UNIFORM REPORTING SYSTEM** ([www.samhsa.gov/data/us\\_map](http://www.samhsa.gov/data/us_map)). States that receive CMHS Block Grants are required to report aggregate data to URS. URS reports include information about utilization of mental health services and client demographic and outcome information.

**NATIONAL SURVEY ON DRUG USE AND HEALTH** (<https://nsduhweb.rti.org>). The NSDUH, managed by SAMHSA, is "an annual nationwide survey involving interviews with approximately 70,000 randomly selected individuals aged 12 and older." NSDUH data are most frequently used by state planners to assess the need for SUD treatment. NSDUH data also include information about mental health conditions.

**TREATMENT EPISODE DATA SET** ([www.samhsa.gov/data/client-level-data-teds/reports?tab=19](http://www.samhsa.gov/data/client-level-data-teds/reports?tab=19)). States that receive Substance Abuse Prevention and Treatment Block Grant funds submit individual client data to TEDS. TEDS includes both admission and discharge data sets, and some 1.5 million admissions are reported annually. TEDS includes information about utilization of SUD treatment services and client demographic and outcome information. TEDS is an admission-based system, and TEDS admissions do not represent individuals. Thus, an individual admitted to treatment twice within a calendar year would be counted as two admissions.

**U.S. CENSUS BUREAU** ([www.census.gov/people](http://www.census.gov/people)). Two main sources of Census Bureau data were used in this report: (1) population estimates and (2) population projections.

# OLDER ADULTS BEHAVIORAL HEALTH PROFILE

---

## South Dakota

# South Dakota

## OLDER ADULTS BEHAVIORAL HEALTH PROFILE

August 2016

### SOUTH DAKOTA'S POPULATION

#### South Dakota Population by Age Group

South Dakota is home to 853,175 people. Of these:

- 300,994 (35.3 percent) are over age 50.
- 183,250 (21.5 percent) are over age 60.
- 89,750 (10.5 percent) are over age 70.
- 38,666 (4.5 percent) are ages 80 and older.


The proportion of women rises fairly steadily in each age group, and women make up 62.6 percent of the 80+ group. The racial/ethnic composition of older South Dakotans is as follows:

#### Race/Ethnicity of South Dakotans Ages 50+

White	AI/AN	Black	Asian	NH/PI	Other	Hispanic
93.2%	4.6%	0.8%	0.6%	0.0%	0.7%	1.3%

Source: U.S. Census Bureau, 2015  
AI/AN stands for American Indian and Alaska Native.  
NH/PI stands for Native Hawaiian and Other Pacific Islander.


Exhibit 1. South Dakota Population by Age Group, 2014



Source: U.S. Census Bureau, 2015

#### The Number of Older South Dakotans Is Growing

Exhibit 2. South Dakota Population by Age Group, 2000–2030



The proportion of South Dakota's population that is 65 and older is growing while the proportion that is younger than 65 is shrinking. The U.S. Census Bureau estimates that 23.1 percent of South Dakota's population will be 65 and older by the year 2030, an increase of 44.6 percent from 2015.

#### Projected Population in South Dakota

Age Group	2015	2025	2030
Under 18	24.6%	24.5%	24.5%
18 to 44	33.1%	30.9%	29.5%
45 to 64	26.2%	23.5%	22.9%
65+	16.1%	21.2%	23.1%

Source: U.S. Census Bureau, 2005

## SUICIDE AMONG OLDER SOUTH DAKOTANS


### Region 8 Suicide Rates Compared With National Rates

Suicide data for South Dakotans of various ages were unavailable for 2013. Therefore, the rates for Region 8 (including Colorado, Montana, North Dakota, Utah, and Wyoming) are used instead.

The suicide rate among individuals ages 50+ in Region 8 was higher than the rate among younger age groups. In 2013, the total suicide rate among all people ages 50+ was 25.7 per 100,000 people (10.8 for women and 41.9 for men). The rate among those ages 50–64 was higher than the rate in the United States.

States vary in their reporting of suicides. The suicide rate is influenced by these reporting practices.


Exhibit 3. Suicide Rates in Region 8 and the United States, 2013



Source: Centers for Disease Control and Prevention (CDC), National Center for Health Statistics (NCHS), Underlying Cause of Death 1999-2013 on CDC WONDER Online Database, released 2015

### Trends in Suicide Rates in Region 8

Exhibit 4. Trends in Suicide Rates in Region 8 by Age Group, 2004–2013



Source: CDC, NCHS, Underlying Cause of Death 1999-2013 on CDC WONDER Online Database, released 2015

Suicide data for South Dakotans of various ages were unavailable. Therefore, the rates for Region 8 are used instead.

The suicide rate among individuals in Region 8 ages 50+ fluctuated from a low of 20.5 per 100,000 in 2004 to a high of 25.8 per 100,000 in 2012. From 2004 to 2013, the rate was generally highest among those in the 50–64 age group.


How a state reports suicides can vary from year to year. The number of suicides is generally low, so even a small difference in reported numbers may make the rate fluctuate widely.

## SUBSTANCE USE DISORDER AND SUBSTANCE USE DISORDER TREATMENT AMONG OLDER SOUTH DAKOTANS

### 30-Day Binge Drinking Among Older South Dakotans

Binge drinking, defined as five or more drinks for men and four or more drinks for women on a single occasion, can lead to serious health problems. Such problems include neurological damage, cardiovascular disease, liver disease, stroke, and poor control of diabetes. Binge drinkers are more likely to take risks such as driving while intoxicated and to experience falls and other accidents. Older people have a lower tolerance for alcohol. Binge drinking decreases with age and occurs more frequently among men than it does among women. As Exhibit 5 shows, 17.0 percent of South Dakota men ages 50–64 reported binge drinking in the past 30 days, while 4.2 percent of those in the 65+ group reported similar behavior.


Exhibit 5. Binge Drinking Rates in South Dakota by Age Group and Sex, 2013



Source: Behavioral Risk Factor Surveillance System (BRFSS), 2013

### Illicit Drug Use Among Older Americans

Exhibit 6. Illicit Drug Use Among Older Americans, 2002–2013



Source: National Survey on Drug Use and Health (NSDUH), 2013  
 Illicit drug use includes illegal drugs and prescription drugs used nonmedically. SAMHSA provides a list of drugs included in its survey in the NSDUH methodological summary.

Nationally, the rate of illicit drug use among older adults ages 50–64 more than doubled from 2002 to 2013. This development partially reflects the entrance into this age group of the baby boom cohort, whose rates of illicit drug use have been higher than those of older cohorts. Although state-specific data are not available, the *South Dakota Behavioral Health Barometer* is available for download from the Substance Abuse and Mental Health Services Administration (SAMHSA) website ([www.samhsa.gov/data/population-data-nsduh/reports?tab=33](http://www.samhsa.gov/data/population-data-nsduh/reports?tab=33)).

# South Dakota


## Admissions to Substance Use Disorder Treatment Among Older South Dakotans

In 2012, there were 2,249 admissions of South Dakotans ages 50 and older to substance use disorder (SUD) treatment in state-funded treatment programs, a rate of 747.2 per 100,000 people ages 50+. This rate was higher than the regional rate and the national average. Men made up 78.0 percent of these admissions. Of all admissions, 51.7 percent were White/Caucasian, 1.5 percent were Black/African American, and 2.8 percent were Hispanic.

The principal sources of referral to treatment among those ages 50 and older were:

Self-Referral	Criminal Justice	Other
16.2%	56.3%	27.5%


**Exhibit 7. SUD Treatment Admissions of Adults Ages 50+ in South Dakota, Region 8, and the United States by Sex, 2012**



Source: Treatment Episode Data Set (TEDS), 2012<sup>1</sup>  
Data include only those clients reported to SAMHSA.

## SUD Treatment Admissions Among South Dakotans Ages 50+ by Insurance Type

**Exhibit 8. SUD Treatment Admissions of Adults Ages 50+ in South Dakota, Region 8, and the United States by Insurance Type, 2012**



Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

In South Dakota, 49.9 percent of older adult admissions to SUD treatment were uninsured, 11.4 percent had Medicaid, 29.6 percent had Medicare, and 9.1 percent had private insurance.

### *SUD Treatment Admissions Among South Dakotans Ages 50+ by Primary Sources of Payment*

Self-Pay	Medicaid	Other
17.2%	0.2%	82.6%

Self-pay data include private insurance. Exhibit 8 and the accompanying text reflect the type of health insurance (if any) clients had at admission; the table above represents primary sources of payment.


<sup>1</sup> TEDS data are collected by states as a condition of the Substance Abuse Prevention and Treatment Block Grant. Guidelines suggest that states report all clients admitted to publicly financed treatment; however, states are inconsistent in applying the guidelines. States may structure and implement different quality controls over the data. For example, states may collect different categories of information to answer TEDS questions. Information is then "walked over" to TEDS definitions.



## Alcohol Use Disorder Treatment Admissions Among South Dakotans Ages 50+

Alcohol was the most frequently cited substance used by older South Dakotans in publicly financed SUD treatment in 2012. It was mentioned as a substance of use in 93.4 percent of admissions among those ages 50+. This was higher than the regional and national rates.

**Exhibit 9. Alcohol Use Treatment Admissions of Adults Ages 50+ in South Dakota, Region 8, and the United States, 2012**




Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

## SUD Treatment Admissions for Non-Alcohol Substance Use

**Exhibit 10. Non-Alcohol Substance Use Treatment Admissions of Adults Ages 50+ in South Dakota, Region 8, and the United States, 2012**

Substances other than alcohol were cited as the primary substances of use for 6.6 percent of older adult admissions to publicly funded treatment in South Dakota.



Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

## Drug-Related Emergency Department Visits Involving Pharmaceutical Misuse and Abuse by Older Adults

SAMHSA periodically releases reports from the Drug Abuse Warning Network (DAWN). DAWN, discontinued in 2011, consisted of a nationwide network of hospital emergency departments (EDs) primarily located in large metropolitan areas. DAWN data consist of professional reviews of ED records to determine the extent to which alcohol and other substance abuse was involved in ED visits. According to the November 25, 2010, *DAWN Report*:


- *In 2004, there were an estimated 115,803 ED visits involving pharmaceutical misuse and abuse by adults aged 50 or older; in 2008, there were 256,097 such visits, representing an increase of 121.1 percent*
- *One fifth (19.7 percent) of ED visits involving pharmaceutical misuse and abuse among older adults were made by persons aged 70 or older*
- *Among ED visits made by older adults, pain relievers were the type of pharmaceutical most commonly involved (43.5 percent), followed by drugs used to treat anxiety or insomnia (31.8 percent) and antidepressants (8.6 percent)*
- *Among patients aged 50 or older who visited the ED for pharmaceutical misuse or abuse, more than half (52.3 percent) were treated and released, and more than one third (37.5 percent) were admitted to the hospital*

Bulleted text in italics above is taken from SAMHSA's Center for Behavioral Health Statistics and Quality. (2010, November 25). *The DAWN Report: Drug-related emergency department visits involving pharmaceutical misuse and abuse by older adults*. Rockville, MD: SAMHSA.

## CO-OCCURRING SUBSTANCE USE AND MENTAL DISORDERS

### Older South Dakotans in SUD Treatment With Co-Occurring Mental Disorders

**Exhibit 11. SUD Treatment Admissions of Adults Ages 50+ With a Co-Occurring Mental Disorder in South Dakota, Region 8, and the United States, 2012**



Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

The national literature shows a strong relationship between substance use and mental disorders. Studies show that 30–80 percent of individuals with a substance use or mental disorder also have a co-occurring disorder.


Exhibit 11 shows the proportion of SUD treatment admissions of South Dakotans ages 50+ with a co-occurring mental disorder. This rate is higher than the regional rate and the national average. However, state reporting practices are a factor in these results.

## MENTAL HEALTH

### Older South Dakotans Reporting Frequent Mental Distress

BRFSS, a household survey conducted in all 50 states and several territories, asks the following question: "Now thinking about your mental health, which includes stress, depression, and problems with emotions, for how many days during the past 30 days was your mental health not good?" Those individuals reporting 14 or more "Yes" days in response to this question experience frequent mental distress (FMD). Exhibit 12 shows that older South Dakotans experience FMD at a rate that is lower than the regional and national rates.


**Exhibit 12. Individuals Reporting Frequent Mental Distress in South Dakota, Region 8, and the United States, 2013**



Source: BRFSS, 2013

### Older South Dakotans Reporting Frequent Mental Distress by Age Group and Sex

**Exhibit 13. South Dakotans Reporting Frequent Mental Distress by Age Group and Sex, 2013**



Source: BRFSS, 2013

Older men in South Dakota were more likely to indulge in binge drinking, but older women were more likely to report that they had FMD (14 days or more per 30-day period). As Exhibit 13 shows, 10.2 percent of women in the 50–64 age group and 6.0 percent in the 65+ age group reported FMD, while 4.6 percent of men in the 50–64 age group and 3.6 percent in the 65+ age group reported FMD.

## Other Measures of Mental Health

BRFSS collected other measures showing risk factors for mental and/or physical illness. These included:

- Social and Emotional Support (2010). BRFSS asked, "How often do you get the social and emotional support you need?" The possible responses were always, usually, sometimes, rarely, or never.
- Life Satisfaction (2010). BRFSS asked, "In general, how satisfied are you with your life?" The possible responses were very satisfied, satisfied, dissatisfied, or very dissatisfied.

Exhibit 14 presents the results of these surveys among older South Dakotans.

**Exhibit 14. BRFSS Measures, 2010**


Indicator	Ages 50+	Ages 50–64	Ages 65+
Rarely or never get social or emotional support	8.1%	5.7%	11.4%
Dissatisfied or very dissatisfied	3.2%	3.3%	3.0%

Source: BRFSS, 2010

## Individuals With Frequent Mental Distress Report High Rates of Poor Physical Health

Older Americans who experienced FMD were more likely to report that their physical health was poor (14 days or more in the past 30-day period when physical health was "not good"). As shown in Exhibit 15, although nearly 11 percent of older Americans with no mental distress reported poor physical health, more than 50 percent of those with FMD reported poor physical health.

**Exhibit 15. Individuals Ages 50+ in the United States Reporting Poor Physical Health by Level of Mental Distress, 2013**




Source: BRFSS, 2013

## Relationship Among Mental Distress, Diabetes, Stroke, Heart Attack, High Blood Pressure, and Coronary Disease

Older Americans who experience FMD are more likely to report that they have health problems. People with FMD were more than twice as likely to report having a stroke than those with some or no mental distress. They experienced coronary disease and heart attack at more than 1.6 times, diabetes at more than 1.4 times, and high blood pressure at nearly 1.2 times the rate of those with some or no mental distress.

**Exhibit 16. Individuals Ages 50+ in the United States With Mental Distress and Serious Health Problems, 2013**



Source: BRFSS, 2013

## Older South Dakotans Admitted to State Mental Health Services

Approximately 5.6 percent of the people served by the South Dakota mental health system were ages 65 and older. This represents more than 850 adults.

Source: Center for Mental Health Services (CMHS) Uniform Reporting System (URS) Output Tables, 2014

## DATA SOURCES

**BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM** ([www.cdc.gov/brfss](http://www.cdc.gov/brfss)). BRFSS, managed by CDC, is the largest ongoing telephone health survey system in the world, tracking health conditions and risk behaviors in the United States annually since 1984. Data are collected on a monthly basis in all 50 states, the District of Columbia, and participating territories. BRFSS data are collected by local jurisdictions and reported to CDC.

**CENTERS FOR DISEASE CONTROL AND PREVENTION, NATIONAL CENTER FOR HEALTH STATISTICS, UNDERLYING CAUSE OF DEATH 1999-2013 ON CDC WONDER ONLINE DATABASE, RELEASED 2015** (<http://wonder.cdc.gov/ucd-icd10.html>). The WONDER online database "contains mortality and population counts for all U.S. counties. Data are based on death certificates for U.S. residents. Each death certificate identifies a single underlying cause of death and demographic data. The number of deaths, crude death rates or age-adjusted death rates, and 95% confidence intervals and standard errors for death rates can be obtained by place of residence (total U.S., region, state and county), age group (single-year-of age, 5-year age groups, 10-year age groups and infant age groups), race, Hispanic ethnicity, gender, year, cause-of-death (4-digit ICD-10 code or group of codes), injury intent and injury mechanism, drug/alcohol induced causes and urbanization categories. Data are also available for place of death, month and week day of death, and whether an autopsy was performed."

**CENTER FOR MENTAL HEALTH SERVICES UNIFORM REPORTING SYSTEM** ([www.samhsa.gov/data/us\\_map](http://www.samhsa.gov/data/us_map)). States that receive CMHS Block Grants are required to report aggregate data to URS. URS reports include information about utilization of mental health services and client demographic and outcome information.

**NATIONAL SURVEY ON DRUG USE AND HEALTH** (<https://nsduhweb.rti.org>). The NSDUH, managed by SAMHSA, is "an annual nationwide survey involving interviews with approximately 70,000 randomly selected individuals aged 12 and older." NSDUH data are most frequently used by state planners to assess the need for SUD treatment. NSDUH data also include information about mental health conditions.

**TREATMENT EPISODE DATA SET** ([www.samhsa.gov/data/client-level-data-teds/reports?tab=19](http://www.samhsa.gov/data/client-level-data-teds/reports?tab=19)). States that receive Substance Abuse Prevention and Treatment Block Grant funds submit individual client data to TEDS. TEDS includes both admission and discharge data sets, and some 1.5 million admissions are reported annually. TEDS includes information about utilization of SUD treatment services and client demographic and outcome information. TEDS is an admission-based system, and TEDS admissions do not represent individuals. Thus, an individual admitted to treatment twice within a calendar year would be counted as two admissions.

**U.S. CENSUS BUREAU** ([www.census.gov/people](http://www.census.gov/people)). Two main sources of Census Bureau data were used in this report: (1) population estimates and (2) population projections.

# OLDER ADULTS BEHAVIORAL HEALTH PROFILE

---

Utah

### UTAH'S POPULATION

#### Utah Population by Age Group

Utah is home to 2,942,902 people. Of these:

- 727,541 (24.7 percent) are over age 50.
- 422,657 (14.4 percent) are over age 60.
- 195,595 (6.6 percent) are over age 70.
- 71,444 (2.4 percent) are ages 80 and older.


The proportion of women rises fairly steadily in each age group, and women make up 57.9 percent of the 80+ group. The racial/ethnic composition of older Utahns is as follows:

#### Race/Ethnicity of Utahns Ages 50+

White	AI/AN	Black	Asian	NH/PI	Other	Hispanic
94.6%	1.1%	0.8%	2.1%	0.6%	0.7%	7.3%

Source: U.S. Census Bureau, 2015  
AI/AN stands for American Indian and Alaska Native.  
NH/PI stands for Native Hawaiian and Other Pacific Islander.


Exhibit 1. Utah Population by Age Group, 2014



Source: U.S. Census Bureau, 2015

#### The Number of Older Utahns Is Growing

Exhibit 2. Utah Population by Age Group, 2000–2030



The proportion of Utah's population that is 65 and older is growing while the proportion that is younger than 65 is shrinking. The U.S. Census Bureau estimates that 13.2 percent of Utah's population will be 65 and older by the year 2030, an increase of 63.9 percent from 2015.

#### Projected Population in Utah

Age Group	2015	2025	2030
Under 18	31.3%	30.4%	30.4%
18 to 44	38.6%	37.0%	36.9%
45 to 64	20.0%	20.0%	19.5%
65+	10.1%	12.6%	13.2%

Source: U.S. Census Bureau, 2005




## SUICIDE AMONG OLDER UTAHNS

### Utah Suicide Rate Compared With Regional and National Rates

The suicide rate among Utahns ages 50 and older is higher than the rate among younger age groups. In 2013, the total suicide rate among all people ages 50+ was 29.9 per 100,000 people (13.0 for women and 48.2 for men). The rate among those ages 50–64 was higher than both the rate in the region (including Colorado, Montana, North Dakota, South Dakota, and Wyoming) and the rate in the United States.

States vary in their reporting of suicides. The suicide rate is influenced by these reporting practices.


Exhibit 3. Suicide Rates in Utah, Region 8, and the United States, 2013



Source: Centers for Disease Control and Prevention (CDC), National Center for Health Statistics (NCHS), Underlying Cause of Death 1999-2013 on CDC WONDER Online Database, released 2015

### Trends in Suicide Rates in Utah

Exhibit 4. Trends in Suicide Rates in Utah by Age Group, 2004–2013



Source: CDC, NCHS, Underlying Cause of Death 1999-2013 on CDC WONDER Online Database, released 2015

The suicide rate among Utahns ages 50+ fluctuated from a low of 18.3 per 100,000 in 2005 to a high of 29.9 per 100,000 in 2013. From 2004 to 2013, the rate was highest among those in the 50–64 age group.


How a state reports suicides can vary from year to year. The number of suicides is generally low, so even a small difference in reported numbers may make the rate fluctuate widely.

## SUBSTANCE USE DISORDER AND SUBSTANCE USE DISORDER TREATMENT AMONG OLDER UTAHNS

### 30-Day Binge Drinking Among Older Utahns

Binge drinking, defined as five or more drinks for men and four or more drinks for women on a single occasion, can lead to serious health problems. Such problems include neurological damage, cardiovascular disease, liver disease, stroke, and poor control of diabetes. Binge drinkers are more likely to take risks such as driving while intoxicated and to experience falls and other accidents. Older people have a lower tolerance for alcohol. Binge drinking decreases with age and occurs more frequently among men than it does among women. As Exhibit 5 shows, 12.1 percent of Utah men ages 50–64 reported binge drinking in the past 30 days, while 4.4 percent of those in the 65+ group reported similar behavior.


Exhibit 5. Binge Drinking Rates in Utah by Age Group and Sex, 2013



Source: Behavioral Risk Factor Surveillance System (BRFSS), 2013

### Illicit Drug Use Among Older Americans

Exhibit 6. Illicit Drug Use Among Older Americans, 2002–2013



Source: National Survey on Drug Use and Health (NSDUH), 2013  
 Illicit drug use includes illegal drugs and prescription drugs used nonmedically. SAMHSA provides a list of drugs included in its survey in the NSDUH methodological summary.

Nationally, the rate of illicit drug use among older adults ages 50–64 more than doubled from 2002 to 2013. This development partially reflects the entrance into this age group of the baby boom cohort, whose rates of illicit drug use have been higher than those of older cohorts. Although state-specific data are not available, the *Utah Behavioral Health Barometer* is available for download from the Substance Abuse and Mental Health Services Administration (SAMHSA) website ([www.samhsa.gov/data/population-data-nsduh/reports?tab=33](http://www.samhsa.gov/data/population-data-nsduh/reports?tab=33)).


## Admissions to Substance Use Disorder Treatment Among Older Utahns

In 2012, there were 1,586 admissions of Utahns ages 50 and older to substance use disorder (SUD) treatment in state-funded treatment programs, a rate of 218 per 100,000 people ages 50+. This rate was lower than the regional rate and the national average. Men made up 82.7 percent of these admissions. Of all admissions, 67.7 percent were White/Caucasian, 7.1 percent were Black/African American, and 13.3 percent were Hispanic.

The principal sources of referral to treatment among those ages 50 and older were:

Self-Referral	Criminal Justice	Other
36.8%	40.8%	22.3%


**Exhibit 7. SUD Treatment Admissions of Adults Ages 50+ in Utah, Region 8, and the United States by Sex, 2012**



Source: Treatment Episode Data Set (TEDS), 2012<sup>1</sup>  
Data include only those clients reported to SAMHSA.

## SUD Treatment Admissions Among Utahns Ages 50+ by Insurance Type

**Exhibit 8. SUD Treatment Admissions of Adults Ages 50+ in Utah, Region 8, and the United States by Insurance Type, 2012**



Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

In Utah, 77.4 percent of older adult admissions to SUD treatment were uninsured, 11.1 percent had Medicaid, 7.2 percent had Medicare, and 4.3 percent had private insurance.

### *SUD Treatment Admissions Among Utahns Ages 50+ by Primary Sources of Payment*

Self-Pay	Medicaid	Other
6.1%	10.1%	83.7%


Self-pay data include private insurance. Exhibit 8 and the accompanying text reflect the type of health insurance (if any) clients had at admission; the table above represents primary sources of payment.

<sup>1</sup> TEDS data are collected by states as a condition of the Substance Abuse Prevention and Treatment Block Grant. Guidelines suggest that states report all clients admitted to publicly financed treatment; however, states are inconsistent in applying the guidelines. States may structure and implement different quality controls over the data. For example, states may collect different categories of information to answer TEDS questions. Information is then "walked over" to TEDS definitions.

## Alcohol Use Disorder Treatment Admissions Among Utahns Ages 50+

Alcohol was the most frequently cited substance used by older Utahns in publicly financed SUD treatment in 2012. It was mentioned as a substance of use in 62.3 percent of admissions among those ages 50+. This was lower than the regional rate and higher than the national rate.

**Exhibit 9. Alcohol Use Treatment Admissions of Adults Ages 50+ in Utah, Region 8, and the United States, 2012**




Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

## SUD Treatment Admissions for Non-Alcohol Substance Use

**Exhibit 10. Non-Alcohol Substance Use Treatment Admissions of Adults Ages 50+ in Utah, Region 8, and the United States, 2012**

Substances other than alcohol were cited as the primary substances of use for 37.7 percent of older adult admissions to publicly funded treatment in Utah.



Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

## Drug-Related Emergency Department Visits Involving Pharmaceutical Misuse and Abuse by Older Adults

SAMHSA periodically releases reports from the Drug Abuse Warning Network (DAWN). DAWN, discontinued in 2011, consisted of a nationwide network of hospital emergency departments (EDs) primarily located in large metropolitan areas. DAWN data consist of professional reviews of ED records to determine the extent to which alcohol and other substance abuse was involved in ED visits. According to the November 25, 2010, *DAWN Report*:


- *In 2004, there were an estimated 115,803 ED visits involving pharmaceutical misuse and abuse by adults aged 50 or older; in 2008, there were 256,097 such visits, representing an increase of 121.1 percent*
- *One fifth (19.7 percent) of ED visits involving pharmaceutical misuse and abuse among older adults were made by persons aged 70 or older*
- *Among ED visits made by older adults, pain relievers were the type of pharmaceutical most commonly involved (43.5 percent), followed by drugs used to treat anxiety or insomnia (31.8 percent) and antidepressants (8.6 percent)*
- *Among patients aged 50 or older who visited the ED for pharmaceutical misuse or abuse, more than half (52.3 percent) were treated and released, and more than one third (37.5 percent) were admitted to the hospital*

Bulleted text in italics above is taken from SAMHSA's Center for Behavioral Health Statistics and Quality. (2010, November 25). *The DAWN Report: Drug-related emergency department visits involving pharmaceutical misuse and abuse by older adults*. Rockville, MD: SAMHSA.

## CO-OCCURRING SUBSTANCE USE AND MENTAL DISORDERS

### Older Utahns in SUD Treatment With Co-Occurring Mental Disorders

**Exhibit 11. SUD Treatment Admissions of Adults Ages 50+ With a Co-Occurring Mental Disorder in Utah, Region 8, and the United States, 2012**



Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

The national literature shows a strong relationship between substance use and mental disorders. Studies show that 30–80 percent of individuals with a substance use or mental disorder also have a co-occurring disorder.


Exhibit 11 shows the proportion of SUD treatment admissions of Utahns ages 50+ with a co-occurring mental disorder. This rate is lower than the regional rate and the national average. However, state reporting practices are a factor in these results.

## MENTAL HEALTH

### Older Utahns Reporting Frequent Mental Distress

BRFSS, a household survey conducted in all 50 states and several territories, asks the following question: "Now thinking about your mental health, which includes stress, depression, and problems with emotions, for how many days during the past 30 days was your mental health not good?" Those individuals reporting 14 or more "Yes" days in response to this question experience frequent mental distress (FMD). Exhibit 12 shows that older Utahns experience FMD at a rate that is higher than the regional rate and lower than the national rates.


**Exhibit 12. Individuals Reporting Frequent Mental Distress in Utah, Region 8, and the United States, 2013**



Source: BRFSS, 2013

### Older Utahns Reporting Frequent Mental Distress by Age Group and Sex

**Exhibit 13. Utahns Reporting Frequent Mental Distress by Age Group and Sex, 2013**



Source: BRFSS, 2013

Older men in Utah were more likely to indulge in binge drinking, but older women were more likely to report that they had FMD (14 days or more per 30-day period). As Exhibit 13 shows, 14.2 percent of women in the 50–64 age group and 8.7 percent in the 65+ age group reported FMD, while 9.2 percent of men in the 50–64 age group and 3.7 percent in the 65+ age group reported FMD.

## Other Measures of Mental Health

BRFSS collected other measures showing risk factors for mental and/or physical illness. These included:

- Social and Emotional Support (2010). BRFSS asked, "How often do you get the social and emotional support you need?" The possible responses were always, usually, sometimes, rarely, or never.
- Life Satisfaction (2010). BRFSS asked, "In general, how satisfied are you with your life?" The possible responses were very satisfied, satisfied, dissatisfied, or very dissatisfied.

Exhibit 14 presents the results of these surveys among older Utahns.

**Exhibit 14. BRFSS Measures, 2010**


Indicator	Ages 50+	Ages 50–64	Ages 65+
Rarely or never get social or emotional support	8.2%	6.9%	10.2%
Dissatisfied or very dissatisfied	4.2%	4.8%	3.2%

Source: BRFSS, 2010

## Individuals With Frequent Mental Distress Report High Rates of Poor Physical Health

Older Americans who experienced FMD were more likely to report that their physical health was poor (14 days or more in the past 30-day period when physical health was "not good"). As shown in Exhibit 15, although nearly 11 percent of older Americans with no mental distress reported poor physical health, more than 50 percent of those with FMD reported poor physical health.

**Exhibit 15. Individuals Ages 50+ in the United States Reporting Poor Physical Health by Level of Mental Distress, 2013**




Source: BRFSS, 2013

## Relationship Among Mental Distress, Diabetes, Stroke, Heart Attack, High Blood Pressure, and Coronary Disease

Older Americans who experience FMD are more likely to report that they have health problems. People with FMD were more than twice as likely to report having a stroke than those with some or no mental distress. They experienced coronary disease and heart attack at more than 1.6 times, diabetes at more than 1.4 times, and high blood pressure at nearly 1.2 times the rate of those with some or no mental distress.

**Exhibit 16. Individuals Ages 50+ in the United States With Mental Distress and Serious Health Problems, 2013**



Source: BRFSS, 2013

## Older Utahns Admitted to State Mental Health Services

Approximately 4.1 percent of the people served by the Utah mental health system were ages 65 and older. This represents more than 2,000 adults.

Source: Center for Mental Health Services (CMHS) Uniform Reporting System (URS) Output Tables, 2014



## DATA SOURCES

**BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM** ([www.cdc.gov/brfss](http://www.cdc.gov/brfss)). BRFSS, managed by CDC, is the largest ongoing telephone health survey system in the world, tracking health conditions and risk behaviors in the United States annually since 1984. Data are collected on a monthly basis in all 50 states, the District of Columbia, and participating territories. BRFSS data are collected by local jurisdictions and reported to CDC.

**CENTERS FOR DISEASE CONTROL AND PREVENTION, NATIONAL CENTER FOR HEALTH STATISTICS, UNDERLYING CAUSE OF DEATH 1999-2013 ON CDC WONDER ONLINE DATABASE, RELEASED 2015** (<http://wonder.cdc.gov/ucd-icd10.html>). The WONDER online database “contains mortality and population counts for all U.S. counties. Data are based on death certificates for U.S. residents. Each death certificate identifies a single underlying cause of death and demographic data. The number of deaths, crude death rates or age-adjusted death rates, and 95% confidence intervals and standard errors for death rates can be obtained by place of residence (total U.S., region, state and county), age group (single-year-of age, 5-year age groups, 10-year age groups and infant age groups), race, Hispanic ethnicity, gender, year, cause-of-death (4-digit ICD-10 code or group of codes), injury intent and injury mechanism, drug/alcohol induced causes and urbanization categories. Data are also available for place of death, month and week day of death, and whether an autopsy was performed.”

**CENTER FOR MENTAL HEALTH SERVICES UNIFORM REPORTING SYSTEM** ([www.samhsa.gov/data/us\\_map](http://www.samhsa.gov/data/us_map)). States that receive CMHS Block Grants are required to report aggregate data to URS. URS reports include information about utilization of mental health services and client demographic and outcome information.

**NATIONAL SURVEY ON DRUG USE AND HEALTH** (<https://nsduhweb.rti.org>). The NSDUH, managed by SAMHSA, is “an annual nationwide survey involving interviews with approximately 70,000 randomly selected individuals aged 12 and older.” NSDUH data are most frequently used by state planners to assess the need for SUD treatment. NSDUH data also include information about mental health conditions.

**TREATMENT EPISODE DATA SET** ([www.samhsa.gov/data/client-level-data-teds/reports?tab=19](http://www.samhsa.gov/data/client-level-data-teds/reports?tab=19)). States that receive Substance Abuse Prevention and Treatment Block Grant funds submit individual client data to TEDS. TEDS includes both admission and discharge data sets, and some 1.5 million admissions are reported annually. TEDS includes information about utilization of SUD treatment services and client demographic and outcome information. TEDS is an admission-based system, and TEDS admissions do not represent individuals. Thus, an individual admitted to treatment twice within a calendar year would be counted as two admissions.

**U.S. CENSUS BUREAU** ([www.census.gov/people](http://www.census.gov/people)). Two main sources of Census Bureau data were used in this report: (1) population estimates and (2) population projections.

# OLDER ADULTS BEHAVIORAL HEALTH PROFILE

---

Wyoming

# Wyoming

## OLDER ADULTS BEHAVIORAL HEALTH PROFILE

August 2016

### WYOMING'S POPULATION

#### Wyoming Population by Age Group

Wyoming is home to 584,153 people. Of these:

- 203,014 (34.8 percent) are over age 50.
- 119,459 (20.4 percent) are over age 60.
- 52,929 (9.1 percent) are over age 70.
- 19,761 (3.4 percent) are ages 80 and older.

The proportion of women rises fairly steadily in each age group, and women make up 59.3 percent of the 80+ group. The racial/ethnic composition of older Wyomingites is as follows:

#### Race/Ethnicity of Wyomingites Ages 50+


White	AI/AN	Black	Asian	NH/PI	Other	Hispanic
95.6%	1.6%	1.1%	0.7%	0.1%	0.9%	5.4%

Source: U.S. Census Bureau, 2015

AI/AN stands for American Indian and Alaska Native.

NH/PI stands for Native Hawaiian and Other Pacific Islander.


Exhibit 1. Wyoming Population by Age Group, 2014



Source: U.S. Census Bureau, 2015

#### The Number of Older Wyomingites Is Growing

Exhibit 2. Wyoming Population by Age Group, 2000–2030



The proportion of Wyoming's population that is 65 and older is growing while the proportion that is younger than 65 is shrinking. The U.S. Census Bureau estimates that 26.5 percent of Wyoming's population will be 65 and older by the year 2030, an increase of 56.0 percent from 2015.

#### Projected Population in Wyoming

Age Group	2015	2025	2030
Under 18	21.9%	20.1%	19.1%
18 to 44	32.9%	30.9%	28.8%
45 to 64	28.4%	24.7%	25.5%
65+	16.8%	24.3%	26.5%

Source: U.S. Census Bureau, 2005

## SUICIDE AMONG OLDER WYOMINGITES


### Region 8 Suicide Rates Compared With National Rates

Suicide data for Wyomingites of various ages were unavailable for 2013. Therefore, the rates for Region 8 (including Colorado, Montana, North Dakota, South Dakota, and Utah) are used instead.

The suicide rate among individuals ages 50+ in Region 8 was higher than the rate among younger age groups. In 2013, the total suicide rate among all people ages 50+ was 25.7 per 100,000 people (10.8 for women and 41.9 for men). The rate among those ages 50–64 was higher than the rate in the United States.

States vary in their reporting of suicides. The suicide rate is influenced by these reporting practices.


**Exhibit 3. Suicide Rates in Region 8 and the United States, 2013**



Source: Centers for Disease Control and Prevention (CDC), National Center for Health Statistics (NCHS), Underlying Cause of Death 1999-2013 on CDC WONDER Online Database, released 2015

### Trends in Suicide Rates in Region 8

**Exhibit 4. Trends in Suicide Rates in Region 8 by Age Group, 2004–2013**



Source: CDC, NCHS, Underlying Cause of Death 1999-2013 on CDC WONDER Online Database, released 2015

Suicide data for Wyomingites of various ages were unavailable. Therefore, the rates for Region 8 are used instead.

The suicide rate among individuals in Region 8 ages 50+ fluctuated from a low of 20.5 per 100,000 in 2004 to a high of 25.8 per 100,000 in 2012. From 2004 to 2013, the rate was generally highest among those in the 50–64 age group.

How a state reports suicides can vary from year to year. The number of suicides is generally low, so even a small difference in reported numbers may make the rate fluctuate widely.


# Wyoming

## SUBSTANCE USE DISORDER AND SUBSTANCE USE DISORDER TREATMENT AMONG OLDER WYOMINGITES

### 30-Day Binge Drinking Among Older Wyomingites

Binge drinking, defined as five or more drinks for men and four or more drinks for women on a single occasion, can lead to serious health problems. Such problems include neurological damage, cardiovascular disease, liver disease, stroke, and poor control of diabetes. Binge drinkers are more likely to take risks such as driving while intoxicated and to experience falls and other accidents. Older people have a lower tolerance for alcohol. Binge drinking decreases with age and occurs more frequently among men than it does among women. As Exhibit 5 shows, 14.3 percent of Wyoming men ages 50–64 reported binge drinking in the past 30 days, while 1.2 percent of those in the 65+ group reported similar behavior.


Exhibit 5. Binge Drinking Rates in Wyoming by Age Group and Sex, 2013



Source: Behavioral Risk Factor Surveillance System (BRFSS), 2013

### Illicit Drug Use Among Older Americans

Exhibit 6. Illicit Drug Use Among Older Americans, 2002–2013



Source: National Survey on Drug Use and Health (NSDUH), 2013  
 Illicit drug use includes illegal drugs and prescription drugs used nonmedically. SAMHSA provides a list of drugs included in its survey in the NSDUH methodological summary.

Nationally, the rate of illicit drug use among older adults ages 50–64 more than doubled from 2002 to 2013. This development partially reflects the entrance into this age group of the baby boom cohort, whose rates of illicit drug use have been higher than those of older cohorts. Although state-specific data are not available, the *Wyoming Behavioral Health Barometer* is available for download from the Substance Abuse and Mental Health Services Administration (SAMHSA) website ([www.samhsa.gov/data/population-data-nsduh/reports?tab=33](http://www.samhsa.gov/data/population-data-nsduh/reports?tab=33)).

# Wyoming


## Admissions to Substance Use Disorder Treatment Among Older Wyomingites

In 2012, there were 640 admissions of Wyomingites ages 50 and older to substance use disorder (SUD) treatment in state-funded treatment programs, a rate of 315.2 per 100,000 people ages 50+. This rate was lower than the regional rate and higher than the national average. Men made up 72.5 percent of these admissions. Of all admissions, 87.5 percent were White/Caucasian, 1.7 percent were Black/African American, and 8.0 percent were Hispanic.

The principal sources of referral to treatment among those ages 50 and older were:

Self-Referral	Criminal Justice	Other
29.1%	40.7%	30.2%


**Exhibit 7. SUD Treatment Admissions of Adults Ages 50+ in Wyoming, Region 8, and the United States by Sex, 2012**



Source: Treatment Episode Data Set (TEDS), 2012<sup>1</sup>  
Data include only those clients reported to SAMHSA.

## SUD Treatment Admissions Among Wyomingites Ages 50+ by Insurance Type

**Exhibit 8. SUD Treatment Admissions of Adults Ages 50+ in Wyoming, Region 8, and the United States by Insurance Type, 2012**



Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

In Wyoming, 84.3 percent of older adult admissions to SUD treatment were uninsured, 5.6 percent had Medicaid, and 10.2 percent had Medicare. There were no reported admissions for older adults who had private insurance.

### *SUD Treatment Admissions Among Wyomingites Ages 50+ by Primary Sources of Payment*

Self-Pay	Medicaid	Other
0.0%	36.1%	63.9%

Self-pay data include private insurance. Exhibit 8 and the accompanying text reflect the type of health insurance (if any) clients had at admission; the table above represents primary sources of payment.


<sup>1</sup> TEDS data are collected by states as a condition of the Substance Abuse Prevention and Treatment Block Grant. Guidelines suggest that states report all clients admitted to publicly financed treatment; however, states are inconsistent in applying the guidelines. States may structure and implement different quality controls over the data. For example, states may collect different categories of information to answer TEDS questions. Information is then "walked over" to TEDS definitions.

# Wyoming

## Alcohol Use Disorder Treatment Admissions Among Wyomingites Ages 50+

Alcohol was the most frequently cited substance used by older Wyomingites in publicly financed SUD treatment in 2012. It was mentioned as a substance of use in 84.7 percent of admissions among those ages 50+. This was lower than the regional rate and higher than the national rate.

**Exhibit 9. Alcohol Use Treatment Admissions of Adults Ages 50+ in Wyoming, Region 8, and the United States, 2012**




Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

## SUD Treatment Admissions for Non-Alcohol Substance Use

**Exhibit 10. Non-Alcohol Substance Use Treatment Admissions of Adults Ages 50+ in Wyoming, Region 8, and the United States, 2012**

Substances other than alcohol were cited as the primary substances of use for 15.3 percent of older adult admissions to publicly funded treatment in Wyoming.



Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

## Drug-Related Emergency Department Visits Involving Pharmaceutical Misuse and Abuse by Older Adults

SAMHSA periodically releases reports from the Drug Abuse Warning Network (DAWN). DAWN, discontinued in 2011, consisted of a nationwide network of hospital emergency departments (EDs) primarily located in large metropolitan areas. DAWN data consist of professional reviews of ED records to determine the extent to which alcohol and other substance abuse was involved in ED visits. According to the November 25, 2010, *DAWN Report*:


- *In 2004, there were an estimated 115,803 ED visits involving pharmaceutical misuse and abuse by adults aged 50 or older; in 2008, there were 256,097 such visits, representing an increase of 121.1 percent*
- *One fifth (19.7 percent) of ED visits involving pharmaceutical misuse and abuse among older adults were made by persons aged 70 or older*
- *Among ED visits made by older adults, pain relievers were the type of pharmaceutical most commonly involved (43.5 percent), followed by drugs used to treat anxiety or insomnia (31.8 percent) and antidepressants (8.6 percent)*
- *Among patients aged 50 or older who visited the ED for pharmaceutical misuse or abuse, more than half (52.3 percent) were treated and released, and more than one third (37.5 percent) were admitted to the hospital*

Bulleted text in italics above is taken from SAMHSA's Center for Behavioral Health Statistics and Quality. (2010, November 25). *The DAWN Report: Drug-related emergency department visits involving pharmaceutical misuse and abuse by older adults*. Rockville, MD: SAMHSA.

## CO-OCCURRING SUBSTANCE USE AND MENTAL DISORDERS

### Older Wyomingites in SUD Treatment With Co-Occurring Mental Disorders

**Exhibit 11. SUD Treatment Admissions of Adults Ages 50+ With a Co-Occurring Mental Disorder in Wyoming, Region 8, and the United States, 2012**



Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

The national literature shows a strong relationship between substance use and mental disorders. Studies show that 30–80 percent of individuals with a substance use or mental disorder also have a co-occurring disorder.

Exhibit 11 shows the proportion of SUD treatment admissions of Wyomingites ages 50+ with a co-occurring mental disorder. This rate is higher than the regional rate and lower than the national average. However, state reporting practices are a factor in these results.




## MENTAL HEALTH

### Older Wyomingites Reporting Frequent Mental Distress

BRFSS, a household survey conducted in all 50 states and several territories, asks the following question: "Now thinking about your mental health, which includes stress, depression, and problems with emotions, for how many days during the past 30 days was your mental health not good?" Those individuals reporting 14 or more "Yes" days in response to this question experience frequent mental distress (FMD). Exhibit 12 shows that older Wyomingites experience FMD at a rate that is lower than the regional and national rates.


**Exhibit 12. Individuals Reporting Frequent Mental Distress in Wyoming, Region 8, and the United States, 2013**



Source: BRFSS, 2013

### Older Wyomingites Reporting Frequent Mental Distress by Age Group and Sex

**Exhibit 13. Wyomingites Reporting Frequent Mental Distress by Age Group and Sex, 2013**



Source: BRFSS, 2013

Older men in Wyoming were more likely to indulge in binge drinking, but older women were more likely to report that they had FMD (14 days or more per 30-day period). As Exhibit 13 shows, 12.2 percent of women in the 50–64 age group and 6.4 percent in the 65+ age group reported FMD, while 8.3 percent of men in the 50–64 age group and 3.9 percent in the 65+ age group reported FMD.

## Other Measures of Mental Health

BRFSS collected other measures showing risk factors for mental and/or physical illness. These included:

- Social and Emotional Support (2010). BRFSS asked, "How often do you get the social and emotional support you need?" The possible responses were always, usually, sometimes, rarely, or never.
- Life Satisfaction (2010). BRFSS asked, "In general, how satisfied are you with your life?" The possible responses were very satisfied, satisfied, dissatisfied, or very dissatisfied.

Exhibit 14 presents the results of these surveys among older Wyomingites.

**Exhibit 14. BRFSS Measures, 2010**


Indicator	Ages 50+	Ages 50–64	Ages 65+
Rarely or never get social or emotional support	7.7%	5.9%	10.8%
Dissatisfied or very dissatisfied	4.1%	4.9%	2.6%

Source: BRFSS, 2010

## Individuals With Frequent Mental Distress Report High Rates of Poor Physical Health

Older Americans who experienced FMD were more likely to report that their physical health was poor (14 days or more in the past 30-day period when physical health was "not good"). As shown in Exhibit 15, although nearly 11 percent of older Americans with no mental distress reported poor physical health, more than 50 percent of those with FMD reported poor physical health.

**Exhibit 15. Individuals Ages 50+ in the United States Reporting Poor Physical Health by Level of Mental Distress, 2013**




Source: BRFSS, 2013

## Relationship Among Mental Distress, Diabetes, Stroke, Heart Attack, High Blood Pressure, and Coronary Disease

Older Americans who experience FMD are more likely to report that they have health problems. People with FMD were more than twice as likely to report having a stroke than those with some or no mental distress. They experienced coronary disease and heart attack at more than 1.6 times, diabetes at more than 1.4 times, and high blood pressure at nearly 1.2 times the rate of those with some or no mental distress.

**Exhibit 16. Individuals Ages 50+ in the United States With Mental Distress and Serious Health Problems, 2013**



Source: BRFSS, 2013

## Older Wyomingites Admitted to State Mental Health Services

Approximately 4.1 percent of the people served by the Wyoming mental health system were ages 65 and older. This represents more than 710 adults.

Source: Center for Mental Health Services (CMHS) Uniform Reporting System (URS) Output Tables, 2014

## DATA SOURCES

**BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM** ([www.cdc.gov/brfss](http://www.cdc.gov/brfss)). BRFSS, managed by CDC, is the largest ongoing telephone health survey system in the world, tracking health conditions and risk behaviors in the United States annually since 1984. Data are collected on a monthly basis in all 50 states, the District of Columbia, and participating territories. BRFSS data are collected by local jurisdictions and reported to CDC.

**CENTERS FOR DISEASE CONTROL AND PREVENTION, NATIONAL CENTER FOR HEALTH STATISTICS, UNDERLYING CAUSE OF DEATH 1999-2013 ON CDC WONDER ONLINE DATABASE, RELEASED 2015** (<http://wonder.cdc.gov/ucd-icd10.html>). The WONDER online database "contains mortality and population counts for all U.S. counties. Data are based on death certificates for U.S. residents. Each death certificate identifies a single underlying cause of death and demographic data. The number of deaths, crude death rates or age-adjusted death rates, and 95% confidence intervals and standard errors for death rates can be obtained by place of residence (total U.S., region, state and county), age group (single-year-of age, 5-year age groups, 10-year age groups and infant age groups), race, Hispanic ethnicity, gender, year, cause-of-death (4-digit ICD-10 code or group of codes), injury intent and injury mechanism, drug/alcohol induced causes and urbanization categories. Data are also available for place of death, month and week day of death, and whether an autopsy was performed."

**CENTER FOR MENTAL HEALTH SERVICES UNIFORM REPORTING SYSTEM** ([www.samhsa.gov/data/us\\_map](http://www.samhsa.gov/data/us_map)). States that receive CMHS Block Grants are required to report aggregate data to URS. URS reports include information about utilization of mental health services and client demographic and outcome information.

**NATIONAL SURVEY ON DRUG USE AND HEALTH** (<https://nsduhweb.rti.org>). The NSDUH, managed by SAMHSA, is "an annual nationwide survey involving interviews with approximately 70,000 randomly selected individuals aged 12 and older." NSDUH data are most frequently used by state planners to assess the need for SUD treatment. NSDUH data also include information about mental health conditions.

**TREATMENT EPISODE DATA SET** ([www.samhsa.gov/data/client-level-data-teds/reports?tab=19](http://www.samhsa.gov/data/client-level-data-teds/reports?tab=19)). States that receive Substance Abuse Prevention and Treatment Block Grant funds submit individual client data to TEDS. TEDS includes both admission and discharge data sets, and some 1.5 million admissions are reported annually. TEDS includes information about utilization of SUD treatment services and client demographic and outcome information. TEDS is an admission-based system, and TEDS admissions do not represent individuals. Thus, an individual admitted to treatment twice within a calendar year would be counted as two admissions.

**U.S. CENSUS BUREAU** ([www.census.gov/people](http://www.census.gov/people)). Two main sources of Census Bureau data were used in this report: (1) population estimates and (2) population projections.