



OLDER ADULTS BEHAVIORAL HEALTH PROFILES

REGION 5

ILLINOIS
INDIANA
MICHIGAN
MINNESOTA
OHIO
WISCONSIN



A Behavioral Health Resource
SAMHSA's State Technical Assistance Contract
August 2016

OLDER ADULTS BEHAVIORAL HEALTH PROFILES

OVERVIEW

The Substance Abuse and Mental Health Services Administration recognizes the importance of behavioral health for older adults and the key role states and communities play in addressing the needs of this vital and growing population. Administrators need clear and concise information to plan, implement, and evaluate effective prevention and treatment efforts in their states.

The Older Adults Behavioral Health Profiles help states and communities identify focus areas for behavioral health plans, select population-level goals, and coordinate and target services to address priority issues. The profiles compare state trends with those in the region and the nation. State and community administrators, planners, and providers can use the information in these profiles and their own data, knowledge, and experience to establish and implement policies that improve the health and future of our valued older citizens.

Disclaimer: Data were current at the time this document was prepared.

OLDER ADULTS BEHAVIORAL HEALTH PROFILE

Illinois

ILLINOIS' POPULATION

Illinois Population by Age Group

Illinois is home to 12,880,580 people.
Of these:

- 4,327,442 (33.6 percent) are over age 50.
- 2,528,593 (19.6 percent) are over age 60.
- 1,206,802 (9.4 percent) are over age 70.
- 481,321 (3.7 percent) are ages 80 and older.


The proportion of women rises fairly steadily in each age group, and women make up 63.9 percent of the 80+ group. The racial/ethnic composition of older Illinoisans is as follows:

Race/Ethnicity of Illinoisans Ages 50+

White	AI/AN	Black	Asian	NH/PI	Other	Hispanic
82.0%	0.4%	12.7%	4.3%	0.0%	0.7%	8.4%

Source: U.S. Census Bureau, 2015
AI/AN stands for American Indian and Alaska Native.
NH/PI stands for Native Hawaiian and Other Pacific Islander.


Exhibit 1. Illinois Population by Age Group, 2014



Source: U.S. Census Bureau, 2015

The Number of Older Illinoisans Is Growing

Exhibit 2. Illinois Population by Age Group, 2000–2030



The proportion of Illinois' population that is 65 and older is growing while the proportion that is younger than 65 is shrinking. The U.S. Census Bureau estimates that 18.0 percent of Illinois' population will be 65 and older by the year 2030, an increase of 35.7 percent from 2015.

Projected Population in Illinois

Age Group	2015	2025	2030
Under 18	24.5%	24.5%	24.3%
18 to 44	36.6%	35.6%	35.1%
45 to 64	25.3%	23.3%	22.6%
65+	13.6%	16.7%	18.0%

Source: U.S. Census Bureau, 2005


SUICIDE AMONG OLDER ILLINOISANS

Illinois Suicide Rate Compared With Regional and National Rates

The suicide rate among Illinoisans ages 50 and older is higher than the rate among younger age groups. In 2013, the total suicide rate among all people ages 50+ was 13.5 per 100,000 people (5.2 for women and 23.3 for men). The rate among those ages 50–64 was lower than both the rate in the region (including Indiana, Michigan, Minnesota, Ohio, and Wisconsin) and the rate in the United States.

States vary in their reporting of suicides. The suicide rate is influenced by these reporting practices.


Exhibit 3. Suicide Rates in Illinois, Region 5, and the United States, 2013



Source: Centers for Disease Control and Prevention (CDC), National Center for Health Statistics (NCHS), Underlying Cause of Death 1999-2013 on CDC WONDER Online Database, released 2015

Trends in Suicide Rates in Illinois

Exhibit 4. Trends in Suicide Rates in Illinois by Age Group, 2004–2013



Source: CDC, NCHS, Underlying Cause of Death 1999-2013 on CDC WONDER Online Database, released 2015

The suicide rate among Illinoisans ages 50+ fluctuated from a low of 10.3 per 100,000 in 2004 and 2006 to a high of 13.9 per 100,000 in 2012. From 2004 to 2013, the rate was highest among those in the 50–64 age group.


How a state reports suicides can vary from year to year. The number of suicides is generally low, so even a small difference in reported numbers may make the rate fluctuate widely.

SUBSTANCE USE DISORDER AND SUBSTANCE USE DISORDER TREATMENT AMONG OLDER ILLINOISANS

30-Day Binge Drinking Among Older Illinoisans

Binge drinking, defined as five or more drinks for men and four or more drinks for women on a single occasion, can lead to serious health problems. Such problems include neurological damage, cardiovascular disease, liver disease, stroke, and poor control of diabetes. Binge drinkers are more likely to take risks such as driving while intoxicated and to experience falls and other accidents. Older people have a lower tolerance for alcohol. Binge drinking decreases with age and occurs more frequently among men than it does among women. As Exhibit 5 shows, 25.3 percent of Illinois men ages 50–64 reported binge drinking in the past 30 days, while 6.9 percent of those in the 65+ group reported similar behavior.


Exhibit 5. Binge Drinking Rates in Illinois by Age Group and Sex, 2013



Source: Behavioral Risk Factor Surveillance System (BRFSS), 2013

Illicit Drug Use Among Older Americans

Exhibit 6. Illicit Drug Use Among Older Americans, 2002–2013



Source: National Survey on Drug Use and Health (NSDUH), 2013
 Illicit drug use includes illegal drugs and prescription drugs used nonmedically. SAMHSA provides a list of drugs included in its survey in the NSDUH methodological summary.

Nationally, the rate of illicit drug use among older adults ages 50–64 more than doubled from 2002 to 2013. This development partially reflects the entrance into this age group of the baby boom cohort, whose rates of illicit drug use have been higher than those of older cohorts. Although state-specific data are not available, the *Illinois Behavioral Health Barometer* is available for download from the Substance Abuse and Mental Health Services Administration (SAMHSA) website (www.samhsa.gov/data/population-data-nsduh/reports?tab=33).


Admissions to Substance Use Disorder Treatment Among Older Illinoisans

In 2012, there were 5,035 admissions of Illinoisans ages 50 and older to substance use disorder (SUD) treatment in state-funded treatment programs, a rate of 116.4 per 100,000 people ages 50+. This rate was lower than the regional rate and the national average. Men made up 65.5 percent of these admissions. Of all admissions, 35.5 percent were White/Caucasian, 60.0 percent were Black/African American, and 6.9 percent were Hispanic.

The principal sources of referral to treatment among those ages 50 and older were:

Self-Referral	Criminal Justice	Other
42.3%	20.9%	36.8%


Exhibit 7. SUD Treatment Admissions of Adults Ages 50+ in Illinois, Region 5, and the United States by Sex, 2012



Source: Treatment Episode Data Set (TEDS), 2012¹
Data include only those clients reported to SAMHSA.

SUD Treatment Admissions Among Illinoisans Ages 50+ by Insurance Type

Exhibit 8. SUD Treatment Admissions of Adults Ages 50+ in Illinois, Region 5, and the United States by Insurance Type, 2012



Source: TEDS, 2012
Data include only those clients reported to SAMHSA.

In Illinois, 61.2 percent of older adult admissions to SUD treatment were uninsured, 32.0 percent had Medicaid, 4.9 percent had Medicare, and 2.0 percent had private insurance.

SUD Treatment Admissions Among Illinoisans Ages 50+ by Primary Sources of Payment

Self-Pay	Medicaid	Other
5.2%	84.7%	10.1%


Self-pay data include private insurance. Exhibit 8 and the accompanying text reflect the type of health insurance (if any) clients had at admission; the table above represents primary sources of payment.

¹ TEDS data are collected by states as a condition of the Substance Abuse Prevention and Treatment Block Grant. Guidelines suggest that states report all clients admitted to publicly financed treatment; however, states are inconsistent in applying the guidelines. States may structure and implement different quality controls over the data. For example, states may collect different categories of information to answer TEDS questions. Information is then "walked over" to TEDS definitions.

Alcohol Use Disorder Treatment Admissions Among Illinoisans Ages 50+

Alcohol was the most frequently cited substance used by older Illinoisans in publicly financed SUD treatment in 2012. It was mentioned as a substance of use in 40.7 percent of admissions among those ages 50+. This was lower than the regional and national rates.

Exhibit 9. Alcohol Use Treatment Admissions of Adults Ages 50+ in Illinois, Region 5, and the United States, 2012




Source: TEDS, 2012
Data include only those clients reported to SAMHSA.

SUD Treatment Admissions for Non-Alcohol Substance Use

Exhibit 10. Non-Alcohol Substance Use Treatment Admissions of Adults Ages 50+ in Illinois, Region 5, and the United States, 2012

Substances other than alcohol were cited as the primary substances of use for 59.3 percent of older adult admissions to publicly funded treatment in Illinois.



Source: TEDS, 2012
Data include only those clients reported to SAMHSA.

Drug-Related Emergency Department Visits Involving Pharmaceutical Misuse and Abuse by Older Adults

SAMHSA periodically releases reports from the Drug Abuse Warning Network (DAWN). DAWN, discontinued in 2011, consisted of a nationwide network of hospital emergency departments (EDs) primarily located in large metropolitan areas. DAWN data consist of professional reviews of ED records to determine the extent to which alcohol and other substance abuse was involved in ED visits. According to the November 25, 2010, *DAWN Report*:


- *In 2004, there were an estimated 115,803 ED visits involving pharmaceutical misuse and abuse by adults aged 50 or older; in 2008, there were 256,097 such visits, representing an increase of 121.1 percent*
- *One fifth (19.7 percent) of ED visits involving pharmaceutical misuse and abuse among older adults were made by persons aged 70 or older*
- *Among ED visits made by older adults, pain relievers were the type of pharmaceutical most commonly involved (43.5 percent), followed by drugs used to treat anxiety or insomnia (31.8 percent) and antidepressants (8.6 percent)*
- *Among patients aged 50 or older who visited the ED for pharmaceutical misuse or abuse, more than half (52.3 percent) were treated and released, and more than one third (37.5 percent) were admitted to the hospital*

Bulleted text in italics above is taken from SAMHSA's Center for Behavioral Health Statistics and Quality. (2010, November 25). *The DAWN Report: Drug-related emergency department visits involving pharmaceutical misuse and abuse by older adults*. Rockville, MD: SAMHSA.

CO-OCCURRING SUBSTANCE USE AND MENTAL DISORDERS

Older Illinoisans in SUD Treatment With Co-Occurring Mental Disorders

Exhibit 11. SUD Treatment Admissions of Adults Ages 50+ With a Co-Occurring Mental Disorder in Illinois, Region 5, and the United States, 2012



Source: TEDS, 2012
Data include only those clients reported to SAMHSA.

The national literature shows a strong relationship between substance use and mental disorders. Studies show that 30–80 percent of individuals with a substance use or mental disorder also have a co-occurring disorder.


Exhibit 11 shows the proportion of SUD treatment admissions of Illinoisans ages 50+ with a co-occurring mental disorder. This rate is lower than the regional rate and the national average. However, state reporting practices are a factor in these results.

MENTAL HEALTH

Older Illinoisans Reporting Frequent Mental Distress

BRFSS, a household survey conducted in all 50 states and several territories, asks the following question: “Now thinking about your mental health, which includes stress, depression, and problems with emotions, for how many days during the past 30 days was your mental health not good?” Those individuals reporting 14 or more “Yes” days in response to this question experience frequent mental distress (FMD). Exhibit 12 shows that Illinoisans in the 50–64 age group experience FMD at a rate that is lower than the regional and national rates, while those in the 65+ age group experience it at a rate that is higher than the regional rate and roughly similar to the national rate.


Exhibit 12. Individuals Reporting Frequent Mental Distress in Illinois, Region 5, and the United States, 2013



Source: BRFSS, 2013

Older Illinoisans Reporting Frequent Mental Distress by Age Group and Sex

Exhibit 13. Illinoisans Reporting Frequent Mental Distress by Age Group and Sex, 2013



Source: BRFSS, 2013

As Exhibit 13 shows, 7.2 percent of men in the 50–64 age group reported FMD (14 days or more per 30-day period) compared with 12.3 percent of women in this age group. Men in the 65+ age group reported a higher rate of FMD than women in the same age group (9.4 percent compared with 6.0 percent). Minnesota is the only other state in the region where men in the 65+ age group reported a higher rate of FMD than did women in this age group.

Other Measures of Mental Health

BRFSS collected other measures showing risk factors for mental and/or physical illness. These included:

- Social and Emotional Support (2010). BRFSS asked, “How often do you get the social and emotional support you need?” The possible responses were always, usually, sometimes, rarely, or never.
- Life Satisfaction (2010). BRFSS asked, “In general, how satisfied are you with your life?” The possible responses were very satisfied, satisfied, dissatisfied, or very dissatisfied.

Exhibit 14 presents the results of these surveys among older Illinoisans.

Exhibit 14. BRFSS Measures, 2010


Indicator	Ages 50+	Ages 50–64	Ages 65+
Rarely or never get social or emotional support	9.6%	7.6%	12.5%
Dissatisfied or very dissatisfied	4.1%	4.9%	3.1%

Source: BRFSS, 2010

Individuals With Frequent Mental Distress Report High Rates of Poor Physical Health

Older Americans who experienced FMD were more likely to report that their physical health was poor (14 days or more in the past 30-day period when physical health was “not good”). As shown in Exhibit 15, although nearly 11 percent of older Americans with no mental distress reported poor physical health, more than 50 percent of those with FMD reported poor physical health.

Exhibit 15. Individuals Ages 50+ in the United States Reporting Poor Physical Health by Level of Mental Distress, 2013




Source: BRFSS, 2013

Relationship Among Mental Distress, Diabetes, Stroke, Heart Attack, High Blood Pressure, and Coronary Disease

Older Americans who experience FMD are more likely to report that they have health problems. People with FMD were more than twice as likely to report having a stroke than those with some or no mental distress. They experienced coronary disease and heart attack at more than 1.6 times, diabetes at more than 1.4 times, and high blood pressure at nearly 1.2 times the rate of those with some or no mental distress.

Exhibit 16. Individuals Ages 50+ in the United States With Mental Distress and Serious Health Problems, 2013



Source: BRFSS, 2013

Older Illinoisans Admitted to State Mental Health Services

Approximately 3.3 percent of the people served by the Illinois mental health system were ages 65 and older. This represents more than 4,470 adults.

Source: Center for Mental Health Services (CMHS) Uniform Reporting System (URS) Output Tables, 2014

DATA SOURCES

BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM (www.cdc.gov/brfss). BRFSS, managed by CDC, is the largest ongoing telephone health survey system in the world, tracking health conditions and risk behaviors in the United States annually since 1984. Data are collected on a monthly basis in all 50 states, the District of Columbia, and participating territories. BRFSS data are collected by local jurisdictions and reported to CDC.

CENTERS FOR DISEASE CONTROL AND PREVENTION, NATIONAL CENTER FOR HEALTH STATISTICS, UNDERLYING CAUSE OF DEATH 1999-2013 ON CDC WONDER ONLINE DATABASE, RELEASED 2015 (<http://wonder.cdc.gov/ucd-icd10.html>). The WONDER online database "contains mortality and population counts for all U.S. counties. Data are based on death certificates for U.S. residents. Each death certificate identifies a single underlying cause of death and demographic data. The number of deaths, crude death rates or age-adjusted death rates, and 95% confidence intervals and standard errors for death rates can be obtained by place of residence (total U.S., region, state and county), age group (single-year-of age, 5-year age groups, 10-year age groups and infant age groups), race, Hispanic ethnicity, gender, year, cause-of-death (4-digit ICD-10 code or group of codes), injury intent and injury mechanism, drug/alcohol induced causes and urbanization categories. Data are also available for place of death, month and week day of death, and whether an autopsy was performed."

CENTER FOR MENTAL HEALTH SERVICES UNIFORM REPORTING SYSTEM (www.samhsa.gov/data/us_map). States that receive CMHS Block Grants are required to report aggregate data to URS. URS reports include information about utilization of mental health services and client demographic and outcome information.

NATIONAL SURVEY ON DRUG USE AND HEALTH (<https://nsduhweb.rti.org>). The NSDUH, managed by SAMHSA, is "an annual nationwide survey involving interviews with approximately 70,000 randomly selected individuals aged 12 and older." NSDUH data are most frequently used by state planners to assess the need for SUD treatment. NSDUH data also include information about mental health conditions.

TREATMENT EPISODE DATA SET (www.samhsa.gov/data/client-level-data-teds/reports?tab=19). States that receive Substance Abuse Prevention and Treatment Block Grant funds submit individual client data to TEDS. TEDS includes both admission and discharge data sets, and some 1.5 million admissions are reported annually. TEDS includes information about utilization of SUD treatment services and client demographic and outcome information. TEDS is an admission-based system, and TEDS admissions do not represent individuals. Thus, an individual admitted to treatment twice within a calendar year would be counted as two admissions.

U.S. CENSUS BUREAU (www.census.gov/people). Two main sources of Census Bureau data were used in this report: (1) population estimates and (2) population projections.

OLDER ADULTS BEHAVIORAL HEALTH PROFILE

Indiana

Indiana

OLDER ADULTS BEHAVIORAL HEALTH PROFILE

August 2016

INDIANA'S POPULATION

Indiana Population by Age Group

Indiana is home to 6,596,855 people. Of these:

- 2,252,368 (34.1 percent) are over age 50.
- 1,330,688 (20.2 percent) are over age 60.
- 630,225 (9.6 percent) are over age 70.
- 243,803 (3.7 percent) are ages 80 and older.

The proportion of women rises fairly steadily in each age group, and women make up 63.7 percent of the 80+ group. The racial/ethnic composition of older Indians is as follows:

Race/Ethnicity of Indians Ages 50+


White	AI/AN	Black	Asian	NH/PI	Other	Hispanic
90.8%	0.3%	7.2%	1.1%	0.0%	0.6%	2.6%

Source: U.S. Census Bureau, 2015

AI/AN stands for American Indian and Alaska Native.

NH/PI stands for Native Hawaiian and Other Pacific Islander.


Exhibit 1. Indiana Population by Age Group, 2014



Source: U.S. Census Bureau, 2015

The Number of Older Indians Is Growing

Exhibit 2. Indiana Population by Age Group, 2000–2030



The proportion of Indiana's population that is 65 and older is growing while the proportion that is younger than 65 is shrinking. The U.S. Census Bureau estimates that 18.1 percent of Indiana's population will be 65 and older by the year 2030, an increase of 35.9 percent from 2015.

Projected Population in Indiana

Age Group	2015	2025	2030
Under 18	24.8%	24.8%	25.0%
18 to 44	35.8%	34.7%	34.2%
45 to 64	25.5%	23.5%	22.7%
65+	13.9%	17.0%	18.1%

Source: U.S. Census Bureau, 2005


SUICIDE AMONG OLDER INDIANIANS

Indiana Suicide Rate Compared With Regional and National Rates

The suicide rate among Indianians ages 50 and older is higher than the rate among younger age groups. In 2013, the total suicide rate among all people ages 50+ was 17.8 per 100,000 people (6.2 for women and 31.2 for men). The rate among those ages 50–64 was higher than both the rate in the region (including Illinois, Michigan, Minnesota, Ohio, and Wisconsin) and the rate in the United States.

States vary in their reporting of suicides. The suicide rate is influenced by these reporting practices.


Exhibit 3. Suicide Rates in Indiana, Region 5, and the United States, 2013



Source: Centers for Disease Control and Prevention (CDC), National Center for Health Statistics (NCHS), Underlying Cause of Death 1999-2013 on CDC WONDER Online Database, released 2015

Trends in Suicide Rates in Indiana

Exhibit 4. Trends in Suicide Rates in Indiana by Age Group, 2004–2013



Source: CDC, NCHS, Underlying Cause of Death 1999-2013 on CDC WONDER Online Database, released 2015

The suicide rate among Indianians ages 50+ fluctuated from a low of 12.8 per 100,000 in 2004 to a high of 17.8 per 100,000 in 2010 and 2013. From 2004 to 2013, the rate was highest among those in the 50–64 age group.


How a state reports suicides can vary from year to year. The number of suicides is generally low, so even a small difference in reported numbers may make the rate fluctuate widely.

SUBSTANCE USE DISORDER AND SUBSTANCE USE DISORDER TREATMENT AMONG OLDER INDIANIANS

30-Day Binge Drinking Among Older Indianians

Binge drinking, defined as five or more drinks for men and four or more drinks for women on a single occasion, can lead to serious health problems. Such problems include neurological damage, cardiovascular disease, liver disease, stroke, and poor control of diabetes. Binge drinkers are more likely to take risks such as driving while intoxicated and to experience falls and other accidents. Older people have a lower tolerance for alcohol. Binge drinking decreases with age and occurs more frequently among men than it does among women. As Exhibit 5 shows, 15.1 percent of Indiana men ages 50–64 reported binge drinking in the past 30 days, while 5.1 percent of those in the 65+ group reported similar behavior.


Exhibit 5. Binge Drinking Rates in Indiana by Age Group and Sex, 2013



Source: Behavioral Risk Factor Surveillance System (BRFSS), 2013

Illicit Drug Use Among Older Americans

Exhibit 6. Illicit Drug Use Among Older Americans, 2002–2013



Source: National Survey on Drug Use and Health (NSDUH), 2013
 Illicit drug use includes illegal drugs and prescription drugs used nonmedically. SAMHSA provides a list of drugs included in its survey in the NSDUH methodological summary.

Nationally, the rate of illicit drug use among older adults ages 50–64 more than doubled from 2002 to 2013. This development partially reflects the entrance into this age group of the baby boom cohort, whose rates of illicit drug use have been higher than those of older cohorts. Although state-specific data are not available, the *Indiana Behavioral Health Barometer* is available for download from the Substance Abuse and Mental Health Services Administration (SAMHSA) website (www.samhsa.gov/data/population-data-nsduh/reports?tab=33).


Admissions to Substance Use Disorder Treatment Among Older Indianians

In 2012, there were 2,718 admissions of Indianians ages 50 and older to substance use disorder (SUD) treatment in state-funded treatment programs, a rate of 120.7 per 100,000 people ages 50+. This rate was lower than the regional rate and the national average. Men made up 70.0 percent of these admissions. Of all admissions, 71.6 percent were White/Caucasian, 25.7 percent were Black/African American, and 3.5 percent were Hispanic.

The principal sources of referral to treatment among those ages 50 and older were:

Self-Referral	Criminal Justice	Other
36.1%	43.9%	20.0%


Exhibit 7. SUD Treatment Admissions of Adults Ages 50+ in Indiana, Region 5, and the United States by Sex, 2012



Source: Treatment Episode Data Set (TEDS), 2012¹
Data include only those clients reported to SAMHSA.

SUD Treatment Admissions Among Indianians Ages 50+ by Insurance Type

Exhibit 8. SUD Treatment Admissions of Adults Ages 50+ in Indiana, Region 5, and the United States by Insurance Type, 2012



Source: TEDS, 2012
Data include only those clients reported to SAMHSA.

In Indiana, 57.5 percent of older adult admissions to SUD treatment were uninsured, 14.4 percent had Medicaid, 20.4 percent had Medicare, and 7.7 percent had private insurance.

SUD Treatment Admissions Among Indianians Ages 50+ by Primary Sources of Payment

Self-Pay	Medicaid	Other
Data not available	Data not available	Data not available


Exhibit 8 and the accompanying text reflect the type of health insurance (if any) clients had at admission; the table above represents primary sources of payment.

¹ TEDS data are collected by states as a condition of the Substance Abuse Prevention and Treatment Block Grant. Guidelines suggest that states report all clients admitted to publicly financed treatment; however, states are inconsistent in applying the guidelines. States may structure and implement different quality controls over the data. For example, states may collect different categories of information to answer TEDS questions. Information is then "walked over" to TEDS definitions.

Alcohol Use Disorder Treatment Admissions Among Indianians Ages 50+

Alcohol was the most frequently cited substance used by older Indianians in publicly financed SUD treatment in 2012. It was mentioned as a substance of use in 65.7 percent of admissions among those ages 50+. This was higher than the regional and national rates.

Exhibit 9. Alcohol Use Treatment Admissions of Adults Ages 50+ in Indiana, Region 5, and the United States, 2012




Source: TEDS, 2012
Data include only those clients reported to SAMHSA.

SUD Treatment Admissions for Non-Alcohol Substance Use

Exhibit 10. Non-Alcohol Substance Use Treatment Admissions of Adults Ages 50+ in Indiana, Region 5, and the United States, 2012

Substances other than alcohol were cited as the primary substances of use for 34.3 percent of older adult admissions to publicly funded treatment in Indiana.



Source: TEDS, 2012
Data include only those clients reported to SAMHSA.

Drug-Related Emergency Department Visits Involving Pharmaceutical Misuse and Abuse by Older Adults

SAMHSA periodically releases reports from the Drug Abuse Warning Network (DAWN). DAWN, discontinued in 2011, consisted of a nationwide network of hospital emergency departments (EDs) primarily located in large metropolitan areas. DAWN data consist of professional reviews of ED records to determine the extent to which alcohol and other substance abuse was involved in ED visits. According to the November 25, 2010, *DAWN Report*:


- *In 2004, there were an estimated 115,803 ED visits involving pharmaceutical misuse and abuse by adults aged 50 or older; in 2008, there were 256,097 such visits, representing an increase of 121.1 percent*
- *One fifth (19.7 percent) of ED visits involving pharmaceutical misuse and abuse among older adults were made by persons aged 70 or older*
- *Among ED visits made by older adults, pain relievers were the type of pharmaceutical most commonly involved (43.5 percent), followed by drugs used to treat anxiety or insomnia (31.8 percent) and antidepressants (8.6 percent)*
- *Among patients aged 50 or older who visited the ED for pharmaceutical misuse or abuse, more than half (52.3 percent) were treated and released, and more than one third (37.5 percent) were admitted to the hospital*

Bulleted text in italics above is taken from SAMHSA's Center for Behavioral Health Statistics and Quality. (2010, November 25). *The DAWN Report: Drug-related emergency department visits involving pharmaceutical misuse and abuse by older adults*. Rockville, MD: SAMHSA.

CO-OCCURRING SUBSTANCE USE AND MENTAL DISORDERS

Older Indianians in SUD Treatment With Co-Occurring Mental Disorders

Exhibit 11. SUD Treatment Admissions of Adults Ages 50+ With a Co-Occurring Mental Disorder in Indiana, Region 5, and the United States, 2012



Source: TEDS, 2012
Data include only those clients reported to SAMHSA.

The national literature shows a strong relationship between substance use and mental disorders. Studies show that 30–80 percent of individuals with a substance use or mental disorder also have a co-occurring disorder.


Exhibit 11 shows the proportion of SUD treatment admissions of Indianians ages 50+ with a co-occurring mental disorder. This rate is higher than the regional rate and the national average. However, state reporting practices are a factor in these results.

MENTAL HEALTH

Older Indianians Reporting Frequent Mental Distress

BRFSS, a household survey conducted in all 50 states and several territories, asks the following question: "Now thinking about your mental health, which includes stress, depression, and problems with emotions, for how many days during the past 30 days was your mental health not good?" Those individuals reporting 14 or more "Yes" days in response to this question experience frequent mental distress (FMD). Exhibit 12 shows that older Indianians experience FMD at a rate that is higher than the regional rate and lower than the national rate.


Exhibit 12. Individuals Reporting Frequent Mental Distress in Indiana, Region 5, and the United States, 2013



Source: BRFSS, 2013

Older Indianians Reporting Frequent Mental Distress by Age Group and Sex

Exhibit 13. Indianians Reporting Frequent Mental Distress by Age Group and Sex, 2013



Source: BRFSS, 2013

Older men in Indiana were more likely to indulge in binge drinking, but older women were more likely to report that they had FMD (14 days or more per 30-day period). As Exhibit 13 shows, 16.1 percent of women in the 50–64 age group and 7.4 percent in the 65+ age group reported FMD, while 9.6 percent of men in the 50–64 age group and 5.7 percent in the 65+ age group reported FMD.

Other Measures of Mental Health

BRFSS collected other measures showing risk factors for mental and/or physical illness. These included:

- Social and Emotional Support (2010). BRFSS asked, “How often do you get the social and emotional support you need?” The possible responses were always, usually, sometimes, rarely, or never.
- Life Satisfaction (2010). BRFSS asked, “In general, how satisfied are you with your life?” The possible responses were very satisfied, satisfied, dissatisfied, or very dissatisfied.

Exhibit 14 presents the results of these surveys among older Indianians.

Exhibit 14. BRFSS Measures, 2010


Indicator	Ages 50+	Ages 50–64	Ages 65+
Rarely or never get social or emotional support	9.2%	7.9%	11.1%
Dissatisfied or very dissatisfied	5.3%	6.8%	3.2%

Source: BRFSS, 2010

Individuals With Frequent Mental Distress Report High Rates of Poor Physical Health

Older Americans who experienced FMD were more likely to report that their physical health was poor (14 days or more in the past 30-day period when physical health was “not good”). As shown in Exhibit 15, although nearly 11 percent of older Americans with no mental distress reported poor physical health, more than 50 percent of those with FMD reported poor physical health.

Exhibit 15. Individuals Ages 50+ in the United States Reporting Poor Physical Health by Level of Mental Distress, 2013




Source: BRFSS, 2013

Relationship Among Mental Distress, Diabetes, Stroke, Heart Attack, High Blood Pressure, and Coronary Disease

Older Americans who experience FMD are more likely to report that they have health problems. People with FMD were more than twice as likely to report having a stroke than those with some or no mental distress. They experienced coronary disease and heart attack at more than 1.6 times, diabetes at more than 1.4 times, and high blood pressure at nearly 1.2 times the rate of those with some or no mental distress.

Exhibit 16. Individuals Ages 50+ in the United States With Mental Distress and Serious Health Problems, 2013



Source: BRFSS, 2013

Older Indianians Admitted to State Mental Health Services

Approximately 2.8 percent of the people served by the Indiana mental health system were ages 65 and older. This represents more than 3,610 adults.

Source: Center for Mental Health Services (CMHS) Uniform Reporting System (URS) Output Tables, 2014

DATA SOURCES

BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM (www.cdc.gov/brfss). BRFSS, managed by CDC, is the largest ongoing telephone health survey system in the world, tracking health conditions and risk behaviors in the United States annually since 1984. Data are collected on a monthly basis in all 50 states, the District of Columbia, and participating territories. BRFSS data are collected by local jurisdictions and reported to CDC.

CENTERS FOR DISEASE CONTROL AND PREVENTION, NATIONAL CENTER FOR HEALTH STATISTICS, UNDERLYING CAUSE OF DEATH 1999-2013 ON CDC WONDER ONLINE DATABASE, RELEASED 2015 (<http://wonder.cdc.gov/ucd-icd10.html>). The WONDER online database “contains mortality and population counts for all U.S. counties. Data are based on death certificates for U.S. residents. Each death certificate identifies a single underlying cause of death and demographic data. The number of deaths, crude death rates or age-adjusted death rates, and 95% confidence intervals and standard errors for death rates can be obtained by place of residence (total U.S., region, state and county), age group (single-year-of age, 5-year age groups, 10-year age groups and infant age groups), race, Hispanic ethnicity, gender, year, cause-of-death (4-digit ICD-10 code or group of codes), injury intent and injury mechanism, drug/alcohol induced causes and urbanization categories. Data are also available for place of death, month and week day of death, and whether an autopsy was performed.”

CENTER FOR MENTAL HEALTH SERVICES UNIFORM REPORTING SYSTEM (www.samhsa.gov/data/us_map). States that receive CMHS Block Grants are required to report aggregate data to URS. URS reports include information about utilization of mental health services and client demographic and outcome information.

NATIONAL SURVEY ON DRUG USE AND HEALTH (<https://nsduhweb.rti.org>). The NSDUH, managed by SAMHSA, is “an annual nationwide survey involving interviews with approximately 70,000 randomly selected individuals aged 12 and older.” NSDUH data are most frequently used by state planners to assess the need for SUD treatment. NSDUH data also include information about mental health conditions.

TREATMENT EPISODE DATA SET (www.samhsa.gov/data/client-level-data-teds/reports?tab=19). States that receive Substance Abuse Prevention and Treatment Block Grant funds submit individual client data to TEDS. TEDS includes both admission and discharge data sets, and some 1.5 million admissions are reported annually. TEDS includes information about utilization of SUD treatment services and client demographic and outcome information. TEDS is an admission-based system, and TEDS admissions do not represent individuals. Thus, an individual admitted to treatment twice within a calendar year would be counted as two admissions.

U.S. CENSUS BUREAU (www.census.gov/people). Two main sources of Census Bureau data were used in this report: (1) population estimates and (2) population projections.

OLDER ADULTS BEHAVIORAL HEALTH PROFILE

Michigan

Michigan

OLDER ADULTS BEHAVIORAL HEALTH PROFILE

August 2016

MICHIGAN'S POPULATION

Michigan Population by Age Group

Michigan is home to 9,909,877 people.
Of these:

- 3,633,119 (36.7 percent) are over age 50.
- 2,163,702 (21.8 percent) are over age 60.
- 1,024,981 (10.3 percent) are over age 70.
- 400,736 (4.0 percent) are ages 80 and older.

The proportion of women rises fairly steadily in each age group, and women make up 63.0 percent of the 80+ group. The racial/ethnic composition of older Michigianians is as follows:

Race/Ethnicity of Michigianians Ages 50+


White	AI/AN	Black	Asian	NH/PI	Other	Hispanic
85.6%	0.6%	11.3%	1.7%	0.0%	0.8%	2.1%

Source: U.S. Census Bureau, 2015

AI/AN stands for American Indian and Alaska Native.

NH/PI stands for Native Hawaiian and Other Pacific Islander.


Exhibit 1. Michigan Population by Age Group, 2014



Source: U.S. Census Bureau, 2015

The Number of Older Michigianians Is Growing

Exhibit 2. Michigan Population by Age Group, 2000–2030



The proportion of Michigan's population that is 65 and older is growing while the proportion that is younger than 65 is shrinking. The U.S. Census Bureau estimates that 19.5 percent of Michigan's population will be 65 and older by the year 2030, an increase of 38.1 percent from 2015.

Projected Population in Michigan

Age Group	2015	2025	2030
Under 18	23.4%	23.1%	22.8%
18 to 44	35.8%	34.5%	33.8%
45 to 64	26.5%	24.4%	24.0%
65+	14.2%	18.0%	19.5%

Source: U.S. Census Bureau, 2005


SUICIDE AMONG OLDER MICHIGANIANS

Michigan Suicide Rate Compared With Regional and National Rates

The suicide rate among Michiganians ages 50 and older is higher than the rate among younger age groups. In 2013, the total suicide rate among all people ages 50+ was 15.1 per 100,000 people (5.8 for women and 25.6 for men). The rate among those ages 50–64 was lower than both the rate in the region (including Illinois, Indiana, Minnesota, Ohio, and Wisconsin) and the rate in the United States.

States vary in their reporting of suicides. The suicide rate is influenced by these reporting practices.


Exhibit 3. Suicide Rates in Michigan, Region 5, and the United States, 2013



Source: Centers for Disease Control and Prevention (CDC), National Center for Health Statistics (NCHS), Underlying Cause of Death 1999–2013 on CDC WONDER Online Database, released 2015

Trends in Suicide Rates in Michigan

Exhibit 4. Trends in Suicide Rates in Michigan by Age Group, 2004–2013



Source: CDC, NCHS, Underlying Cause of Death 1999–2013 on CDC WONDER Online Database, released 2015

The suicide rate among Michiganians ages 50+ fluctuated from a low of 12.8 per 100,000 in 2004 to a high of 16.0 per 100,000 in 2010. From 2004 to 2013, the rate was highest among those in the 50–64 age group.


How a state reports suicides can vary from year to year. The number of suicides is generally low, so even a small difference in reported numbers may make the rate fluctuate widely.

SUBSTANCE USE DISORDER AND SUBSTANCE USE DISORDER TREATMENT AMONG OLDER MICHIGANIANS

30-Day Binge Drinking Among Older Michiganians

Binge drinking, defined as five or more drinks for men and four or more drinks for women on a single occasion, can lead to serious health problems. Such problems include neurological damage, cardiovascular disease, liver disease, stroke, and poor control of diabetes. Binge drinkers are more likely to take risks such as driving while intoxicated and to experience falls and other accidents. Older people have a lower tolerance for alcohol. Binge drinking decreases with age and occurs more frequently among men than it does among women. As Exhibit 5 shows, 19.6 percent of Michigan men ages 50–64 reported binge drinking in the past 30 days, while 10.3 percent of those in the 65+ group reported similar behavior.


Exhibit 5. Binge Drinking Rates in Michigan by Age Group and Sex, 2013



Source: Behavioral Risk Factor Surveillance System (BRFSS), 2013

Illicit Drug Use Among Older Americans

Exhibit 6. Illicit Drug Use Among Older Americans, 2002–2013



Source: National Survey on Drug Use and Health (NSDUH), 2013

Illicit drug use includes illegal drugs and prescription drugs used nonmedically. SAMHSA provides a list of drugs included in its survey in the NSDUH methodological summary.

Nationally, the rate of illicit drug use among older adults ages 50–64 more than doubled from 2002 to 2013. This development partially reflects the entrance into this age group of the baby boom cohort, whose rates of illicit drug use have been higher than those of older cohorts. Although state-specific data are not available, the *Michigan Behavioral Health Barometer* is available for download from the Substance Abuse and Mental Health Services Administration (SAMHSA) website (www.samhsa.gov/data/population-data-nsduh/reports?tab=33).


Admissions to Substance Use Disorder Treatment Among Older Michigianians

In 2012, there were 8,619 admissions of Michigianians ages 50 and older to substance use disorder (SUD) treatment in state-funded treatment programs, a rate of 237.2 per 100,000 people ages 50+. This rate was higher than the regional rate and lower than the national average. Men made up 70.0 percent of these admissions. Of all admissions, 48.6 percent were White/Caucasian, 48.7 percent were Black/African American, and 1.7 percent were Hispanic.

The principal sources of referral to treatment among those ages 50 and older were:

Self-Referral	Criminal Justice	Other
46.8%	14.6%	38.6%


Exhibit 7. SUD Treatment Admissions of Adults Ages 50+ in Michigan, Region 5, and the United States by Sex, 2012



Source: Treatment Episode Data Set (TEDS), 2012¹
Data include only those clients reported to SAMHSA.

SUD Treatment Admissions Among Individuals in Region 5 Ages 50+ by Insurance Type

Exhibit 8. SUD Treatment Admissions of Adults Ages 50+ in Region 5 and the United States by Insurance Type, 2012



Source: TEDS, 2012
Data include only those clients reported to SAMHSA.

States may choose not to report some of the fields to TEDS, including information on treatment admission insurance types. These data were not reported by Michigan in 2012. Therefore, the rates for Region 5 are used instead.

In Region 5, 59.9 percent of older adult admissions to SUD treatment were uninsured, 25.9 percent had Medicaid, 10.2 percent had Medicare, and 3.9 percent had private insurance.

SUD Treatment Admissions Among Michigianians Ages 50+ by Primary Sources of Payment

Self-Pay	Medicaid	Other
Data not available	Data not available	Data not available


Exhibit 8 and the accompanying text reflect the type of health insurance (if any) clients had at admission; the table above represents primary sources of payment.

¹ TEDS data are collected by states as a condition of the Substance Abuse Prevention and Treatment Block Grant. Guidelines suggest that states report all clients admitted to publicly financed treatment; however, states are inconsistent in applying the guidelines. States may structure and implement different quality controls over the data. For example, states may collect different categories of information to answer TEDS questions. Information is then "walked over" to TEDS definitions.

Alcohol Use Disorder Treatment Admissions Among Michiganians Ages 50+

Alcohol was the most frequently cited substance used by older Michiganians in publicly financed SUD treatment in 2012. It was mentioned as a substance of use in 49.3 percent of admissions among those ages 50+. This was lower than the regional and national rates.

Exhibit 9. Alcohol Use Treatment Admissions of Adults Ages 50+ in Michigan, Region 5, and the United States, 2012




Source: TEDS, 2012
Data include only those clients reported to SAMHSA.

SUD Treatment Admissions for Non-Alcohol Substance Use

Exhibit 10. Non-Alcohol Substance Use Treatment Admissions of Adults Ages 50+ in Michigan, Region 5, and the United States, 2012

Substances other than alcohol were cited as the primary substances of use for 50.7 percent of older adult admissions to publicly funded treatment in Michigan.



Source: TEDS, 2012
Data include only those clients reported to SAMHSA.

Drug-Related Emergency Department Visits Involving Pharmaceutical Misuse and Abuse by Older Adults

SAMHSA periodically releases reports from the Drug Abuse Warning Network (DAWN). DAWN, discontinued in 2011, consisted of a nationwide network of hospital emergency departments (EDs) primarily located in large metropolitan areas. DAWN data consist of professional reviews of ED records to determine the extent to which alcohol and other substance abuse was involved in ED visits. According to the November 25, 2010, *DAWN Report*:


- *In 2004, there were an estimated 115,803 ED visits involving pharmaceutical misuse and abuse by adults aged 50 or older; in 2008, there were 256,097 such visits, representing an increase of 121.1 percent*
- *One fifth (19.7 percent) of ED visits involving pharmaceutical misuse and abuse among older adults were made by persons aged 70 or older*
- *Among ED visits made by older adults, pain relievers were the type of pharmaceutical most commonly involved (43.5 percent), followed by drugs used to treat anxiety or insomnia (31.8 percent) and antidepressants (8.6 percent)*
- *Among patients aged 50 or older who visited the ED for pharmaceutical misuse or abuse, more than half (52.3 percent) were treated and released, and more than one third (37.5 percent) were admitted to the hospital*

Bulleted text in italics above is taken from SAMHSA's Center for Behavioral Health Statistics and Quality. (2010, November 25). *The DAWN Report: Drug-related emergency department visits involving pharmaceutical misuse and abuse by older adults*. Rockville, MD: SAMHSA.

CO-OCCURRING SUBSTANCE USE AND MENTAL DISORDERS

Older Michiganians in SUD Treatment With Co-Occurring Mental Disorders

Exhibit 11. SUD Treatment Admissions of Adults Ages 50+ With a Co-Occurring Mental Disorder in Michigan, Region 5, and the United States, 2012



Source: TEDS, 2012
Data include only those clients reported to SAMHSA.

The national literature shows a strong relationship between substance use and mental disorders. Studies show that 30–80 percent of individuals with a substance use or mental disorder also have a co-occurring disorder.


Exhibit 11 shows the proportion of SUD treatment admissions of Michiganians ages 50+ with a co-occurring mental disorder. This rate is lower than the regional rate and the national average. However, state reporting practices are a factor in these results.

MENTAL HEALTH

Older Michiganians Reporting Frequent Mental Distress

BRFSS, a household survey conducted in all 50 states and several territories, asks the following question: "Now thinking about your mental health, which includes stress, depression, and problems with emotions, for how many days during the past 30 days was your mental health not good?" Those individuals reporting 14 or more "Yes" days in response to this question experience frequent mental distress (FMD). Exhibit 12 shows that Michiganians in the 50–64 age group experience FMD at a rate that is higher than the regional and national rates, while those in the 65+ age group experience it at a rate that is higher than the regional rate and lower than the national rate.


Exhibit 12. Individuals Reporting Frequent Mental Distress in Michigan, Region 5, and the United States, 2013



Source: BRFSS, 2013

Older Michiganians Reporting Frequent Mental Distress by Age Group and Sex

Exhibit 13. Michiganians Reporting Frequent Mental Distress by Age Group and Sex, 2013



Source: BRFSS, 2013

Older men in Michigan were more likely to indulge in binge drinking, but older women were more likely to report that they had FMD (14 days or more per 30-day period). As Exhibit 13 shows, 15.7 percent of women in the 50–64 age group and 6.8 percent in the 65+ age group reported FMD, while 11.9 percent of men in the 50–64 age group and 6.7 percent in the 65+ age group reported FMD.

Other Measures of Mental Health

BRFSS collected other measures showing risk factors for mental and/or physical illness. These included:

- Social and Emotional Support (2010). BRFSS asked, "How often do you get the social and emotional support you need?" The possible responses were always, usually, sometimes, rarely, or never.
- Life Satisfaction (2010). BRFSS asked, "In general, how satisfied are you with your life?" The possible responses were very satisfied, satisfied, dissatisfied, or very dissatisfied.

Exhibit 14 presents the results of these surveys among older Michigianians.

Exhibit 14. BRFSS Measures, 2010


Indicator	Ages 50+	Ages 50–64	Ages 65+
Rarely or never get social or emotional support	7.3%	5.3%	10.4%
Dissatisfied or very dissatisfied	4.2%	5.2%	2.8%

Source: BRFSS, 2010

Individuals With Frequent Mental Distress Report High Rates of Poor Physical Health

Older Americans who experienced FMD were more likely to report that their physical health was poor (14 days or more in the past 30-day period when physical health was "not good"). As shown in Exhibit 15, although nearly 11 percent of older Americans with no mental distress reported poor physical health, more than 50 percent of those with FMD reported poor physical health.

Exhibit 15. Individuals Ages 50+ in the United States Reporting Poor Physical Health by Level of Mental Distress, 2013




Source: BRFSS, 2013

Relationship Among Mental Distress, Diabetes, Stroke, Heart Attack, High Blood Pressure, and Coronary Disease

Older Americans who experience FMD are more likely to report that they have health problems. People with FMD were more than twice as likely to report having a stroke than those with some or no mental distress. They experienced coronary disease and heart attack at more than 1.6 times, diabetes at more than 1.4 times, and high blood pressure at nearly 1.2 times the rate of those with some or no mental distress.

Exhibit 16. Individuals Ages 50+ in the United States With Mental Distress and Serious Health Problems, 2013



Source: BRFSS, 2013

Older Michiganians Admitted to State Mental Health Services

Approximately 4.8 percent of the people served by the Michigan mental health system were ages 65 and older. This represents more than 11,860 adults.

Source: Center for Mental Health Services (CMHS) Uniform Reporting System (URS) Output Tables, 2014

DATA SOURCES

BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM (www.cdc.gov/brfss). BRFSS, managed by CDC, is the largest ongoing telephone health survey system in the world, tracking health conditions and risk behaviors in the United States annually since 1984. Data are collected on a monthly basis in all 50 states, the District of Columbia, and participating territories. BRFSS data are collected by local jurisdictions and reported to CDC.

CENTERS FOR DISEASE CONTROL AND PREVENTION, NATIONAL CENTER FOR HEALTH STATISTICS, UNDERLYING CAUSE OF DEATH 1999-2013 ON CDC WONDER ONLINE DATABASE, RELEASED 2015 (<http://wonder.cdc.gov/ucd-icd10.html>). The WONDER online database "contains mortality and population counts for all U.S. counties. Data are based on death certificates for U.S. residents. Each death certificate identifies a single underlying cause of death and demographic data. The number of deaths, crude death rates or age-adjusted death rates, and 95% confidence intervals and standard errors for death rates can be obtained by place of residence (total U.S., region, state and county), age group (single-year-of age, 5-year age groups, 10-year age groups and infant age groups), race, Hispanic ethnicity, gender, year, cause-of-death (4-digit ICD-10 code or group of codes), injury intent and injury mechanism, drug/alcohol induced causes and urbanization categories. Data are also available for place of death, month and week day of death, and whether an autopsy was performed."

CENTER FOR MENTAL HEALTH SERVICES UNIFORM REPORTING SYSTEM (www.samhsa.gov/data/us_map). States that receive CMHS Block Grants are required to report aggregate data to URS. URS reports include information about utilization of mental health services and client demographic and outcome information.

NATIONAL SURVEY ON DRUG USE AND HEALTH (<https://nsduhweb.rti.org>). The NSDUH, managed by SAMHSA, is "an annual nationwide survey involving interviews with approximately 70,000 randomly selected individuals aged 12 and older." NSDUH data are most frequently used by state planners to assess the need for SUD treatment. NSDUH data also include information about mental health conditions.

TREATMENT EPISODE DATA SET (www.samhsa.gov/data/client-level-data-teds/reports?tab=19). States that receive Substance Abuse Prevention and Treatment Block Grant funds submit individual client data to TEDS. TEDS includes both admission and discharge data sets, and some 1.5 million admissions are reported annually. TEDS includes information about utilization of SUD treatment services and client demographic and outcome information. TEDS is an admission-based system, and TEDS admissions do not represent individuals. Thus, an individual admitted to treatment twice within a calendar year would be counted as two admissions.

U.S. CENSUS BUREAU (www.census.gov/people). Two main sources of Census Bureau data were used in this report: (1) population estimates and (2) population projections.

OLDER ADULTS BEHAVIORAL HEALTH PROFILE

Minnesota

Minnesota

OLDER ADULTS BEHAVIORAL HEALTH PROFILE

August 2016

MINNESOTA'S POPULATION

Minnesota Population by Age Group

Minnesota is home to 5,457,173 people. Of these:

- 1,897,121 (34.8 percent) are over age 50.
- 1,107,299 (20.3 percent) are over age 60.
- 529,695 (9.7 percent) are over age 70.
- 216,507 (4.0 percent) are ages 80 and older.


The proportion of women rises fairly steadily in each age group, and women make up 62.2 percent of the 80+ group. The racial/ethnic composition of older Minnesotans is as follows:

Race/Ethnicity of Minnesotans Ages 50+

White	AI/AN	Black	Asian	NH/PI	Other	Hispanic
93.5%	0.8%	2.9%	2.2%	0.0%	0.6%	1.7%

Source: U.S. Census Bureau, 2015
AI/AN stands for American Indian and Alaska Native.
NH/PI stands for Native Hawaiian and Other Pacific Islander.


Exhibit 1. Minnesota Population by Age Group, 2014



Source: U.S. Census Bureau, 2015

The Number of Older Minnesotans Is Growing

Exhibit 2. Minnesota Population by Age Group, 2000–2030



The proportion of Minnesota's population that is 65 and older is growing while the proportion that is younger than 65 is shrinking. The U.S. Census Bureau estimates that 18.9 percent of Minnesota's population will be 65 and older by the year 2030, an increase of 54.1 percent from 2015.

Projected Population in Minnesota

Age Group	2015	2025	2030
Under 18	23.8%	24.1%	23.9%
18 to 44	36.5%	35.3%	34.2%
45 to 64	26.0%	23.2%	23.0%
65+	13.7%	17.4%	18.9%

Source: U.S. Census Bureau, 2005


SUICIDE AMONG OLDER MINNESOTANS

Minnesota Suicide Rate Compared With Regional and National Rates

The suicide rate among Minnesotans ages 50 and older is higher than the rate among younger age groups. In 2013, the total suicide rate among all people ages 50+ was 16.1 per 100,000 people (6.4 for women and 26.9 for men). The rate among those ages 50–64 was higher than the rate in the region (including Illinois, Indiana, Michigan, Ohio, and Wisconsin) and lower than the rate in the United States.

States vary in their reporting of suicides. The suicide rate is influenced by these reporting practices.


Exhibit 3. Suicide Rates in Minnesota, Region 5, and the United States, 2013



Source: Centers for Disease Control and Prevention (CDC), National Center for Health Statistics (NCHS), Underlying Cause of Death 1999-2013 on CDC WONDER Online Database, released 2015

Trends in Suicide Rates in Minnesota

Exhibit 4. Trends in Suicide Rates in Minnesota by Age Group, 2004-2013



Source: CDC, NCHS, Underlying Cause of Death 1999-2013 on CDC WONDER Online Database, released 2015

The suicide rate among Minnesotans ages 50+ fluctuated from a low of 10.9 per 100,000 in 2004 to a high of 16.4 per 100,000 in 2011. From 2004 to 2013, the rate was highest among those in the 50–64 age group.


How a state reports suicides can vary from year to year. The number of suicides is generally low, so even a small difference in reported numbers may make the rate fluctuate widely.

SUBSTANCE USE DISORDER AND SUBSTANCE USE DISORDER TREATMENT AMONG OLDER MINNESOTANS

30-Day Binge Drinking Among Older Minnesotans

Binge drinking, defined as five or more drinks for men and four or more drinks for women on a single occasion, can lead to serious health problems. Such problems include neurological damage, cardiovascular disease, liver disease, stroke, and poor control of diabetes. Binge drinkers are more likely to take risks such as driving while intoxicated and to experience falls and other accidents. Older people have a lower tolerance for alcohol. Binge drinking decreases with age and occurs more frequently among men than it does among women. As Exhibit 5 shows, 23.2 percent of Minnesota men ages 50–64 reported binge drinking in the past 30 days, while 6.5 percent of those in the 65+ group reported similar behavior.


Exhibit 5. Binge Drinking Rates in Minnesota by Age Group and Sex, 2013



Source: Behavioral Risk Factor Surveillance System (BRFSS), 2013

Illicit Drug Use Among Older Americans

Exhibit 6. Illicit Drug Use Among Older Americans, 2002–2013



Source: National Survey on Drug Use and Health (NSDUH), 2013
 Illicit drug use includes illegal drugs and prescription drugs used nonmedically. SAMHSA provides a list of drugs included in its survey in the NSDUH methodological summary.

Nationally, the rate of illicit drug use among older adults ages 50–64 more than doubled from 2002 to 2013. This development partially reflects the entrance into this age group of the baby boom cohort, whose rates of illicit drug use have been higher than those of older cohorts. Although state-specific data are not available, the *Minnesota Behavioral Health Barometer* is available for download from the Substance Abuse and Mental Health Services Administration (SAMHSA) website (www.samhsa.gov/data/population-data-nsduh/reports?tab=33).


Admissions to Substance Use Disorder Treatment Among Older Minnesotans

In 2012, there were 7,341 admissions of Minnesotans ages 50 and older to substance use disorder (SUD) treatment in state-funded treatment programs, a rate of 387.0 per 100,000 people ages 50+. This rate was higher than the regional rate and the national average. Men made up 66.2 percent of these admissions. Of all admissions, 79.5 percent were White/Caucasian, 13.1 percent were Black/African American, and 2.1 percent were Hispanic.

The principal sources of referral to treatment among those ages 50 and older were:

Self-Referral	Criminal Justice	Other
36.5%	17.6%	45.9%


Exhibit 7. SUD Treatment Admissions of Adults Ages 50+ in Minnesota, Region 5, and the United States by Sex, 2012



Source: Treatment Episode Data Set (TEDS), 2012¹
Data include only those clients reported to SAMHSA.

SUD Treatment Admissions Among Individuals in Region 5 Ages 50+ by Insurance Type

Exhibit 8. SUD Treatment Admissions of Adults Ages 50+ in Region 5 and the United States by Insurance Type, 2012



Source: TEDS, 2012
Data include only those clients reported to SAMHSA.

States may choose not to report some of the fields to TEDS, including information on treatment admission insurance types. These data were not reported by Minnesota in 2012. Therefore, the rates for Region 5 are used instead.

In Region 5, 59.9 percent of older adult admissions to SUD treatment were uninsured, 25.9 percent had Medicaid, 10.2 percent had Medicare, and 3.9 percent had private insurance.

SUD Treatment Admissions Among Minnesotans Ages 50+ by Primary Sources of Payment

Self-Pay	Medicaid	Other
Data not available	Data not available	Data not available


Exhibit 8 and the accompanying text reflect the type of health insurance (if any) clients had at admission; the table above represents primary sources of payment.

¹ TEDS data are collected by states as a condition of the Substance Abuse Prevention and Treatment Block Grant. Guidelines suggest that states report all clients admitted to publicly financed treatment; however, states are inconsistent in applying the guidelines. States may structure and implement different quality controls over the data. For example, states may collect different categories of information to answer TEDS questions. Information is then "walked over" to TEDS definitions.

Alcohol Use Disorder Treatment Admissions Among Minnesotans Ages 50+

Alcohol was the most frequently cited substance used by older Minnesotans in publicly financed SUD treatment in 2012. It was mentioned as a substance of use in 76.6 percent of admissions among those ages 50+. This was higher than the regional and national rates.

Exhibit 9. Alcohol Use Treatment Admissions of Adults Ages 50+ in Minnesota, Region 5, and the United States, 2012




Source: TEDS, 2012
Data include only those clients reported to SAMHSA.

SUD Treatment Admissions for Non-Alcohol Substance Use

Exhibit 10. Non-Alcohol Substance Use Treatment Admissions of Adults Ages 50+ in Minnesota, Region 5, and the United States, 2012

Substances other than alcohol were cited as the primary substances of use for 23.4 percent of older adult admissions to publicly funded treatment in Minnesota.



Source: TEDS, 2012
Data include only those clients reported to SAMHSA.

Drug-Related Emergency Department Visits Involving Pharmaceutical Misuse and Abuse by Older Adults

SAMHSA periodically releases reports from the Drug Abuse Warning Network (DAWN). DAWN, discontinued in 2011, consisted of a nationwide network of hospital emergency departments (EDs) primarily located in large metropolitan areas. DAWN data consist of professional reviews of ED records to determine the extent to which alcohol and other substance abuse was involved in ED visits. According to the November 25, 2010, *DAWN Report*:


- *In 2004, there were an estimated 115,803 ED visits involving pharmaceutical misuse and abuse by adults aged 50 or older; in 2008, there were 256,097 such visits, representing an increase of 121.1 percent*
- *One fifth (19.7 percent) of ED visits involving pharmaceutical misuse and abuse among older adults were made by persons aged 70 or older*
- *Among ED visits made by older adults, pain relievers were the type of pharmaceutical most commonly involved (43.5 percent), followed by drugs used to treat anxiety or insomnia (31.8 percent) and antidepressants (8.6 percent)*
- *Among patients aged 50 or older who visited the ED for pharmaceutical misuse or abuse, more than half (52.3 percent) were treated and released, and more than one third (37.5 percent) were admitted to the hospital*

Bulleted text in italics above is taken from SAMHSA's Center for Behavioral Health Statistics and Quality. (2010, November 25). *The DAWN Report: Drug-related emergency department visits involving pharmaceutical misuse and abuse by older adults*. Rockville, MD: SAMHSA.

CO-OCCURRING SUBSTANCE USE AND MENTAL DISORDERS

Older Minnesotans in SUD Treatment With Co-Occurring Mental Disorders

Exhibit 11. SUD Treatment Admissions of Adults Ages 50+ With a Co-Occurring Mental Disorder in Region 5 and the United States, 2012



Source: TEDS, 2012
Data include only those clients reported to SAMHSA.

The national literature shows a strong relationship between substance use and mental disorders. Studies show that 30–80 percent of individuals with a substance use or mental disorder also have a co-occurring disorder.


States may choose not to report some of the fields to TEDS, including information on SUD treatment admissions with a co-occurring disorder. These data were not reported by Minnesota in 2012. Therefore, Exhibit 11 shows regional and national figures only.

MENTAL HEALTH

Older Minnesotans Reporting Frequent Mental Distress

BRFSS, a household survey conducted in all 50 states and several territories, asks the following question: "Now thinking about your mental health, which includes stress, depression, and problems with emotions, for how many days during the past 30 days was your mental health not good?" Those individuals reporting 14 or more "Yes" days in response to this question experience frequent mental distress (FMD). Exhibit 12 shows that older Minnesotans experience FMD at a rate that is lower than the regional and national rates.


Exhibit 12. Individuals Reporting Frequent Mental Distress in Minnesota, Region 5, and the United States, 2013



Source: BRFSS, 2013

Older Minnesotans Reporting Frequent Mental Distress by Age Group and Sex

Exhibit 13. Minnesotans Reporting Frequent Mental Distress by Age Group and Sex, 2013



Source: BRFSS, 2013

As Exhibit 13 shows, 6.4 percent of men in the 50–64 age group reported FMD (14 days or more per 30-day period) compared with 12.6 percent of women in this age group. Men in the 65+ age group reported a higher rate of FMD than women in the same age group (4.3 percent compared with 3.3 percent). Illinois is the only other state in the region where men in the 65+ age group reported a higher rate of FMD than did women in this age group.

Other Measures of Mental Health

BRFSS collected other measures showing risk factors for mental and/or physical illness. These included:

- Social and Emotional Support (2010). BRFSS asked, "How often do you get the social and emotional support you need?" The possible responses were always, usually, sometimes, rarely, or never.
- Life Satisfaction (2010). BRFSS asked, "In general, how satisfied are you with your life?" The possible responses were very satisfied, satisfied, dissatisfied, or very dissatisfied.

Exhibit 14 presents the results of these surveys among older Minnesotans.

Exhibit 14. BRFSS Measures, 2010


Indicator	Ages 50+	Ages 50–64	Ages 65+
Rarely or never get social or emotional support	6.7%	5.0%	9.5%
Dissatisfied or very dissatisfied	4.3%	5.6%	2.2%

Source: BRFSS, 2010

Individuals With Frequent Mental Distress Report High Rates of Poor Physical Health

Older Americans who experienced FMD were more likely to report that their physical health was poor (14 days or more in the past 30-day period when physical health was "not good"). As shown in Exhibit 15, although nearly 11 percent of older Americans with no mental distress reported poor physical health, more than 50 percent of those with FMD reported poor physical health.

Exhibit 15. Individuals Ages 50+ in the United States Reporting Poor Physical Health by Level of Mental Distress, 2013




Source: BRFSS, 2013

Relationship Among Mental Distress, Diabetes, Stroke, Heart Attack, High Blood Pressure, and Coronary Disease

Older Americans who experience FMD are more likely to report that they have health problems. People with FMD were more than twice as likely to report having a stroke than those with some or no mental distress. They experienced coronary disease and heart attack at more than 1.6 times, diabetes at more than 1.4 times, and high blood pressure at nearly 1.2 times the rate of those with some or no mental distress.

Exhibit 16. Individuals Ages 50+ in the United States With Mental Distress and Serious Health Problems, 2013



Source: BRFSS, 2013

Older Minnesotans Admitted to State Mental Health Services

Approximately 5.8 percent of the people served by the Minnesota mental health system were ages 65 and older. This represents more than 11,660 adults.

Source: Center for Mental Health Services (CMHS) Uniform Reporting System (URS) Output Tables, 2014

DATA SOURCES

BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM (www.cdc.gov/brfss). BRFSS, managed by CDC, is the largest ongoing telephone health survey system in the world, tracking health conditions and risk behaviors in the United States annually since 1984. Data are collected on a monthly basis in all 50 states, the District of Columbia, and participating territories. BRFSS data are collected by local jurisdictions and reported to CDC.

CENTERS FOR DISEASE CONTROL AND PREVENTION, NATIONAL CENTER FOR HEALTH STATISTICS, UNDERLYING CAUSE OF DEATH 1999-2013 ON CDC WONDER ONLINE DATABASE, RELEASED 2015 (<http://wonder.cdc.gov/ucd-icd10.html>). The WONDER online database “contains mortality and population counts for all U.S. counties. Data are based on death certificates for U.S. residents. Each death certificate identifies a single underlying cause of death and demographic data. The number of deaths, crude death rates or age-adjusted death rates, and 95% confidence intervals and standard errors for death rates can be obtained by place of residence (total U.S., region, state and county), age group (single-year-of age, 5-year age groups, 10-year age groups and infant age groups), race, Hispanic ethnicity, gender, year, cause-of-death (4-digit ICD-10 code or group of codes), injury intent and injury mechanism, drug/alcohol induced causes and urbanization categories. Data are also available for place of death, month and week day of death, and whether an autopsy was performed.”

CENTER FOR MENTAL HEALTH SERVICES UNIFORM REPORTING SYSTEM (www.samhsa.gov/data/us_map). States that receive CMHS Block Grants are required to report aggregate data to URS. URS reports include information about utilization of mental health services and client demographic and outcome information.

NATIONAL SURVEY ON DRUG USE AND HEALTH (<https://nsduhweb.rti.org>). The NSDUH, managed by SAMHSA, is “an annual nationwide survey involving interviews with approximately 70,000 randomly selected individuals aged 12 and older.” NSDUH data are most frequently used by state planners to assess the need for SUD treatment. NSDUH data also include information about mental health conditions.

TREATMENT EPISODE DATA SET (www.samhsa.gov/data/client-level-data-teds/reports?tab=19). States that receive Substance Abuse Prevention and Treatment Block Grant funds submit individual client data to TEDS. TEDS includes both admission and discharge data sets, and some 1.5 million admissions are reported annually. TEDS includes information about utilization of SUD treatment services and client demographic and outcome information. TEDS is an admission-based system, and TEDS admissions do not represent individuals. Thus, an individual admitted to treatment twice within a calendar year would be counted as two admissions.

U.S. CENSUS BUREAU (www.census.gov/people). Two main sources of Census Bureau data were used in this report: (1) population estimates and (2) population projections.

OLDER ADULTS BEHAVIORAL HEALTH PROFILE

Ohio

OHIO'S POPULATION

Ohio Population by Age Group

Ohio is home to 11,594,163 people. Of these:

- 4,222,362 (36.4 percent) are over age 50.
- 2,527,164 (21.8 percent) are over age 60.
- 1,214,105 (10.5 percent) are over age 70.
- 481,093 (4.1 percent) are ages 80 and older.


The proportion of women rises fairly steadily in each age group, and women make up 63.5 percent of the 80+ group. The racial/ethnic composition of older Ohioans is as follows:

Race/Ethnicity of Ohioans Ages 50+

White	AI/AN	Black	Asian	NH/PI	Other	Hispanic
87.9%	0.2%	10.0%	1.2%	0.0%	0.7%	1.5%

Source: U.S. Census Bureau, 2015
AI/AN stands for American Indian and Alaska Native.
NH/PI stands for Native Hawaiian and Other Pacific Islander.


Exhibit 1. Ohio Population by Age Group, 2014



Source: U.S. Census Bureau, 2015

The Number of Older Ohioans Is Growing

Exhibit 2. Ohio Population by Age Group, 2000–2030



The proportion of Ohio's population that is 65 and older is growing while the proportion that is younger than 65 is shrinking. The U.S. Census Bureau estimates that 20.4 percent of Ohio's population will be 65 and older by the year 2030, an increase of 33.4 percent from 2015.

Projected Population in Ohio

Age Group	2015	2025	2030
Under 18	23.4%	23.0%	22.9%
18 to 44	34.8%	34.0%	33.4%
45 to 64	26.6%	24.0%	23.4%
65+	15.2%	19.0%	20.4%

Source: U.S. Census Bureau, 2005


SUICIDE AMONG OLDER OHIOANS

Ohio Suicide Rate Compared With Regional and National Rates

The suicide rate among Ohioans ages 50 and older is higher than the rate among younger age groups. In 2013, the total suicide rate among all people ages 50+ was 16.7 per 100,000 people (6.4 for women and 28.7 for men). The rate among those ages 50–64 was higher than the rate in the region (including Illinois, Indiana, Michigan, Minnesota, and Wisconsin) and lower than the rate in the United States.

States vary in their reporting of suicides. The suicide rate is influenced by these reporting practices.


Exhibit 3. Suicide Rates in Ohio, Region 5, and the United States, 2013



Source: Centers for Disease Control and Prevention (CDC), National Center for Health Statistics (NCHS), Underlying Cause of Death 1999–2013 on CDC WONDER Online Database, released 2015

Trends in Suicide Rates in Ohio

Exhibit 4. Trends in Suicide Rates in Ohio by Age Group, 2004–2013



Source: CDC, NCHS, Underlying Cause of Death 1999–2013 on CDC WONDER Online Database, released 2015

The suicide rate among Ohioans ages 50+ fluctuated from a low of 12.2 per 100,000 in 2009 to a high of 17.0 per 100,000 in 2012. From 2004 to 2013, the rate was highest among those in the 50–64 age group.


How a state reports suicides can vary from year to year. The number of suicides is generally low, so even a small difference in reported numbers may make the rate fluctuate widely.

SUBSTANCE USE DISORDER AND SUBSTANCE USE DISORDER TREATMENT AMONG OLDER OHIOANS

30-Day Binge Drinking Among Older Ohioans

Binge drinking, defined as five or more drinks for men and four or more drinks for women on a single occasion, can lead to serious health problems. Such problems include neurological damage, cardiovascular disease, liver disease, stroke, and poor control of diabetes. Binge drinkers are more likely to take risks such as driving while intoxicated and to experience falls and other accidents. Older people have a lower tolerance for alcohol. Binge drinking decreases with age and occurs more frequently among men than it does among women. As Exhibit 5 shows, 18.4 percent of Ohio men ages 50–64 reported binge drinking in the past 30 days, while 6.1 percent of those in the 65+ group reported similar behavior.


Exhibit 5. Binge Drinking Rates in Ohio by Age Group and Sex, 2013



Source: Behavioral Risk Factor Surveillance System (BRFSS), 2013

Illicit Drug Use Among Older Americans

Exhibit 6. Illicit Drug Use Among Older Americans, 2002–2013



Source: National Survey on Drug Use and Health (NSDUH), 2013

Illicit drug use includes illegal drugs and prescription drugs used nonmedically. SAMHSA provides a list of drugs included in its survey in the NSDUH methodological summary.

Nationally, the rate of illicit drug use among older adults ages 50–64 more than doubled from 2002 to 2013. This development partially reflects the entrance into this age group of the baby boom cohort, whose rates of illicit drug use have been higher than those of older cohorts. Although state-specific data are not available, the *Ohio Behavioral Health Barometer* is available for download from the Substance Abuse and Mental Health Services Administration (SAMHSA) website (www.samhsa.gov/data/population-data-nsduh/reports?tab=33).


Admissions to Substance Use Disorder Treatment Among Older Ohioans

In 2012, there were 6,077 admissions of Ohioans ages 50 and older to substance use disorder (SUD) treatment in state-funded treatment programs, a rate of 143.9 per 100,000 people ages 50+. This rate was lower than the regional rate and the national average. Men made up 69.9 percent of these admissions. Of all admissions, 63.4 percent were White/Caucasian, 35.9 percent were Black/African American, and 1.1 percent were Hispanic.

The principal sources of referral to treatment among those ages 50 and older were:

Self-Referral	Criminal Justice	Other
33.0%	40.4%	26.5%


Exhibit 7. SUD Treatment Admissions of Adults Ages 50+ in Ohio, Region 5, and the United States by Sex, 2012



Source: Treatment Episode Data Set (TEDS), 2012¹
Data include only those clients reported to SAMHSA.

SUD Treatment Admissions Among Individuals in Region 5 Ages 50+ by Insurance Type

Exhibit 8. SUD Treatment Admissions of Adults Ages 50+ in Region 5 and the United States by Insurance Type, 2012



Source: TEDS, 2012
Data include only those clients reported to SAMHSA.

States may choose not to report some of the fields to TEDS, including information on treatment admission insurance types. These data were not reported by Ohio in 2012. Therefore, the rates for Region 5 are used instead.

In Region 5, 59.9 percent of older adult admissions to SUD treatment were uninsured, 25.9 percent had Medicaid, 10.2 percent had Medicare, and 3.9 percent had private insurance.

SUD Treatment Admissions Among Ohioans Ages 50+ by Primary Sources of Payment

Self-Pay	Medicaid	Other
Data not available	Data not available	Data not available


Exhibit 8 and the accompanying text reflect the type of health insurance (if any) clients had at admission; the table above represents primary sources of payment.

¹ TEDS data are collected by states as a condition of the Substance Abuse Prevention and Treatment Block Grant. Guidelines suggest that states report all clients admitted to publicly financed treatment; however, states are inconsistent in applying the guidelines. States may structure and implement different quality controls over the data. For example, states may collect different categories of information to answer TEDS questions. Information is then "walked over" to TEDS definitions.

Alcohol Use Disorder Treatment Admissions Among Ohioans Ages 50+

Alcohol was the most frequently cited substance used by older Ohioans in publicly financed SUD treatment in 2012. It was mentioned as a substance of use in 55.2 percent of admissions among those ages 50+. This was lower than the regional and national rates.

Exhibit 9. Alcohol Use Treatment Admissions of Adults Ages 50+ in Ohio, Region 5, and the United States, 2012




Source: TEDS, 2012
Data include only those clients reported to SAMHSA.

SUD Treatment Admissions for Non-Alcohol Substance Use

Exhibit 10. Non-Alcohol Substance Use Treatment Admissions of Adults Ages 50+ in Ohio, Region 5, and the United States, 2012

Substances other than alcohol were cited as the primary substances of use for 44.8 percent of older adult admissions to publicly funded treatment in Ohio.



Source: TEDS, 2012
Data include only those clients reported to SAMHSA.

Drug-Related Emergency Department Visits Involving Pharmaceutical Misuse and Abuse by Older Adults

SAMHSA periodically releases reports from the Drug Abuse Warning Network (DAWN). DAWN, discontinued in 2011, consisted of a nationwide network of hospital emergency departments (EDs) primarily located in large metropolitan areas. DAWN data consist of professional reviews of ED records to determine the extent to which alcohol and other substance abuse was involved in ED visits. According to the November 25, 2010, *DAWN Report*:


- *In 2004, there were an estimated 115,803 ED visits involving pharmaceutical misuse and abuse by adults aged 50 or older; in 2008, there were 256,097 such visits, representing an increase of 121.1 percent*
- *One fifth (19.7 percent) of ED visits involving pharmaceutical misuse and abuse among older adults were made by persons aged 70 or older*
- *Among ED visits made by older adults, pain relievers were the type of pharmaceutical most commonly involved (43.5 percent), followed by drugs used to treat anxiety or insomnia (31.8 percent) and antidepressants (8.6 percent)*
- *Among patients aged 50 or older who visited the ED for pharmaceutical misuse or abuse, more than half (52.3 percent) were treated and released, and more than one third (37.5 percent) were admitted to the hospital*

Bulleted text in italics above is taken from SAMHSA's Center for Behavioral Health Statistics and Quality. (2010, November 25). *The DAWN Report: Drug-related emergency department visits involving pharmaceutical misuse and abuse by older adults*. Rockville, MD: SAMHSA.

CO-OCCURRING SUBSTANCE USE AND MENTAL DISORDERS

Older Ohioans in SUD Treatment With Co-Occurring Mental Disorders

Exhibit 11. SUD Treatment Admissions of Adults Ages 50+ With a Co-Occurring Mental Disorder in Ohio, Region 5, and the United States, 2012



Source: TEDS, 2012
Data include only those clients reported to SAMHSA.

The national literature shows a strong relationship between substance use and mental disorders. Studies show that 30–80 percent of individuals with a substance use or mental disorder also have a co-occurring disorder.


Exhibit 11 shows the proportion of SUD treatment admissions of Ohioans ages 50+ with a co-occurring mental disorder. This rate is higher than the regional rate and lower than the national average. However, state reporting practices are a factor in these results.

MENTAL HEALTH

Older Ohioans Reporting Frequent Mental Distress

BRFSS, a household survey conducted in all 50 states and several territories, asks the following question: "Now thinking about your mental health, which includes stress, depression, and problems with emotions, for how many days during the past 30 days was your mental health not good?" Those individuals reporting 14 or more "Yes" days in response to this question experience frequent mental distress (FMD). Exhibit 12 shows that older Ohioans experience FMD at a rate that is higher than the regional rate and roughly similar to the national rate.


Exhibit 12. Individuals Reporting Frequent Mental Distress in Ohio, Region 5, and the United States, 2013



Source: BRFSS, 2013

Older Ohioans Reporting Frequent Mental Distress by Age Group and Sex

Exhibit 13. Ohioans Reporting Frequent Mental Distress by Age Group and Sex, 2013



Source: BRFSS, 2013

Older men in Ohio were more likely to indulge in binge drinking, but older women were more likely to report that they had FMD (14 days or more per 30-day period). As Exhibit 13 shows, 16.1 percent of women in the 50–64 age group and 8.1 percent in the 65+ age group reported FMD, while 10.3 percent of men in the 50–64 age group and 5.1 percent in the 65+ age group reported FMD.

Other Measures of Mental Health

BRFSS collected other measures showing risk factors for mental and/or physical illness. These included:

- Social and Emotional Support (2010). BRFSS asked, “How often do you get the social and emotional support you need?” The possible responses were always, usually, sometimes, rarely, or never.
- Life Satisfaction (2010). BRFSS asked, “In general, how satisfied are you with your life?” The possible responses were very satisfied, satisfied, dissatisfied, or very dissatisfied.

Exhibit 14 presents the results of these surveys among older Ohioans.

Exhibit 14. BRFSS Measures, 2010


Indicator	Ages 50+	Ages 50–64	Ages 65+
Rarely or never get social or emotional support	9.4%	7.6%	11.9%
Dissatisfied or very dissatisfied	5.8%	7.3%	3.6%

Source: BRFSS, 2010

Individuals With Frequent Mental Distress Report High Rates of Poor Physical Health

Older Americans who experienced FMD were more likely to report that their physical health was poor (14 days or more in the past 30-day period when physical health was “not good”). As shown in Exhibit 15, although nearly 11 percent of older Americans with no mental distress reported poor physical health, more than 50 percent of those with FMD reported poor physical health.

Exhibit 15. Individuals Ages 50+ in the United States Reporting Poor Physical Health by Level of Mental Distress, 2013




Source: BRFSS, 2013

Relationship Among Mental Distress, Diabetes, Stroke, Heart Attack, High Blood Pressure, and Coronary Disease

Older Americans who experience FMD are more likely to report that they have health problems. People with FMD were more than twice as likely to report having a stroke than those with some or no mental distress. They experienced coronary disease and heart attack at more than 1.6 times, diabetes at more than 1.4 times, and high blood pressure at nearly 1.2 times the rate of those with some or no mental distress.

Exhibit 16. Individuals Ages 50+ in the United States With Mental Distress and Serious Health Problems, 2013



Source: BRFSS, 2013

Older Ohioans Admitted to State Mental Health Services

Approximately 3.2 percent of the people served by the Ohio mental health system were ages 65 and older. This represents more than 12,570 adults.

Source: Center for Mental Health Services (CMHS) Uniform Reporting System (URS) Output Tables, 2014

DATA SOURCES

BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM (www.cdc.gov/brfss). BRFSS, managed by CDC, is the largest ongoing telephone health survey system in the world, tracking health conditions and risk behaviors in the United States annually since 1984. Data are collected on a monthly basis in all 50 states, the District of Columbia, and participating territories. BRFSS data are collected by local jurisdictions and reported to CDC.

CENTERS FOR DISEASE CONTROL AND PREVENTION, NATIONAL CENTER FOR HEALTH STATISTICS, UNDERLYING CAUSE OF DEATH 1999-2013 ON CDC WONDER ONLINE DATABASE, RELEASED 2015 (<http://wonder.cdc.gov/ucd-icd10.html>). The WONDER online database "contains mortality and population counts for all U.S. counties. Data are based on death certificates for U.S. residents. Each death certificate identifies a single underlying cause of death and demographic data. The number of deaths, crude death rates or age-adjusted death rates, and 95% confidence intervals and standard errors for death rates can be obtained by place of residence (total U.S., region, state and county), age group (single-year-of age, 5-year age groups, 10-year age groups and infant age groups), race, Hispanic ethnicity, gender, year, cause-of-death (4-digit ICD-10 code or group of codes), injury intent and injury mechanism, drug/alcohol induced causes and urbanization categories. Data are also available for place of death, month and week day of death, and whether an autopsy was performed."

CENTER FOR MENTAL HEALTH SERVICES UNIFORM REPORTING SYSTEM (www.samhsa.gov/data/us_map). States that receive CMHS Block Grants are required to report aggregate data to URS. URS reports include information about utilization of mental health services and client demographic and outcome information.

NATIONAL SURVEY ON DRUG USE AND HEALTH (<https://nsduhweb.rti.org>). The NSDUH, managed by SAMHSA, is "an annual nationwide survey involving interviews with approximately 70,000 randomly selected individuals aged 12 and older." NSDUH data are most frequently used by state planners to assess the need for SUD treatment. NSDUH data also include information about mental health conditions.

TREATMENT EPISODE DATA SET (www.samhsa.gov/data/client-level-data-teds/reports?tab=19). States that receive Substance Abuse Prevention and Treatment Block Grant funds submit individual client data to TEDS. TEDS includes both admission and discharge data sets, and some 1.5 million admissions are reported annually. TEDS includes information about utilization of SUD treatment services and client demographic and outcome information. TEDS is an admission-based system, and TEDS admissions do not represent individuals. Thus, an individual admitted to treatment twice within a calendar year would be counted as two admissions.

U.S. CENSUS BUREAU (www.census.gov/people). Two main sources of Census Bureau data were used in this report: (1) population estimates and (2) population projections.

OLDER ADULTS BEHAVIORAL HEALTH PROFILE

Wisconsin

Wisconsin

OLDER ADULTS BEHAVIORAL HEALTH PROFILE

August 2016

WISCONSIN'S POPULATION

Wisconsin Population by Age Group

Wisconsin is home to 5,757,564 people. Of these:

- 2,091,687 (36.3 percent) are over age 50.
- 1,236,999 (21.5 percent) are over age 60.
- 594,110 (10.3 percent) are over age 70.
- 241,714 (4.2 percent) are ages 80 and older.

The proportion of women rises fairly steadily in each age group, and women make up 62.6 percent of the 80+ group. The racial/ethnic composition of older Wisconsinites is as follows:

Race/Ethnicity of Wisconsinites Ages 50+


White	AI/AN	Black	Asian	NH/PI	Other	Hispanic
93.7%	0.7%	3.9%	1.2%	0.0%	0.5%	2.3%

Source: U.S. Census Bureau, 2015

AI/AN stands for American Indian and Alaska Native.

NH/PI stands for Native Hawaiian and Other Pacific Islander.


Exhibit 1. Wisconsin Population by Age Group, 2014



Source: U.S. Census Bureau, 2015

The Number of Older Wisconsinites Is Growing

Exhibit 2. Wisconsin Population by Age Group, 2000–2030



The proportion of Wisconsin's population that is 65 and older is growing while the proportion that is younger than 65 is shrinking. The U.S. Census Bureau estimates that 21.3 percent of Wisconsin's population will be 65 and older by the year 2030, an increase of 48.8 percent from 2015.

Projected Population in Wisconsin

Age Group	2015	2025	2030
Under 18	22.8%	22.6%	22.2%
18 to 44	35.1%	33.4%	32.3%
45 to 64	27.0%	24.5%	24.1%
65+	15.0%	19.4%	21.3%

Source: U.S. Census Bureau, 2005


SUICIDE AMONG OLDER WISCONSINITES

Wisconsin Suicide Rate Compared With Regional and National Rates

The suicide rate among Wisconsinites ages 50 and older is higher than the rate among younger age groups. In 2013, the total suicide rate among all people ages 50+ was 18.0 per 100,000 people (6.1 for women and 31.1 for men). The rate among those ages 50–64 was higher than both the rate in the region (including Illinois, Indiana, Michigan, Minnesota, and Ohio) and the rate in the United States.

States vary in their reporting of suicides. The suicide rate is influenced by these reporting practices.


Exhibit 3. Suicide Rates in Wisconsin, Region 5, and the United States, 2013



Source: Centers for Disease Control and Prevention (CDC), National Center for Health Statistics (NCHS), Underlying Cause of Death 1999-2013 on CDC WONDER Online Database, released 2015

Trends in Suicide Rates in Wisconsin

Exhibit 4. Trends in Suicide Rates in Wisconsin by Age Group, 2004–2013



Source: CDC, NCHS, Underlying Cause of Death 1999-2013 on CDC WONDER Online Database, released 2015

The suicide rate among Wisconsinites ages 50+ fluctuated from a low of 12.3 per 100,000 in 2005 to a high of 18.0 per 100,000 in 2013. From 2004 to 2013, the rate was generally highest among those in the 50–64 age group.


How a state reports suicides can vary from year to year. The number of suicides is generally low, so even a small difference in reported numbers may make the rate fluctuate widely.

SUBSTANCE USE DISORDER AND SUBSTANCE USE DISORDER TREATMENT AMONG OLDER WISCONSINITES

30-Day Binge Drinking Among Older Wisconsinites

Binge drinking, defined as five or more drinks for men and four or more drinks for women on a single occasion, can lead to serious health problems. Such problems include neurological damage, cardiovascular disease, liver disease, stroke, and poor control of diabetes. Binge drinkers are more likely to take risks such as driving while intoxicated and to experience falls and other accidents. Older people have a lower tolerance for alcohol. Binge drinking decreases with age and occurs more frequently among men than it does among women. As Exhibit 5 shows, 23.9 percent of Wisconsin men ages 50–64 reported binge drinking in the past 30 days, while 7.5 percent of those in the 65+ group reported similar behavior.


Exhibit 5. Binge Drinking Rates in Wisconsin by Age Group and Sex, 2013



Source: Behavioral Risk Factor Surveillance System (BRFSS), 2013

Illicit Drug Use Among Older Americans

Exhibit 6. Illicit Drug Use Among Older Americans, 2002–2013



Source: National Survey on Drug Use and Health (NSDUH), 2013
 Illicit drug use includes illegal drugs and prescription drugs used nonmedically. SAMHSA provides a list of drugs included in its survey in the NSDUH methodological summary.

Nationally, the rate of illicit drug use among older adults ages 50–64 more than doubled from 2002 to 2013. This development partially reflects the entrance into this age group of the baby boom cohort, whose rates of illicit drug use have been higher than those of older cohorts. Although state-specific data are not available, the *Wisconsin Behavioral Health Barometer* is available for download from the Substance Abuse and Mental Health Services Administration (SAMHSA) website (www.samhsa.gov/data/population-data-nsduh/reports?tab=33).


Admissions to Substance Use Disorder Treatment Among Older Wisconsinites

In 2012, there were 5,732 admissions of Wisconsinites ages 50 and older to substance use disorder (SUD) treatment in state-funded treatment programs, a rate of 274.0 per 100,000 people ages 50+. This rate was higher than the regional rate and the national average. Men made up 74.7 percent of these admissions. Of all admissions, 73.7 percent were White/Caucasian, 24.4 percent were Black/African American, and 3.1 percent were Hispanic.

The principal sources of referral to treatment among those ages 50 and older were:

Self-Referral	Criminal Justice	Other
26.0%	54.5%	19.5%


Exhibit 7. SUD Treatment Admissions of Adults Ages 50+ in Wisconsin, Region 5, and the United States by Sex, 2012



Source: Treatment Episode Data Set (TEDS), 2012¹
Data include only those clients reported to SAMHSA.

SUD Treatment Admissions Among Individuals in Region 5 Ages 50+ by Insurance Type

Exhibit 8. SUD Treatment Admissions of Adults Ages 50+ in Region 5 and the United States by Insurance Type, 2012



Source: TEDS, 2012
Data include only those clients reported to SAMHSA.

States may choose not to report some of the fields to TEDS, including information on treatment admission insurance types. These data were not reported by Wisconsin in 2012. Therefore, the rates for Region 5 are used instead.

In Region 5, 59.9 percent of older adult admissions to SUD treatment were uninsured, 25.9 percent had Medicaid, 10.2 percent had Medicare, and 3.9 percent had private insurance.

SUD Treatment Admissions Among Wisconsinites Ages 50+ by Primary Sources of Payment

Self-Pay	Medicaid	Other
Data not available	Data not available	Data not available


Exhibit 8 and the accompanying text reflect the type of health insurance (if any) clients had at admission; the table above represents primary sources of payment.

¹ TEDS data are collected by states as a condition of the Substance Abuse Prevention and Treatment Block Grant. Guidelines suggest that states report all clients admitted to publicly financed treatment; however, states are inconsistent in applying the guidelines. States may structure and implement different quality controls over the data. For example, states may collect different categories of information to answer TEDS questions. Information is then "walked over" to TEDS definitions.

Alcohol Use Disorder Treatment Admissions Among Wisconsinites Ages 50+

Alcohol was the most frequently cited substance used by older Wisconsinites in publicly financed SUD treatment in 2012. It was mentioned as a substance of use in 87.2 percent of admissions among those ages 50+. This was higher than the regional and national rates.


Exhibit 9. Alcohol Use Treatment Admissions of Adults Ages 50+ in Wisconsin, Region 5, and the United States, 2012



Source: TEDS, 2012
Data include only those clients reported to SAMHSA.

SUD Treatment Admissions for Non-Alcohol Substance Use

Exhibit 10. Non-Alcohol Substance Use Treatment Admissions of Adults Ages 50+ in Wisconsin, Region 5, and the United States, 2012



Source: TEDS, 2012
Data include only those clients reported to SAMHSA.

Substances other than alcohol were cited as the primary substances of use for 12.8 percent of older adult admissions to publicly funded treatment in Wisconsin.

Drug-Related Emergency Department Visits Involving Pharmaceutical Misuse and Abuse by Older Adults

SAMHSA periodically releases reports from the Drug Abuse Warning Network (DAWN). DAWN, discontinued in 2011, consisted of a nationwide network of hospital emergency departments (EDs) primarily located in large metropolitan areas. DAWN data consist of professional reviews of ED records to determine the extent to which alcohol and other substance abuse was involved in ED visits. According to the November 25, 2010, *DAWN Report*:


- *In 2004, there were an estimated 115,803 ED visits involving pharmaceutical misuse and abuse by adults aged 50 or older; in 2008, there were 256,097 such visits, representing an increase of 121.1 percent*
- *One fifth (19.7 percent) of ED visits involving pharmaceutical misuse and abuse among older adults were made by persons aged 70 or older*
- *Among ED visits made by older adults, pain relievers were the type of pharmaceutical most commonly involved (43.5 percent), followed by drugs used to treat anxiety or insomnia (31.8 percent) and antidepressants (8.6 percent)*
- *Among patients aged 50 or older who visited the ED for pharmaceutical misuse or abuse, more than half (52.3 percent) were treated and released, and more than one third (37.5 percent) were admitted to the hospital*

Bulleted text in italics above is taken from SAMHSA's Center for Behavioral Health Statistics and Quality. (2010, November 25). *The DAWN Report: Drug-related emergency department visits involving pharmaceutical misuse and abuse by older adults*. Rockville, MD: SAMHSA.

CO-OCCURRING SUBSTANCE USE AND MENTAL DISORDERS

Older Wisconsinites in SUD Treatment With Co-Occurring Mental Disorders

Exhibit 11. SUD Treatment Admissions of Adults Ages 50+ With a Co-Occurring Mental Disorder in Region 5 and the United States, 2012



Source: TEDS, 2012
Data include only those clients reported to SAMHSA.

The national literature shows a strong relationship between substance use and mental disorders. Studies show that 30–80 percent of individuals with a substance use or mental disorder also have a co-occurring disorder.


States may choose not to report some of the fields to TEDS, including information on SUD treatment admissions with a co-occurring disorder. These data were not reported by Wisconsin in 2012. Therefore, Exhibit 11 shows regional and national figures only.

MENTAL HEALTH

Older Wisconsinites Reporting Frequent Mental Distress

BRFSS, a household survey conducted in all 50 states and several territories, asks the following question: "Now thinking about your mental health, which includes stress, depression, and problems with emotions, for how many days during the past 30 days was your mental health not good?" Those individuals reporting 14 or more "Yes" days in response to this question experience frequent mental distress (FMD). Exhibit 12 shows that older Wisconsinites experience FMD at a rate that is lower than the regional and national rates.


Exhibit 12. Individuals Reporting Frequent Mental Distress in Wisconsin, Region 5, and the United States, 2013



Source: BRFSS, 2013

Older Wisconsinites Reporting Frequent Mental Distress by Age Group and Sex

Exhibit 13. Wisconsinites Reporting Frequent Mental Distress by Age Group and Sex, 2013



Source: BRFSS, 2013

Older men in Wisconsin were more likely to indulge in binge drinking, but older women were more likely to report that they had FMD (14 days or more per 30-day period). As Exhibit 13 shows, 14.6 percent of women in the 50–64 age group and 4.7 percent in the 65+ age group reported FMD, while 8.2 percent of men in the 50–64 age group and 3.0 percent in the 65+ age group reported FMD.

Other Measures of Mental Health

BRFSS collected other measures showing risk factors for mental and/or physical illness. These included:

- Social and Emotional Support (2010). BRFSS asked, "How often do you get the social and emotional support you need?" The possible responses were always, usually, sometimes, rarely, or never.
- Life Satisfaction (2010). BRFSS asked, "In general, how satisfied are you with your life?" The possible responses were very satisfied, satisfied, dissatisfied, or very dissatisfied.

Exhibit 14 presents the results of these surveys among older Wisconsinites.

Exhibit 14. BRFSS Measures, 2010


Indicator	Ages 50+	Ages 50–64	Ages 65+
Rarely or never get social or emotional support	6.9%	4.4%	10.7%
Dissatisfied or very dissatisfied	5.3%	6.6%	3.4%

Source: BRFSS, 2010

Individuals With Frequent Mental Distress Report High Rates of Poor Physical Health

Older Americans who experienced FMD were more likely to report that their physical health was poor (14 days or more in the past 30-day period when physical health was "not good"). As shown in Exhibit 15, although nearly 11 percent of older Americans with no mental distress reported poor physical health, more than 50 percent of those with FMD reported poor physical health.

Exhibit 15. Individuals Ages 50+ in the United States Reporting Poor Physical Health by Level of Mental Distress, 2013




Source: BRFSS, 2013

Relationship Among Mental Distress, Diabetes, Stroke, Heart Attack, High Blood Pressure, and Coronary Disease

Older Americans who experience FMD are more likely to report that they have health problems. People with FMD were more than twice as likely to report having a stroke than those with some or no mental distress. They experienced coronary disease and heart attack at more than 1.6 times, diabetes at more than 1.4 times, and high blood pressure at nearly 1.2 times the rate of those with some or no mental distress.

Exhibit 16. Individuals Ages 50+ in the United States With Mental Distress and Serious Health Problems, 2013



Source: BRFSS, 2013

Older Wisconsinites Admitted to State Mental Health Services

Approximately 7.2 percent of the people served by the Wisconsin mental health system were ages 65 and older. This represents more than 7,300 adults.

Source: Center for Mental Health Services (CMHS) Uniform Reporting System (URS) Output Tables, 2014

DATA SOURCES

BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM (www.cdc.gov/brfss). BRFSS, managed by CDC, is the largest ongoing telephone health survey system in the world, tracking health conditions and risk behaviors in the United States annually since 1984. Data are collected on a monthly basis in all 50 states, the District of Columbia, and participating territories. BRFSS data are collected by local jurisdictions and reported to CDC.

CENTERS FOR DISEASE CONTROL AND PREVENTION, NATIONAL CENTER FOR HEALTH STATISTICS, UNDERLYING CAUSE OF DEATH 1999-2013 ON CDC WONDER ONLINE DATABASE, RELEASED 2015 (<http://wonder.cdc.gov/ucd-icd10.html>). The WONDER online database "contains mortality and population counts for all U.S. counties. Data are based on death certificates for U.S. residents. Each death certificate identifies a single underlying cause of death and demographic data. The number of deaths, crude death rates or age-adjusted death rates, and 95% confidence intervals and standard errors for death rates can be obtained by place of residence (total U.S., region, state and county), age group (single-year-of age, 5-year age groups, 10-year age groups and infant age groups), race, Hispanic ethnicity, gender, year, cause-of-death (4-digit ICD-10 code or group of codes), injury intent and injury mechanism, drug/alcohol induced causes and urbanization categories. Data are also available for place of death, month and week day of death, and whether an autopsy was performed."

CENTER FOR MENTAL HEALTH SERVICES UNIFORM REPORTING SYSTEM (www.samhsa.gov/data/us_map). States that receive CMHS Block Grants are required to report aggregate data to URS. URS reports include information about utilization of mental health services and client demographic and outcome information.

NATIONAL SURVEY ON DRUG USE AND HEALTH (<https://nsduhweb.rti.org>). The NSDUH, managed by SAMHSA, is "an annual nationwide survey involving interviews with approximately 70,000 randomly selected individuals aged 12 and older." NSDUH data are most frequently used by state planners to assess the need for SUD treatment. NSDUH data also include information about mental health conditions.

TREATMENT EPISODE DATA SET (www.samhsa.gov/data/client-level-data-teds/reports?tab=19). States that receive Substance Abuse Prevention and Treatment Block Grant funds submit individual client data to TEDS. TEDS includes both admission and discharge data sets, and some 1.5 million admissions are reported annually. TEDS includes information about utilization of SUD treatment services and client demographic and outcome information. TEDS is an admission-based system, and TEDS admissions do not represent individuals. Thus, an individual admitted to treatment twice within a calendar year would be counted as two admissions.

U.S. CENSUS BUREAU (www.census.gov/people). Two main sources of Census Bureau data were used in this report: (1) population estimates and (2) population projections.