



# OLDER ADULTS BEHAVIORAL HEALTH PROFILES

## REGION 6

ARKANSAS  
LOUISIANA  
NEW MEXICO  
OKLAHOMA  
TEXAS



A Behavioral Health Resource  
SAMHSA's State Technical Assistance Contract  
August 2016

# OLDER ADULTS BEHAVIORAL HEALTH PROFILES

---

## OVERVIEW

The Substance Abuse and Mental Health Services Administration recognizes the importance of behavioral health for older adults and the key role states and communities play in addressing the needs of this vital and growing population. Administrators need clear and concise information to plan, implement, and evaluate effective prevention and treatment efforts in their states.

The Older Adults Behavioral Health Profiles help states and communities identify focus areas for behavioral health plans, select population-level goals, and coordinate and target services to address priority issues. The profiles compare state trends with those in the region and the nation. State and community administrators, planners, and providers can use the information in these profiles and their own data, knowledge, and experience to establish and implement policies that improve the health and future of our valued older citizens.

Disclaimer: Data were current at the time this document was prepared.

# OLDER ADULTS BEHAVIORAL HEALTH PROFILE

---

## Arkansas

# Arkansas

## OLDER ADULTS BEHAVIORAL HEALTH PROFILE

August 2016

### ARKANSAS' POPULATION

#### Arkansas Population by Age Group

Arkansas is home to 2,966,369 people. Of these:

- 1,039,212 (35.0 percent) are over age 50.
- 641,099 (21.6 percent) are over age 60.
- 312,786 (10.5 percent) are over age 70.
- 112,772 (3.8 percent) are ages 80 and older.

The proportion of women rises fairly steadily in each age group, and women make up 62.2 percent of the 80+ group. The racial/ethnic composition of older Arkansans is as follows:

#### Race/Ethnicity of Arkansans Ages 50+


White	AI/AN	Black	Asian	NH/PI	Other	Hispanic
85.2%	0.7%	12.1%	0.9%	0.1%	1.0%	2.5%

Source: U.S. Census Bureau, 2015

AI/AN stands for American Indian and Alaska Native.

NH/PI stands for Native Hawaiian and Other Pacific Islander.


Exhibit 1. Arkansas Population by Age Group, 2014



Source: U.S. Census Bureau, 2015

#### The Number of Older Arkansans Is Growing

Exhibit 2. Arkansas Population by Age Group, 2000–2030



The proportion of Arkansas' population that is 65 and older is growing while the proportion that is younger than 65 is shrinking. The U.S. Census Bureau estimates that 20.3 percent of Arkansas' population will be 65 and older by the year 2030, an increase of 40.3 percent from 2015.

#### Projected Population in Arkansas

Age Group	2015	2025	2030
Under 18	24.3%	24.0%	24.2%
18 to 44	33.6%	32.4%	31.8%
45 to 64	26.3%	24.7%	23.7%
65+	15.8%	19.0%	20.3%

Source: U.S. Census Bureau, 2005


## SUICIDE AMONG OLDER ARKANSANS

### Arkansas Suicide Rate Compared With Regional and National Rates

The suicide rate among Arkansans ages 50 and older is higher than the rate among younger age groups. In 2013, the total suicide rate among all people ages 50+ was 21.9 per 100,000 people (8.0 for women and 37.7 for men). The rate among those ages 50–64 was higher than both the rate in the region (including Louisiana, New Mexico, Oklahoma, and Texas) and the rate in the United States.

States vary in their reporting of suicides. The suicide rate is influenced by these reporting practices.


Exhibit 3. Suicide Rates in Arkansas, Region 6, and the United States, 2013



Source: Centers for Disease Control and Prevention (CDC), National Center for Health Statistics (NCHS), Underlying Cause of Death 1999–2013 on CDC WONDER Online Database, released 2015

### Trends in Suicide Rates in Arkansas

Exhibit 4. Trends in Suicide Rates in Arkansas by Age Group, 2004–2013



Source: CDC, NCHS, Underlying Cause of Death 1999–2013 on CDC WONDER Online Database, released 2015

The suicide rate among Arkansans ages 50+ fluctuated from a low of 16.1 per 100,000 in 2007 to a high of 21.9 per 100,000 in 2013. From 2004 to 2013, the rate was generally highest among those in the 50–64 age group.


How a state reports suicides can vary from year to year. The number of suicides is generally low, so even a small difference in reported numbers may make the rate fluctuate widely.

## SUBSTANCE USE DISORDER AND SUBSTANCE USE DISORDER TREATMENT AMONG OLDER ARKANSANS

### 30-Day Binge Drinking Among Older Arkansans

Binge drinking, defined as five or more drinks for men and four or more drinks for women on a single occasion, can lead to serious health problems. Such problems include neurological damage, cardiovascular disease, liver disease, stroke, and poor control of diabetes. Binge drinkers are more likely to take risks such as driving while intoxicated and to experience falls and other accidents. Older people have a lower tolerance for alcohol. Binge drinking decreases with age and occurs more frequently among men than it does among women. As Exhibit 5 shows, 13.7 percent of Arkansas men ages 50–64 reported binge drinking in the past 30 days, while 4.2 percent of those in the 65+ group reported similar behavior.


Exhibit 5. Binge Drinking Rates in Arkansas by Age Group and Sex, 2013



Source: Behavioral Risk Factor Surveillance System (BRFSS), 2013

### Illicit Drug Use Among Older Americans

Exhibit 6. Illicit Drug Use Among Older Americans, 2002–2013



Source: National Survey on Drug Use and Health (NSDUH), 2013

Illicit drug use includes illegal drugs and prescription drugs used nonmedically. SAMHSA provides a list of drugs included in its survey in the NSDUH methodological summary.

Nationally, the rate of illicit drug use among older adults ages 50–64 more than doubled from 2002 to 2013. This development partially reflects the entrance into this age group of the baby boom cohort, whose rates of illicit drug use have been higher than those of older cohorts. Although state-specific data are not available, the *Arkansas Behavioral Health Barometer* is available for download from the Substance Abuse and Mental Health Services Administration (SAMHSA) website ([www.samhsa.gov/data/population-data-nsduh/reports?tab=33](http://www.samhsa.gov/data/population-data-nsduh/reports?tab=33)).


## Admissions to Substance Use Disorder Treatment Among Older Arkansans

In 2012, there were 1,242 admissions of Arkansans ages 50 and older to substance use disorder (SUD) treatment in state-funded treatment programs, a rate of 119.5 per 100,000 people ages 50+. This rate was higher than the regional rate and lower than the national average. Men made up 74.8 percent of these admissions. Of all admissions, 75.2 percent were White/Caucasian, 23.0 percent were Black/African American, and 0.8 percent were Hispanic.

The principal sources of referral to treatment among those ages 50 and older were:

Self-Referral	Criminal Justice	Other
47.9%	32.9%	19.2%


**Exhibit 7. SUD Treatment Admissions of Adults Ages 50+ in Arkansas, Region 6, and the United States by Sex, 2012**



Source: Treatment Episode Data Set (TEDS), 2012<sup>1</sup>  
Data include only those clients reported to SAMHSA.

## SUD Treatment Admissions Among Arkansans Ages 50+ by Insurance Type

**Exhibit 8. SUD Treatment Admissions of Adults Ages 50+ in Arkansas, Region 6, and the United States by Insurance Type, 2012**



Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

In Arkansas, 70.2 percent of older adult admissions to SUD treatment were uninsured, 6.7 percent had Medicaid, 13.4 percent had Medicare, and 9.7 percent had private insurance.

### *SUD Treatment Admissions Among Arkansans Ages 50+ by Primary Sources of Payment*

Self-Pay	Medicaid	Other
28.2%	0.2%	71.5%


Self-pay data include private insurance. Exhibit 8 and the accompanying text reflect the type of health insurance (if any) clients had at admission; the table above represents primary sources of payment.

<sup>1</sup> TEDS data are collected by states as a condition of the Substance Abuse Prevention and Treatment Block Grant. Guidelines suggest that states report all clients admitted to publicly financed treatment; however, states are inconsistent in applying the guidelines. States may structure and implement different quality controls over the data. For example, states may collect different categories of information to answer TEDS questions. Information is then "walked over" to TEDS definitions.

## Alcohol Use Disorder Treatment Admissions Among Arkansans Ages 50+

Alcohol was the most frequently cited substance used by older Arkansans in publicly financed SUD treatment in 2012. It was mentioned as a substance of use in 64.1 percent of admissions among those ages 50+. This was higher than the regional and national rates.


**Exhibit 9. Alcohol Use Treatment Admissions of Adults Ages 50+ in Arkansas, Region 6, and the United States, 2012**



Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

## SUD Treatment Admissions for Non-Alcohol Substance Use

**Exhibit 10. Non-Alcohol Substance Use Treatment Admissions of Adults Ages 50+ in Arkansas, Region 6, and the United States, 2012**



Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

Substances other than alcohol were cited as the primary substances of use for 35.9 percent of older adult admissions to publicly funded treatment in Arkansas.



## Drug-Related Emergency Department Visits Involving Pharmaceutical Misuse and Abuse by Older Adults

SAMHSA periodically releases reports from the Drug Abuse Warning Network (DAWN). DAWN, discontinued in 2011, consisted of a nationwide network of hospital emergency departments (EDs) primarily located in large metropolitan areas. DAWN data consist of professional reviews of ED records to determine the extent to which alcohol and other substance abuse was involved in ED visits. According to the November 25, 2010, *DAWN Report*:


- *In 2004, there were an estimated 115,803 ED visits involving pharmaceutical misuse and abuse by adults aged 50 or older; in 2008, there were 256,097 such visits, representing an increase of 121.1 percent*
- *One fifth (19.7 percent) of ED visits involving pharmaceutical misuse and abuse among older adults were made by persons aged 70 or older*
- *Among ED visits made by older adults, pain relievers were the type of pharmaceutical most commonly involved (43.5 percent), followed by drugs used to treat anxiety or insomnia (31.8 percent) and antidepressants (8.6 percent)*
- *Among patients aged 50 or older who visited the ED for pharmaceutical misuse or abuse, more than half (52.3 percent) were treated and released, and more than one third (37.5 percent) were admitted to the hospital*

Bulleted text in italics above is taken from SAMHSA's Center for Behavioral Health Statistics and Quality. (2010, November 25). *The DAWN Report: Drug-related emergency department visits involving pharmaceutical misuse and abuse by older adults*. Rockville, MD: SAMHSA.

## CO-OCCURRING SUBSTANCE USE AND MENTAL DISORDERS

### Older Arkansans in SUD Treatment With Co-Occurring Mental Disorders

**Exhibit 11. SUD Treatment Admissions of Adults Ages 50+ With a Co-Occurring Mental Disorder in Arkansas, Region 6, and the United States, 2012**



Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

The national literature shows a strong relationship between substance use and mental disorders. Studies show that 30–80 percent of individuals with a substance use or mental disorder also have a co-occurring disorder.


Exhibit 11 shows the proportion of SUD treatment admissions of Arkansans ages 50+ with a co-occurring mental disorder. This rate is lower than the regional rate and the national average. However, state reporting practices are a factor in these results.

## MENTAL HEALTH

### Older Arkansans Reporting Frequent Mental Distress

BRFSS, a household survey conducted in all 50 states and several territories, asks the following question: "Now thinking about your mental health, which includes stress, depression, and problems with emotions, for how many days during the past 30 days was your mental health not good?" Those individuals reporting 14 or more "Yes" days in response to this question experience frequent mental distress (FMD). Exhibit 12 shows that older Arkansans experience FMD at a rate that is higher than the regional and national rates.


**Exhibit 12. Individuals Reporting Frequent Mental Distress in Arkansas, Region 6, and the United States, 2013**



Source: BRFSS, 2013

### Older Arkansans Reporting Frequent Mental Distress by Age Group and Sex

**Exhibit 13. Arkansans Reporting Frequent Mental Distress by Age Group and Sex, 2013**



Source: BRFSS, 2013

Older men in Arkansas were more likely to indulge in binge drinking, but older women were more likely to report that they had FMD (14 days or more per 30-day period). As Exhibit 13 shows, 19.6 percent of women in the 50–64 age group and 10.8 percent in the 65+ age group reported FMD, while 13.6 percent of men in the 50–64 age group and 5.3 percent in the 65+ age group reported FMD.

## Other Measures of Mental Health

BRFSS collected other measures showing risk factors for mental and/or physical illness. These included:

- Social and Emotional Support (2010). BRFSS asked, "How often do you get the social and emotional support you need?" The possible responses were always, usually, sometimes, rarely, or never.
- Life Satisfaction (2010). BRFSS asked, "In general, how satisfied are you with your life?" The possible responses were very satisfied, satisfied, dissatisfied, or very dissatisfied.

Exhibit 14 presents the results of these surveys among older Arkansans.

**Exhibit 14. BRFSS Measures, 2010**


Indicator	Ages 50+	Ages 50–64	Ages 65+
Rarely or never get social or emotional support	9.3%	7.7%	11.6%
Dissatisfied or very dissatisfied	4.5%	5.5%	3.2%

Source: BRFSS, 2010

## Individuals With Frequent Mental Distress Report High Rates of Poor Physical Health

Older Americans who experienced FMD were more likely to report that their physical health was poor (14 days or more in the past 30-day period when physical health was "not good"). As shown in Exhibit 15, although nearly 11 percent of older Americans with no mental distress reported poor physical health, more than 50 percent of those with FMD reported poor physical health.

**Exhibit 15. Individuals Ages 50+ in the United States Reporting Poor Physical Health by Level of Mental Distress, 2013**




Source: BRFSS, 2013

## Relationship Among Mental Distress, Diabetes, Stroke, Heart Attack, High Blood Pressure, and Coronary Disease

Older Americans who experience FMD are more likely to report that they have health problems. People with FMD were more than twice as likely to report having a stroke than those with some or no mental distress. They experienced coronary disease and heart attack at more than 1.6 times, diabetes at more than 1.4 times, and high blood pressure at nearly 1.2 times the rate of those with some or no mental distress.

**Exhibit 16. Individuals Ages 50+ in the United States With Mental Distress and Serious Health Problems, 2013**



Source: BRFSS, 2013

## Older Arkansans Admitted to State Mental Health Services

Approximately 3.3 percent of the people served by the Arkansas mental health system were ages 65 and older. This represents more than 2,240 adults.

Source: Center for Mental Health Services (CMHS) Uniform Reporting System (URS) Output Tables, 2014

## DATA SOURCES

**BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM** ([www.cdc.gov/brfss](http://www.cdc.gov/brfss)). BRFSS, managed by CDC, is the largest ongoing telephone health survey system in the world, tracking health conditions and risk behaviors in the United States annually since 1984. Data are collected on a monthly basis in all 50 states, the District of Columbia, and participating territories. BRFSS data are collected by local jurisdictions and reported to CDC.

**CENTERS FOR DISEASE CONTROL AND PREVENTION, NATIONAL CENTER FOR HEALTH STATISTICS, UNDERLYING CAUSE OF DEATH 1999-2013 ON CDC WONDER ONLINE DATABASE, RELEASED 2015** (<http://wonder.cdc.gov/ucd-icd10.html>). The WONDER online database "contains mortality and population counts for all U.S. counties. Data are based on death certificates for U.S. residents. Each death certificate identifies a single underlying cause of death and demographic data. The number of deaths, crude death rates or age-adjusted death rates, and 95% confidence intervals and standard errors for death rates can be obtained by place of residence (total U.S., region, state and county), age group (single-year-of age, 5-year age groups, 10-year age groups and infant age groups), race, Hispanic ethnicity, gender, year, cause-of-death (4-digit ICD-10 code or group of codes), injury intent and injury mechanism, drug/alcohol induced causes and urbanization categories. Data are also available for place of death, month and week day of death, and whether an autopsy was performed."

**CENTER FOR MENTAL HEALTH SERVICES UNIFORM REPORTING SYSTEM** ([www.samhsa.gov/data/us\\_map](http://www.samhsa.gov/data/us_map)). States that receive CMHS Block Grants are required to report aggregate data to URS. URS reports include information about utilization of mental health services and client demographic and outcome information.

**NATIONAL SURVEY ON DRUG USE AND HEALTH** (<https://nsduhweb.rti.org>). The NSDUH, managed by SAMHSA, is "an annual nationwide survey involving interviews with approximately 70,000 randomly selected individuals aged 12 and older." NSDUH data are most frequently used by state planners to assess the need for SUD treatment. NSDUH data also include information about mental health conditions.

**TREATMENT EPISODE DATA SET** ([www.samhsa.gov/data/client-level-data-teds/reports?tab=19](http://www.samhsa.gov/data/client-level-data-teds/reports?tab=19)). States that receive Substance Abuse Prevention and Treatment Block Grant funds submit individual client data to TEDS. TEDS includes both admission and discharge data sets, and some 1.5 million admissions are reported annually. TEDS includes information about utilization of SUD treatment services and client demographic and outcome information. TEDS is an admission-based system, and TEDS admissions do not represent individuals. Thus, an individual admitted to treatment twice within a calendar year would be counted as two admissions.

**U.S. CENSUS BUREAU** ([www.census.gov/people](http://www.census.gov/people)). Two main sources of Census Bureau data were used in this report: (1) population estimates and (2) population projections.

# OLDER ADULTS BEHAVIORAL HEALTH PROFILE

---

## Louisiana

# Louisiana

## OLDER ADULTS BEHAVIORAL HEALTH PROFILE

August 2016

### LOUISIANA'S POPULATION

#### Louisiana Population by Age Group

Louisiana is home to 4,649,676 people. Of these:

- 1,544,074 (33.2 percent) are over age 50.
- 904,073 (19.4 percent) are over age 60.
- 416,473 (9.0 percent) are over age 70.
- 151,994 (3.3 percent) are ages 80 and older.

The proportion of women rises fairly steadily in each age group, and women make up 63.3 percent of the 80+ group. The racial/ethnic composition of older Louisianians is as follows:

#### Race/Ethnicity of Louisianians Ages 50+


White	AI/AN	Black	Asian	NH/PI	Other	Hispanic
70.3%	0.6%	27.0%	1.3%	0.0%	0.7%	2.8%

Source: U.S. Census Bureau, 2015

AI/AN stands for American Indian and Alaska Native.

NH/PI stands for Native Hawaiian and Other Pacific Islander.


Exhibit 1. Louisiana Population by Age Group, 2014



Source: U.S. Census Bureau, 2015

#### The Number of Older Louisianians Is Growing

Exhibit 2. Louisiana Population by Age Group, 2000–2030



The proportion of Louisiana's population that is 65 and older is growing while the proportion that is younger than 65 is shrinking. The U.S. Census Bureau estimates that 19.7 percent of Louisiana's population will be 65 and older by the year 2030, an increase of 42.2 percent from 2015.

#### Projected Population in Louisiana

Age Group	2015	2025	2030
Under 18	25.2%	24.2%	23.9%
18 to 44	35.1%	34.5%	33.7%
45 to 64	25.5%	23.1%	22.7%
65+	14.2%	18.2%	19.7%

Source: U.S. Census Bureau, 2005


## SUICIDE AMONG OLDER LOUISIANIANS

### Louisiana Suicide Rate Compared With Regional and National Rates

The suicide rate among Louisianians ages 50 and older is higher than the rate among younger age groups. In 2013, the total suicide rate among all people ages 50+ was 17.4 per 100,000 people (7.1 for women and 29.3 for men). The rate among those ages 50–64 was lower than both the rate in the region (including Arkansas, New Mexico, Oklahoma, and Texas) and the rate in the United States.

States vary in their reporting of suicides. The suicide rate is influenced by these reporting practices.


Exhibit 3. Suicide Rates in Louisiana, Region 6, and the United States, 2013



Source: Centers for Disease Control and Prevention (CDC), National Center for Health Statistics (NCHS), Underlying Cause of Death 1999–2013 on CDC WONDER Online Database, released 2015

### Trends in Suicide Rates in Louisiana

Exhibit 4. Trends in Suicide Rates in Louisiana by Age Group, 2004–2013



Source: CDC, NCHS, Underlying Cause of Death 1999–2013 on CDC WONDER Online Database, released 2015

The suicide rate among Louisianians ages 50+ fluctuated from a low of 12.7 per 100,000 in 2006 to a high of 17.4 per 100,000 in 2013. From 2004 to 2013, the rate was generally highest among those in the 50–64 age group.

How a state reports suicides can vary from year to year. The number of suicides is generally low, so even a small difference in reported numbers may make the rate fluctuate widely.




## SUBSTANCE USE DISORDER AND SUBSTANCE USE DISORDER TREATMENT AMONG OLDER LOUISIANIANS

### 30-Day Binge Drinking Among Older Louisianians

Binge drinking, defined as five or more drinks for men and four or more drinks for women on a single occasion, can lead to serious health problems. Such problems include neurological damage, cardiovascular disease, liver disease, stroke, and poor control of diabetes. Binge drinkers are more likely to take risks such as driving while intoxicated and to experience falls and other accidents. Older people have a lower tolerance for alcohol. Binge drinking decreases with age and occurs more frequently among men than it does among women. As Exhibit 5 shows, 18.8 percent of Louisiana men ages 50–64 reported binge drinking in the past 30 days, while 7.7 percent of those in the 65+ group reported similar behavior.


Exhibit 5. Binge Drinking Rates in Louisiana by Age Group and Sex, 2013



Source: Behavioral Risk Factor Surveillance System (BRFSS), 2013

### Illicit Drug Use Among Older Americans

Exhibit 6. Illicit Drug Use Among Older Americans, 2002–2013



Source: National Survey on Drug Use and Health (NSDUH), 2013  
Illicit drug use includes illegal drugs and prescription drugs used nonmedically. SAMHSA provides a list of drugs included in its survey in the NSDUH methodological summary.

Nationally, the rate of illicit drug use among older adults ages 50–64 more than doubled from 2002 to 2013. This development partially reflects the entrance into this age group of the baby boom cohort, whose rates of illicit drug use have been higher than those of older cohorts. Although state-specific data are not available, the *Louisiana Behavioral Health Barometer* is available for download from the Substance Abuse and Mental Health Services Administration (SAMHSA) website ([www.samhsa.gov/data/population-data-nsduh/reports?tab=33](http://www.samhsa.gov/data/population-data-nsduh/reports?tab=33)).


## Admissions to Substance Use Disorder Treatment Among Older Louisianians

In 2012, there were 1,897 admissions of Louisianians ages 50 and older to substance use disorder (SUD) treatment in state-funded treatment programs, a rate of 122.9 per 100,000 people ages 50+. This rate was higher than the regional rate and lower than the national average. Men made up 69.5 percent of these admissions. Of all admissions, 55.1 percent were White/Caucasian, 43.2 percent were Black/African American, and 1.6 percent were Hispanic.

The principal sources of referral to treatment among those ages 50 and older were:

Self-Referral	Criminal Justice	Other
48.8%	14.7%	36.4%


**Exhibit 7. SUD Treatment Admissions of Adults Ages 50+ in Louisiana, Region 6, and the United States by Sex, 2012**



Source: Treatment Episode Data Set (TEDS), 2012<sup>1</sup>  
Data include only those clients reported to SAMHSA.

## SUD Treatment Admissions Among Louisianians Ages 50+ by Insurance Type

**Exhibit 8. SUD Treatment Admissions of Adults Ages 50+ in Louisiana, Region 6, and the United States by Insurance Type, 2012**



Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

In Louisiana, 77.0 percent of older adult admissions to SUD treatment were uninsured, 10.5 percent had Medicaid, 6.0 percent had Medicare, and 6.5 percent had private insurance.

### *SUD Treatment Admissions Among Louisianians Ages 50+ by Primary Sources of Payment*

Self-Pay	Medicaid	Other
12.4%	3.7%	83.9%


Self-pay data include private insurance. Exhibit 8 and the accompanying text reflect the type of health insurance (if any) clients had at admission; the table above represents primary sources of payment.

<sup>1</sup> TEDS data are collected by states as a condition of the Substance Abuse Prevention and Treatment Block Grant. Guidelines suggest that states report all clients admitted to publicly financed treatment; however, states are inconsistent in applying the guidelines. States may structure and implement different quality controls over the data. For example, states may collect different categories of information to answer TEDS questions. Information is then "walked over" to TEDS definitions.

## Alcohol Use Disorder Treatment Admissions Among Louisianians Ages 50+

Alcohol was the most frequently cited substance used by older Louisianians in publicly financed SUD treatment in 2012. It was mentioned as a substance of use in 48.1 percent of admissions among those ages 50+. This was lower than the regional and national rates.

**Exhibit 9. Alcohol Use Treatment Admissions of Adults Ages 50+ in Louisiana, Region 6, and the United States, 2012**




Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

## SUD Treatment Admissions for Non-Alcohol Substance Use

**Exhibit 10. Non-Alcohol Substance Use Treatment Admissions of Adults Ages 50+ in Louisiana, Region 6, and the United States, 2012**

Substances other than alcohol were cited as the primary substances of use for 51.9 percent of older adult admissions to publicly funded treatment in Louisiana.



Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

## Drug-Related Emergency Department Visits Involving Pharmaceutical Misuse and Abuse by Older Adults

SAMHSA periodically releases reports from the Drug Abuse Warning Network (DAWN). DAWN, discontinued in 2011, consisted of a nationwide network of hospital emergency departments (EDs) primarily located in large metropolitan areas. DAWN data consist of professional reviews of ED records to determine the extent to which alcohol and other substance abuse was involved in ED visits. According to the November 25, 2010, *DAWN Report*:


- *In 2004, there were an estimated 115,803 ED visits involving pharmaceutical misuse and abuse by adults aged 50 or older; in 2008, there were 256,097 such visits, representing an increase of 121.1 percent*
- *One fifth (19.7 percent) of ED visits involving pharmaceutical misuse and abuse among older adults were made by persons aged 70 or older*
- *Among ED visits made by older adults, pain relievers were the type of pharmaceutical most commonly involved (43.5 percent), followed by drugs used to treat anxiety or insomnia (31.8 percent) and antidepressants (8.6 percent)*
- *Among patients aged 50 or older who visited the ED for pharmaceutical misuse or abuse, more than half (52.3 percent) were treated and released, and more than one third (37.5 percent) were admitted to the hospital*

Bulleted text in italics above is taken from SAMHSA's Center for Behavioral Health Statistics and Quality. (2010, November 25). *The DAWN Report: Drug-related emergency department visits involving pharmaceutical misuse and abuse by older adults*. Rockville, MD: SAMHSA.

## CO-OCCURRING SUBSTANCE USE AND MENTAL DISORDERS

### Older Louisianians in SUD Treatment With Co-Occurring Mental Disorders

**Exhibit 11. SUD Treatment Admissions of Adults Ages 50+ With a Co-Occurring Mental Disorder in Louisiana, Region 6, and the United States, 2012**



Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

The national literature shows a strong relationship between substance use and mental disorders. Studies show that 30–80 percent of individuals with a substance use or mental disorder also have a co-occurring disorder.


Exhibit 11 shows the proportion of SUD treatment admissions of Louisianians ages 50+ with a co-occurring mental disorder. This rate is lower than the regional rate and the national average. However, state reporting practices are a factor in these results.

## MENTAL HEALTH

### Older Louisianians Reporting Frequent Mental Distress

BRFSS, a household survey conducted in all 50 states and several territories, asks the following question: "Now thinking about your mental health, which includes stress, depression, and problems with emotions, for how many days during the past 30 days was your mental health not good?" Those individuals reporting 14 or more "Yes" days in response to this question experience frequent mental distress (FMD). Exhibit 12 shows that older Louisianians experience FMD at a rate that is higher than the regional and national rates.


**Exhibit 12. Individuals Reporting Frequent Mental Distress in Louisiana, Region 6, and the United States, 2013**



Source: BRFSS, 2013

### Older Louisianians Reporting Frequent Mental Distress by Age Group and Sex

**Exhibit 13. Louisianians Reporting Frequent Mental Distress by Age Group and Sex, 2013**



Source: BRFSS, 2013

Older men in Louisiana were more likely to indulge in binge drinking, but older women were more likely to report that they had FMD (14 days or more per 30-day period). As Exhibit 13 shows, 16.5 percent of women in the 50–64 age group and 8.6 percent in the 65+ age group reported FMD, while 12.5 percent of men in the 50–64 age group and 6.1 percent in the 65+ age group reported FMD.

## Other Measures of Mental Health

BRFSS collected other measures showing risk factors for mental and/or physical illness. These included:

- Social and Emotional Support (2010). BRFSS asked, "How often do you get the social and emotional support you need?" The possible responses were always, usually, sometimes, rarely, or never.
- Life Satisfaction (2010). BRFSS asked, "In general, how satisfied are you with your life?" The possible responses were very satisfied, satisfied, dissatisfied, or very dissatisfied.

Exhibit 14 presents the results of these surveys among older Louisianians.

**Exhibit 14. BRFSS Measures, 2010**


Indicator	Ages 50+	Ages 50–64	Ages 65+
Rarely or never get social or emotional support	10.4%	9.3%	11.9%
Dissatisfied or very dissatisfied	5.0%	6.0%	3.5%

Source: BRFSS, 2010

## Individuals With Frequent Mental Distress Report High Rates of Poor Physical Health

Older Americans who experienced FMD were more likely to report that their physical health was poor (14 days or more in the past 30-day period when physical health was "not good"). As shown in Exhibit 15, although nearly 11 percent of older Americans with no mental distress reported poor physical health, more than 50 percent of those with FMD reported poor physical health.

**Exhibit 15. Individuals Ages 50+ in the United States Reporting Poor Physical Health by Level of Mental Distress, 2013**




Source: BRFSS, 2013

## Relationship Among Mental Distress, Diabetes, Stroke, Heart Attack, High Blood Pressure, and Coronary Disease

Older Americans who experience FMD are more likely to report that they have health problems. People with FMD were more than twice as likely to report having a stroke than those with some or no mental distress. They experienced coronary disease and heart attack at more than 1.6 times, diabetes at more than 1.4 times, and high blood pressure at nearly 1.2 times the rate of those with some or no mental distress.

**Exhibit 16. Individuals Ages 50+ in the United States With Mental Distress and Serious Health Problems, 2013**



Source: BRFSS, 2013

## Older Louisianians Admitted to State Mental Health Services

Approximately 2.7 percent of the people served by the Louisiana mental health system were ages 65 and older. This represents more than 1,320 adults.

Source: Center for Mental Health Services (CMHS) Uniform Reporting System (URS) Output Tables, 2014

## DATA SOURCES

**BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM** ([www.cdc.gov/brfss](http://www.cdc.gov/brfss)). BRFSS, managed by CDC, is the largest ongoing telephone health survey system in the world, tracking health conditions and risk behaviors in the United States annually since 1984. Data are collected on a monthly basis in all 50 states, the District of Columbia, and participating territories. BRFSS data are collected by local jurisdictions and reported to CDC.

**CENTERS FOR DISEASE CONTROL AND PREVENTION, NATIONAL CENTER FOR HEALTH STATISTICS, UNDERLYING CAUSE OF DEATH 1999-2013 ON CDC WONDER ONLINE DATABASE, RELEASED 2015** (<http://wonder.cdc.gov/ucd-icd10.html>). The WONDER online database "contains mortality and population counts for all U.S. counties. Data are based on death certificates for U.S. residents. Each death certificate identifies a single underlying cause of death and demographic data. The number of deaths, crude death rates or age-adjusted death rates, and 95% confidence intervals and standard errors for death rates can be obtained by place of residence (total U.S., region, state and county), age group (single-year-of age, 5-year age groups, 10-year age groups and infant age groups), race, Hispanic ethnicity, gender, year, cause-of-death (4-digit ICD-10 code or group of codes), injury intent and injury mechanism, drug/alcohol induced causes and urbanization categories. Data are also available for place of death, month and week day of death, and whether an autopsy was performed."

**CENTER FOR MENTAL HEALTH SERVICES UNIFORM REPORTING SYSTEM** ([www.samhsa.gov/data/us\\_map](http://www.samhsa.gov/data/us_map)). States that receive CMHS Block Grants are required to report aggregate data to URS. URS reports include information about utilization of mental health services and client demographic and outcome information.

**NATIONAL SURVEY ON DRUG USE AND HEALTH** (<https://nsduhweb.rti.org>). The NSDUH, managed by SAMHSA, is "an annual nationwide survey involving interviews with approximately 70,000 randomly selected individuals aged 12 and older." NSDUH data are most frequently used by state planners to assess the need for SUD treatment. NSDUH data also include information about mental health conditions.

**TREATMENT EPISODE DATA SET** ([www.samhsa.gov/data/client-level-data-teds/reports?tab=19](http://www.samhsa.gov/data/client-level-data-teds/reports?tab=19)). States that receive Substance Abuse Prevention and Treatment Block Grant funds submit individual client data to TEDS. TEDS includes both admission and discharge data sets, and some 1.5 million admissions are reported annually. TEDS includes information about utilization of SUD treatment services and client demographic and outcome information. TEDS is an admission-based system, and TEDS admissions do not represent individuals. Thus, an individual admitted to treatment twice within a calendar year would be counted as two admissions.

**U.S. CENSUS BUREAU** ([www.census.gov/people](http://www.census.gov/people)). Two main sources of Census Bureau data were used in this report: (1) population estimates and (2) population projections.



# OLDER ADULTS BEHAVIORAL HEALTH PROFILE

---

## New Mexico

# New Mexico

## OLDER ADULTS BEHAVIORAL HEALTH PROFILE

August 2016

### NEW MEXICO'S POPULATION

#### New Mexico Population by Age Group

New Mexico is home to 2,085,572 people. Of these:

- 730,227 (35.0 percent) are over age 50.
- 447,604 (21.5 percent) are over age 60.
- 209,966 (10.1 percent) are over age 70.
- 75,957 (3.6 percent) are ages 80 and older.

The proportion of women rises fairly steadily in each age group, and women make up 59.3 percent of the 80+ group. The racial/ethnic composition of older New Mexicans is as follows:

#### Race/Ethnicity of New Mexicans Ages 50+


White	AI/AN	Black	Asian	NH/PI	Other	Hispanic
88.3%	7.2%	1.8%	1.4%	0.1%	1.2%	35.7%

Source: U.S. Census Bureau, 2015

AI/AN stands for American Indian and Alaska Native.

NH/PI stands for Native Hawaiian and Other Pacific Islander.


Exhibit 1. New Mexico Population by Age Group, 2014



Source: U.S. Census Bureau, 2015

#### The Number of Older New Mexicans Is Growing

Exhibit 2. New Mexico Population by Age Group, 2000–2030



The proportion of New Mexico's population that is 65 and older is growing while the proportion that is younger than 65 is shrinking. The U.S. Census Bureau estimates that 26.4 percent of New Mexico's population will be 65 and older by the year 2030, an increase of 61.6 percent from 2015.

#### Projected Population in New Mexico

Age Group	2015	2025	2030
Under 18	23.7%	22.7%	21.7%
18 to 44	32.0%	29.7%	28.6%
45 to 64	27.4%	23.9%	23.3%
65+	16.8%	23.6%	26.4%

Source: U.S. Census Bureau, 2005


## SUICIDE AMONG OLDER NEW MEXICANS

### New Mexico Suicide Rate Compared With Regional and National Rates

The suicide rate among New Mexicans ages 50 and older is higher than the rate among younger age groups. In 2013, the total suicide rate among all people ages 50+ was 27.5 per 100,000 people (14.4 for women and 42.3 for men). The rate among those ages 50–64 was higher than both the rate in the region (including Arkansas, Louisiana, Oklahoma, and Texas) and the rate in the United States.

States vary in their reporting of suicides. The suicide rate is influenced by these reporting practices.


**Exhibit 3. Suicide Rates in New Mexico, Region 6, and the United States, 2013**



Source: Centers for Disease Control and Prevention (CDC), National Center for Health Statistics (NCHS), Underlying Cause of Death 1999-2013 on CDC WONDER Online Database, released 2015

### Trends in Suicide Rates in New Mexico

**Exhibit 4. Trends in Suicide Rates in New Mexico by Age Group, 2004–2013**



Source: CDC, NCHS, Underlying Cause of Death 1999-2013 on CDC WONDER Online Database, released 2015

The suicide rate among New Mexicans ages 50+ fluctuated from a low of 20.3 per 100,000 in 2006 to a high of 27.5 per 100,000 in 2013. From 2004 to 2013, the rate was generally highest among those in the 50–64 age group.


How a state reports suicides can vary from year to year. The number of suicides is generally low, so even a small difference in reported numbers may make the rate fluctuate widely.

## SUBSTANCE USE DISORDER AND SUBSTANCE USE DISORDER TREATMENT AMONG OLDER NEW MEXICANS

### 30-Day Binge Drinking Among Older New Mexicans

Binge drinking, defined as five or more drinks for men and four or more drinks for women on a single occasion, can lead to serious health problems. Such problems include neurological damage, cardiovascular disease, liver disease, stroke, and poor control of diabetes. Binge drinkers are more likely to take risks such as driving while intoxicated and to experience falls and other accidents. Older people have a lower tolerance for alcohol. Binge drinking decreases with age and occurs more frequently among men than it does among women. As Exhibit 5 shows, 15.2 percent of New Mexico men ages 50–64 reported binge drinking in the past 30 days, while 5.9 percent of those in the 65+ group reported similar behavior.


Exhibit 5. Binge Drinking Rates in New Mexico by Age Group and Sex, 2013



Source: Behavioral Risk Factor Surveillance System (BRFSS), 2013

### Illicit Drug Use Among Older Americans

Exhibit 6. Illicit Drug Use Among Older Americans, 2002–2013



Source: National Survey on Drug Use and Health (NSDUH), 2013  
 Illicit drug use includes illegal drugs and prescription drugs used nonmedically. SAMHSA provides a list of drugs included in its survey in the NSDUH methodological summary.

Nationally, the rate of illicit drug use among older adults ages 50–64 more than doubled from 2002 to 2013. This development partially reflects the entrance into this age group of the baby boom cohort, whose rates of illicit drug use have been higher than those of older cohorts. Although state-specific data are not available, the *New Mexico Behavioral Health Barometer* is available for download from the Substance Abuse and Mental Health Services Administration (SAMHSA) website ([www.samhsa.gov/data/population-data-nsduh/reports?tab=33](http://www.samhsa.gov/data/population-data-nsduh/reports?tab=33)).


## Admissions to Substance Use Disorder Treatment Among Older New Mexicans

In 2012, there were 948 admissions of New Mexicans ages 50 and older to substance use disorder (SUD) treatment in state-funded treatment programs, a rate of 129.8 per 100,000 people ages 50+. This rate was higher than the regional rate and lower than the national average. Men made up 70.1 percent of these admissions. Of all admissions, 78.6 percent were White/Caucasian, 1.8 percent were Black/African American, and 47.7 percent were Hispanic.

The principal sources of referral to treatment among those ages 50 and older were:

Self-Referral	Criminal Justice	Other
34.3%	41.4%	24.3%


**Exhibit 7. SUD Treatment Admissions of Adults Ages 50+ in New Mexico, Region 6, and the United States by Sex, 2012**



Source: Treatment Episode Data Set (TEDS), 2012<sup>1</sup>  
Data include only those clients reported to SAMHSA.

## SUD Treatment Admissions Among New Mexicans Ages 50+ by Insurance Type

**Exhibit 8. SUD Treatment Admissions of Adults Ages 50+ in New Mexico, Region 6, and the United States by Insurance Type, 2012**



Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

In New Mexico, 71.3 percent of older adult admissions to SUD treatment were uninsured, 1.9 percent had Medicaid, 21.8 percent had Medicare, and 5.0 percent had private insurance.

### *SUD Treatment Admissions Among New Mexicans Ages 50+ by Primary Sources of Payment*

Self-Pay	Medicaid	Other
15.5%	3.6%	80.8%


Self-pay data include private insurance. Exhibit 8 and the accompanying text reflect the type of health insurance (if any) clients had at admission; the table above represents primary sources of payment.

<sup>1</sup> TEDS data are collected by states as a condition of the Substance Abuse Prevention and Treatment Block Grant. Guidelines suggest that states report all clients admitted to publicly financed treatment; however, states are inconsistent in applying the guidelines. States may structure and implement different quality controls over the data. For example, states may collect different categories of information to answer TEDS questions. Information is then "walked over" to TEDS definitions.

## Alcohol Use Disorder Treatment Admissions Among New Mexicans Ages 50+

Alcohol was the most frequently cited substance used by older New Mexicans in publicly financed SUD treatment in 2012. It was mentioned as a substance of use in 74.1 percent of admissions among those ages 50+. This was higher than the regional and national rates.

**Exhibit 9. Alcohol Use Treatment Admissions of Adults Ages 50+ in New Mexico, Region 6, and the United States, 2012**




Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

## SUD Treatment Admissions for Non-Alcohol Substance Use

**Exhibit 10. Non-Alcohol Substance Use Treatment Admissions of Adults Ages 50+ in New Mexico, Region 6, and the United States, 2012**

Substances other than alcohol were cited as the primary substances of use for 25.9 percent of older adult admissions to publicly funded treatment in New Mexico.



Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

## Drug-Related Emergency Department Visits Involving Pharmaceutical Misuse and Abuse by Older Adults

SAMHSA periodically releases reports from the Drug Abuse Warning Network (DAWN). DAWN, discontinued in 2011, consisted of a nationwide network of hospital emergency departments (EDs) primarily located in large metropolitan areas. DAWN data consist of professional reviews of ED records to determine the extent to which alcohol and other substance abuse was involved in ED visits. According to the November 25, 2010, *DAWN Report*:


- *In 2004, there were an estimated 115,803 ED visits involving pharmaceutical misuse and abuse by adults aged 50 or older; in 2008, there were 256,097 such visits, representing an increase of 121.1 percent*
- *One fifth (19.7 percent) of ED visits involving pharmaceutical misuse and abuse among older adults were made by persons aged 70 or older*
- *Among ED visits made by older adults, pain relievers were the type of pharmaceutical most commonly involved (43.5 percent), followed by drugs used to treat anxiety or insomnia (31.8 percent) and antidepressants (8.6 percent)*
- *Among patients aged 50 or older who visited the ED for pharmaceutical misuse or abuse, more than half (52.3 percent) were treated and released, and more than one third (37.5 percent) were admitted to the hospital*

Bulleted text in italics above is taken from SAMHSA's Center for Behavioral Health Statistics and Quality. (2010, November 25). *The DAWN Report: Drug-related emergency department visits involving pharmaceutical misuse and abuse by older adults*. Rockville, MD: SAMHSA.

## CO-OCCURRING SUBSTANCE USE AND MENTAL DISORDERS

### Older New Mexicans in SUD Treatment With Co-Occurring Mental Disorders

**Exhibit 11. SUD Treatment Admissions of Adults Ages 50+ With a Co-Occurring Mental Disorder in New Mexico, Region 6, and the United States, 2012**



Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

The national literature shows a strong relationship between substance use and mental disorders. Studies show that 30–80 percent of individuals with a substance use or mental disorder also have a co-occurring disorder.


Exhibit 11 shows the proportion of SUD treatment admissions of New Mexicans ages 50+ with a co-occurring mental disorder. This rate is lower than the regional rate and the national average. However, state reporting practices are a factor in these results.

## MENTAL HEALTH

### Older New Mexicans Reporting Frequent Mental Distress

BRFSS, a household survey conducted in all 50 states and several territories, asks the following question: "Now thinking about your mental health, which includes stress, depression, and problems with emotions, for how many days during the past 30 days was your mental health not good?" Those individuals reporting 14 or more "Yes" days in response to this question experience frequent mental distress (FMD). Exhibit 12 shows that New Mexicans in the 50–64 age group experience FMD at a rate that is higher than the regional and national rates while those in the 65+ age group experience it at a rate that is lower than the regional and national rates.


**Exhibit 12. Individuals Reporting Frequent Mental Distress in New Mexico, Region 6, and the United States, 2013**



Source: BRFSS, 2013

### Older New Mexicans Reporting Frequent Mental Distress by Age Group and Sex

**Exhibit 13. New Mexicans Reporting Frequent Mental Distress by Age Group and Sex, 2013**



Source: BRFSS, 2013

As Exhibit 13 shows, 13.7 percent of men in the 50–64 age group reported FMD (14 days or more per 30-day period) compared with 16.4 percent of women in this age group. In the 65+ age group, men reported a higher rate of FMD than did women (6.7 percent compared with 6.5 percent), in contrast to the other states in the region, where women in this age group were more likely to report FMD.



## Other Measures of Mental Health

BRFSS collected other measures showing risk factors for mental and/or physical illness. These included:

- Social and Emotional Support (2010). BRFSS asked, "How often do you get the social and emotional support you need?" The possible responses were always, usually, sometimes, rarely, or never.
- Life Satisfaction (2010). BRFSS asked, "In general, how satisfied are you with your life?" The possible responses were very satisfied, satisfied, dissatisfied, or very dissatisfied.

Exhibit 14 presents the results of these surveys among older New Mexicans.

**Exhibit 14. BRFSS Measures, 2010**


Indicator	Ages 50+	Ages 50–64	Ages 65+
Rarely or never get social or emotional support	9.7%	8.7%	11.1%
Dissatisfied or very dissatisfied	5.0%	6.0%	3.7%

Source: BRFSS, 2010

## Individuals With Frequent Mental Distress Report High Rates of Poor Physical Health

Older Americans who experienced FMD were more likely to report that their physical health was poor (14 days or more in the past 30-day period when physical health was "not good"). As shown in Exhibit 15, although nearly 11 percent of older Americans with no mental distress reported poor physical health, more than 50 percent of those with FMD reported poor physical health.

**Exhibit 15. Individuals Ages 50+ in the United States Reporting Poor Physical Health by Level of Mental Distress, 2013**




Source: BRFSS, 2013

## Relationship Among Mental Distress, Diabetes, Stroke, Heart Attack, High Blood Pressure, and Coronary Disease

Older Americans who experience FMD are more likely to report that they have health problems. People with FMD were more than twice as likely to report having a stroke than those with some or no mental distress. They experienced coronary disease and heart attack at more than 1.6 times, diabetes at more than 1.4 times, and high blood pressure at nearly 1.2 times the rate of those with some or no mental distress.

**Exhibit 16. Individuals Ages 50+ in the United States With Mental Distress and Serious Health Problems, 2013**



Source: BRFSS, 2013

## Older New Mexicans Admitted to State Mental Health Services

Approximately 3.2 percent of the people served by the New Mexico mental health system were ages 65 and older. This represents more than 2,940 adults.

Source: Center for Mental Health Services (CMHS) Uniform Reporting System (URS) Output Tables, 2014

## DATA SOURCES

**BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM** ([www.cdc.gov/brfss](http://www.cdc.gov/brfss)). BRFSS, managed by CDC, is the largest ongoing telephone health survey system in the world, tracking health conditions and risk behaviors in the United States annually since 1984. Data are collected on a monthly basis in all 50 states, the District of Columbia, and participating territories. BRFSS data are collected by local jurisdictions and reported to CDC.

**CENTERS FOR DISEASE CONTROL AND PREVENTION, NATIONAL CENTER FOR HEALTH STATISTICS, UNDERLYING CAUSE OF DEATH 1999-2013 ON CDC WONDER ONLINE DATABASE, RELEASED 2015** (<http://wonder.cdc.gov/ucd-icd10.html>). The WONDER online database "contains mortality and population counts for all U.S. counties. Data are based on death certificates for U.S. residents. Each death certificate identifies a single underlying cause of death and demographic data. The number of deaths, crude death rates or age-adjusted death rates, and 95% confidence intervals and standard errors for death rates can be obtained by place of residence (total U.S., region, state and county), age group (single-year-of age, 5-year age groups, 10-year age groups and infant age groups), race, Hispanic ethnicity, gender, year, cause-of-death (4-digit ICD-10 code or group of codes), injury intent and injury mechanism, drug/alcohol induced causes and urbanization categories. Data are also available for place of death, month and week day of death, and whether an autopsy was performed."

**CENTER FOR MENTAL HEALTH SERVICES UNIFORM REPORTING SYSTEM** ([www.samhsa.gov/data/us\\_map](http://www.samhsa.gov/data/us_map)). States that receive CMHS Block Grants are required to report aggregate data to URS. URS reports include information about utilization of mental health services and client demographic and outcome information.

**NATIONAL SURVEY ON DRUG USE AND HEALTH** (<https://nsduhweb.rti.org>). The NSDUH, managed by SAMHSA, is "an annual nationwide survey involving interviews with approximately 70,000 randomly selected individuals aged 12 and older." NSDUH data are most frequently used by state planners to assess the need for SUD treatment. NSDUH data also include information about mental health conditions.

**TREATMENT EPISODE DATA SET** ([www.samhsa.gov/data/client-level-data-teds/reports?tab=19](http://www.samhsa.gov/data/client-level-data-teds/reports?tab=19)). States that receive Substance Abuse Prevention and Treatment Block Grant funds submit individual client data to TEDS. TEDS includes both admission and discharge data sets, and some 1.5 million admissions are reported annually. TEDS includes information about utilization of SUD treatment services and client demographic and outcome information. TEDS is an admission-based system, and TEDS admissions do not represent individuals. Thus, an individual admitted to treatment twice within a calendar year would be counted as two admissions.

**U.S. CENSUS BUREAU** ([www.census.gov/people](http://www.census.gov/people)). Two main sources of Census Bureau data were used in this report: (1) population estimates and (2) population projections.

# OLDER ADULTS BEHAVIORAL HEALTH PROFILE

---

# Oklahoma

# Oklahoma

## OLDER ADULTS BEHAVIORAL HEALTH PROFILE

August 2016

### OKLAHOMA'S POPULATION

#### Oklahoma Population by Age Group

Oklahoma is home to 3,878,051 people. Of these:

- 1,300,187 (33.5 percent) are over age 50.
- 783,474 (20.2 percent) are over age 60.
- 378,264 (9.8 percent) are over age 70.
- 139,263 (3.6 percent) are ages 80 and older.


The proportion of women rises fairly steadily in each age group, and women make up 61.7 percent of the 80+ group. The racial/ethnic composition of older Oklahomans is as follows:

#### Race/Ethnicity of Oklahomans Ages 50+

White	AI/AN	Black	Asian	NH/PI	Other	Hispanic
83.0%	6.5%	5.9%	1.4%	0.1%	3.1%	3.7%

Source: U.S. Census Bureau, 2015  
AI/AN stands for American Indian and Alaska Native.  
NH/PI stands for Native Hawaiian and Other Pacific Islander.


Exhibit 1. Oklahoma Population by Age Group, 2014



Source: U.S. Census Bureau, 2015

#### The Number of Older Oklahomans Is Growing

Exhibit 2. Oklahoma Population by Age Group, 2000–2030



The proportion of Oklahoma's population that is 65 and older is growing while the proportion that is younger than 65 is shrinking. The U.S. Census Bureau estimates that 19.4 percent of Oklahoma's population will be 65 and older by the year 2030, an increase of 36.8 percent from 2015.

#### Projected Population in Oklahoma

Age Group	2015	2025	2030
Under 18	25.0%	24.9%	25.0%
18 to 44	34.4%	33.7%	33.4%
45 to 64	25.5%	23.0%	22.2%
65+	15.1%	18.4%	19.4%

Source: U.S. Census Bureau, 2005


## SUICIDE AMONG OLDER OKLAHOMANS

### Oklahoma Suicide Rate Compared With Regional and National Rates

The suicide rate among Oklahomans ages 50 and older is higher than the rate among younger age groups. In 2013, the total suicide rate among all people ages 50+ was 21.4 per 100,000 people (8.5 for women and 36.0 for men). The rate among those ages 50–64 was higher than both the rate in the region (including Arkansas, Louisiana, New Mexico, and Texas) and the rate in the United States.

States vary in their reporting of suicides. The suicide rate is influenced by these reporting practices.


Exhibit 3. Suicide Rates in Oklahoma, Region 6, and the United States, 2013



Source: Centers for Disease Control and Prevention (CDC), National Center for Health Statistics (NCHS), Underlying Cause of Death 1999-2013 on CDC WONDER Online Database, released 2015

### Trends in Suicide Rates in Oklahoma

Exhibit 4. Trends in Suicide Rates in Oklahoma by Age Group, 2004–2013



Source: CDC, NCHS, Underlying Cause of Death 1999-2013 on CDC WONDER Online Database, released 2015

The suicide rate among Oklahomans ages 50+ fluctuated from a low of 17.1 per 100,000 in 2005 to a high of 22.0 per 100,000 in 2011. From 2004 to 2013, the rate was generally highest among those in the 50–64 age group.


How a state reports suicides can vary from year to year. The number of suicides is generally low, so even a small difference in reported numbers may make the rate fluctuate widely.

## SUBSTANCE USE DISORDER AND SUBSTANCE USE DISORDER TREATMENT AMONG OLDER OKLAHOMANS

### 30-Day Binge Drinking Among Older Oklahomans

Binge drinking, defined as five or more drinks for men and four or more drinks for women on a single occasion, can lead to serious health problems. Such problems include neurological damage, cardiovascular disease, liver disease, stroke, and poor control of diabetes. Binge drinkers are more likely to take risks such as driving while intoxicated and to experience falls and other accidents. Older people have a lower tolerance for alcohol. Binge drinking decreases with age and occurs more frequently among men than it does among women. As Exhibit 5 shows, 11.5 percent of Oklahoma men ages 50–64 reported binge drinking in the past 30 days, while 3.2 percent of those in the 65+ group reported similar behavior.


Exhibit 5. Binge Drinking Rates in Oklahoma by Age Group and Sex, 2013



Source: Behavioral Risk Factor Surveillance System (BRFSS), 2013

### Illicit Drug Use Among Older Americans

Exhibit 6. Illicit Drug Use Among Older Americans, 2002–2013



Source: National Survey on Drug Use and Health (NSDUH), 2013  
 Illicit drug use includes illegal drugs and prescription drugs used nonmedically. SAMHSA provides a list of drugs included in its survey in the NSDUH methodological summary.

Nationally, the rate of illicit drug use among older adults ages 50–64 more than doubled from 2002 to 2013. This development partially reflects the entrance into this age group of the baby boom cohort, whose rates of illicit drug use have been higher than those of older cohorts. Although state-specific data are not available, the *Oklahoma Behavioral Health Barometer* is available for download from the Substance Abuse and Mental Health Services Administration (SAMHSA) website ([www.samhsa.gov/data/population-data-nsduh/reports?tab=33](http://www.samhsa.gov/data/population-data-nsduh/reports?tab=33)).


## Admissions to Substance Use Disorder Treatment Among Older Oklahomans

In 2012, there were 1,377 admissions of Oklahomans ages 50 and older to substance use disorder (SUD) treatment in state-funded treatment programs, a rate of 105.9 per 100,000 people ages 50+. This rate was higher than the regional rate and lower than the national average. Men made up 65.5 percent of these admissions. Of all admissions, 70.6 percent were White/Caucasian, 16.8 percent were Black/African American, and 2.3 percent were Hispanic.

The principal sources of referral to treatment among those ages 50 and older were:

Self-Referral	Criminal Justice	Other
48.1%	38.3%	13.6%


**Exhibit 7. SUD Treatment Admissions of Adults Ages 50+ in Oklahoma, Region 6, and the United States by Sex, 2012**



Source: Treatment Episode Data Set (TEDS), 2012<sup>1</sup>  
Data include only those clients reported to SAMHSA.

## SUD Treatment Admissions Among Oklahomans Ages 50+ by Insurance Type

**Exhibit 8. SUD Treatment Admissions of Adults Ages 50+ in Oklahoma, Region 6, and the United States by Insurance Type, 2012**



Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

In Oklahoma, there were no reported admissions for older adults who were uninsured or who had private insurance, 73.4 percent had Medicaid, and 26.6 percent had Medicare.

### *SUD Treatment Admissions Among Oklahomans Ages 50+ by Primary Sources of Payment*

Self-Pay	Medicaid	Other
Data not available	Data not available	Data not available

Exhibit 8 and the accompanying text reflect the type of health insurance (if any) clients had at admission; the table above represents primary sources of payment.


<sup>1</sup> TEDS data are collected by states as a condition of the Substance Abuse Prevention and Treatment Block Grant. Guidelines suggest that states report all clients admitted to publicly financed treatment; however, states are inconsistent in applying the guidelines. States may structure and implement different quality controls over the data. For example, states may collect different categories of information to answer TEDS questions. Information is then "walked over" to TEDS definitions.



## Alcohol Use Disorder Treatment Admissions Among Oklahomans Ages 50+

Alcohol was the most frequently cited substance used by older Oklahomans in publicly financed SUD treatment in 2012. It was mentioned as a substance of use in 57.2 percent of admissions among those ages 50+. This was higher than the regional rate and lower than the national rate.

**Exhibit 9. Alcohol Use Treatment Admissions of Adults Ages 50+ in Oklahoma, Region 6, and the United States, 2012**




Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

## SUD Treatment Admissions for Non-Alcohol Substance Use

**Exhibit 10. Non-Alcohol Substance Use Treatment Admissions of Adults Ages 50+ in Oklahoma, Region 6, and the United States, 2012**

Substances other than alcohol were cited as the primary substances of use for 42.8 percent of older adult admissions to publicly funded treatment in Oklahoma.



Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

## Drug-Related Emergency Department Visits Involving Pharmaceutical Misuse and Abuse by Older Adults

SAMHSA periodically releases reports from the Drug Abuse Warning Network (DAWN). DAWN, discontinued in 2011, consisted of a nationwide network of hospital emergency departments (EDs) primarily located in large metropolitan areas. DAWN data consist of professional reviews of ED records to determine the extent to which alcohol and other substance abuse was involved in ED visits. According to the November 25, 2010, *DAWN Report*:


- *In 2004, there were an estimated 115,803 ED visits involving pharmaceutical misuse and abuse by adults aged 50 or older; in 2008, there were 256,097 such visits, representing an increase of 121.1 percent*
- *One fifth (19.7 percent) of ED visits involving pharmaceutical misuse and abuse among older adults were made by persons aged 70 or older*
- *Among ED visits made by older adults, pain relievers were the type of pharmaceutical most commonly involved (43.5 percent), followed by drugs used to treat anxiety or insomnia (31.8 percent) and antidepressants (8.6 percent)*
- *Among patients aged 50 or older who visited the ED for pharmaceutical misuse or abuse, more than half (52.3 percent) were treated and released, and more than one third (37.5 percent) were admitted to the hospital*

Bulleted text in italics above is taken from SAMHSA's Center for Behavioral Health Statistics and Quality. (2010, November 25). *The DAWN Report: Drug-related emergency department visits involving pharmaceutical misuse and abuse by older adults*. Rockville, MD: SAMHSA.

## CO-OCCURRING SUBSTANCE USE AND MENTAL DISORDERS

### Older Oklahomans in SUD Treatment With Co-Occurring Mental Disorders

**Exhibit 11. SUD Treatment Admissions of Adults Ages 50+ With a Co-Occurring Mental Disorder in Oklahoma, Region 6, and the United States, 2012**



Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

The national literature shows a strong relationship between substance use and mental disorders. Studies show that 30–80 percent of individuals with a substance use or mental disorder also have a co-occurring disorder.


Exhibit 11 shows the proportion of SUD treatment admissions of Oklahomans ages 50+ with a co-occurring mental disorder. This rate is higher than the regional rate and the national average. However, state reporting practices are a factor in these results.

## MENTAL HEALTH

### Older Oklahomans Reporting Frequent Mental Distress

BRFSS, a household survey conducted in all 50 states and several territories, asks the following question: "Now thinking about your mental health, which includes stress, depression, and problems with emotions, for how many days during the past 30 days was your mental health not good?" Those individuals reporting 14 or more "Yes" days in response to this question experience frequent mental distress (FMD). Exhibit 12 shows that older Oklahomans experience FMD at a rate that is higher than the regional and national rates.


**Exhibit 12. Individuals Reporting Frequent Mental Distress in Oklahoma, Region 6, and the United States, 2013**



Source: BRFSS, 2013

### Older Oklahomans Reporting Frequent Mental Distress by Age Group and Sex

**Exhibit 13. Oklahomans Reporting Frequent Mental Distress by Age Group and Sex, 2013**



Source: BRFSS, 2013

Older men in Oklahoma were more likely to indulge in binge drinking, but older women were more likely to report that they had FMD (14 days or more per 30-day period). As Exhibit 13 shows, 18.2 percent of women in the 50–64 age group and 10.1 percent in the 65+ age group reported FMD, while 14.4 percent of men in the 50–64 age group and 7.0 percent in the 65+ age group reported FMD.

## Other Measures of Mental Health

BRFSS collected other measures showing risk factors for mental and/or physical illness. These included:

- Social and Emotional Support (2010). BRFSS asked, "How often do you get the social and emotional support you need?" The possible responses were always, usually, sometimes, rarely, or never.
- Life Satisfaction (2010). BRFSS asked, "In general, how satisfied are you with your life?" The possible responses were very satisfied, satisfied, dissatisfied, or very dissatisfied.

Exhibit 14 presents the results of these surveys among older Oklahomans.

**Exhibit 14. BRFSS Measures, 2010**


Indicator	Ages 50+	Ages 50–64	Ages 65+
Rarely or never get social or emotional support	9.0%	8.1%	10.4%
Dissatisfied or very dissatisfied	6.0%	7.7%	3.7%

Source: BRFSS, 2010

## Individuals With Frequent Mental Distress Report High Rates of Poor Physical Health

Older Americans who experienced FMD were more likely to report that their physical health was poor (14 days or more in the past 30-day period when physical health was "not good"). As shown in Exhibit 15, although nearly 11 percent of older Americans with no mental distress reported poor physical health, more than 50 percent of those with FMD reported poor physical health.

**Exhibit 15. Individuals Ages 50+ in the United States Reporting Poor Physical Health by Level of Mental Distress, 2013**




Source: BRFSS, 2013

## Relationship Among Mental Distress, Diabetes, Stroke, Heart Attack, High Blood Pressure, and Coronary Disease

Older Americans who experience FMD are more likely to report that they have health problems. People with FMD were more than twice as likely to report having a stroke than those with some or no mental distress. They experienced coronary disease and heart attack at more than 1.6 times, diabetes at more than 1.4 times, and high blood pressure at nearly 1.2 times the rate of those with some or no mental distress.

**Exhibit 16. Individuals Ages 50+ in the United States With Mental Distress and Serious Health Problems, 2013**



Source: BRFSS, 2013

## Older Oklahomans Admitted to State Mental Health Services

Approximately 1.9 percent of the people served by the Oklahoma mental health system were ages 65 and older. This represents more than 1,490 adults.

Source: Center for Mental Health Services (CMHS) Uniform Reporting System (URS) Output Tables, 2014

## DATA SOURCES

**BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM** ([www.cdc.gov/brfss](http://www.cdc.gov/brfss)). BRFSS, managed by CDC, is the largest ongoing telephone health survey system in the world, tracking health conditions and risk behaviors in the United States annually since 1984. Data are collected on a monthly basis in all 50 states, the District of Columbia, and participating territories. BRFSS data are collected by local jurisdictions and reported to CDC.

**CENTERS FOR DISEASE CONTROL AND PREVENTION, NATIONAL CENTER FOR HEALTH STATISTICS, UNDERLYING CAUSE OF DEATH 1999-2013 ON CDC WONDER ONLINE DATABASE, RELEASED 2015** (<http://wonder.cdc.gov/ucd-icd10.html>). The WONDER online database “contains mortality and population counts for all U.S. counties. Data are based on death certificates for U.S. residents. Each death certificate identifies a single underlying cause of death and demographic data. The number of deaths, crude death rates or age-adjusted death rates, and 95% confidence intervals and standard errors for death rates can be obtained by place of residence (total U.S., region, state and county), age group (single-year-of age, 5-year age groups, 10-year age groups and infant age groups), race, Hispanic ethnicity, gender, year, cause-of-death (4-digit ICD-10 code or group of codes), injury intent and injury mechanism, drug/alcohol induced causes and urbanization categories. Data are also available for place of death, month and week day of death, and whether an autopsy was performed.”

**CENTER FOR MENTAL HEALTH SERVICES UNIFORM REPORTING SYSTEM** ([www.samhsa.gov/data/us\\_map](http://www.samhsa.gov/data/us_map)). States that receive CMHS Block Grants are required to report aggregate data to URS. URS reports include information about utilization of mental health services and client demographic and outcome information.

**NATIONAL SURVEY ON DRUG USE AND HEALTH** (<https://nsduhweb.rti.org>). The NSDUH, managed by SAMHSA, is “an annual nationwide survey involving interviews with approximately 70,000 randomly selected individuals aged 12 and older.” NSDUH data are most frequently used by state planners to assess the need for SUD treatment. NSDUH data also include information about mental health conditions.

**TREATMENT EPISODE DATA SET** ([www.samhsa.gov/data/client-level-data-teds/reports?tab=19](http://www.samhsa.gov/data/client-level-data-teds/reports?tab=19)). States that receive Substance Abuse Prevention and Treatment Block Grant funds submit individual client data to TEDS. TEDS includes both admission and discharge data sets, and some 1.5 million admissions are reported annually. TEDS includes information about utilization of SUD treatment services and client demographic and outcome information. TEDS is an admission-based system, and TEDS admissions do not represent individuals. Thus, an individual admitted to treatment twice within a calendar year would be counted as two admissions.

**U.S. CENSUS BUREAU** ([www.census.gov/people](http://www.census.gov/people)). Two main sources of Census Bureau data were used in this report: (1) population estimates and (2) population projections.

# OLDER ADULTS BEHAVIORAL HEALTH PROFILE

---

Texas

### TEXAS' POPULATION

#### Texas Population by Age Group

Texas is home to 26,956,958 people. Of these:

- 7,826,161 (29.0 percent) are over age 50.
- 4,442,101 (16.5 percent) are over age 60.
- 2,023,234 (7.5 percent) are over age 70.
- 735,408 (2.7 percent) are ages 80 and older.


The proportion of women rises fairly steadily in each age group, and women make up 61.9 percent of the 80+ group. The racial/ethnic composition of older Texans is as follows:

#### Race/Ethnicity of Texans Ages 50+

White	AI/AN	Black	Asian	NH/PI	Other	Hispanic
83.7%	0.8%	10.8%	3.8%	0.1%	0.9%	25.6%

Source: U.S. Census Bureau, 2015  
AI/AN stands for American Indian and Alaska Native.  
NH/PI stands for Native Hawaiian and Other Pacific Islander.


Exhibit 1. Texas Population by Age Group, 2014



Source: U.S. Census Bureau, 2015

#### The Number of Older Texans Is Growing

Exhibit 2. Texas Population by Age Group, 2000–2030



The proportion of Texas' population that is 65 and older is growing while the proportion that is younger than 65 is shrinking. The U.S. Census Bureau estimates that 15.6 percent of Texas' population will be 65 and older by the year 2030, an increase of 66.6 percent from 2015.

#### Projected Population in Texas

Age Group	2015	2025	2030
Under 18	27.7%	27.4%	27.0%
18 to 44	37.0%	36.4%	36.3%
45 to 64	23.5%	21.6%	21.1%
65+	11.7%	14.6%	15.6%

Source: U.S. Census Bureau, 2005




## SUICIDE AMONG OLDER TEXANS

### Texas Suicide Rate Compared With Regional and National Rates

The suicide rate among Texans ages 50 and older is higher than the rate among younger age groups. In 2013, the total suicide rate among all people ages 50+ was 17.0 per 100,000 people (6.5 for women and 28.9 for men). The rate among those ages 50–64 was lower than both the rate in the region (including Arkansas, Louisiana, New Mexico, and Oklahoma) and the rate in the United States.

States vary in their reporting of suicides. The suicide rate is influenced by these reporting practices.


Exhibit 3. Suicide Rates in Texas, Region 6, and the United States, 2013



Source: Centers for Disease Control and Prevention (CDC), National Center for Health Statistics (NCHS), Underlying Cause of Death 1999-2013 on CDC WONDER Online Database, released 2015

### Trends in Suicide Rates in Texas

Exhibit 4. Trends in Suicide Rates in Texas by Age Group, 2004-2013



Source: CDC, NCHS, Underlying Cause of Death 1999-2013 on CDC WONDER Online Database, released 2015

The suicide rate among Texans ages 50+ fluctuated from a low of 14.2 per 100,000 in 2007 to a high of 17.2 per 100,000 in 2012. From 2004 to 2013, the rate was generally highest among those in the 50–64 age group.


How a state reports suicides can vary from year to year. The number of suicides is generally low, so even a small difference in reported numbers may make the rate fluctuate widely.

## SUBSTANCE USE DISORDER AND SUBSTANCE USE DISORDER TREATMENT AMONG OLDER TEXANS

### 30-Day Binge Drinking Among Older Texans

Binge drinking, defined as five or more drinks for men and four or more drinks for women on a single occasion, can lead to serious health problems. Such problems include neurological damage, cardiovascular disease, liver disease, stroke, and poor control of diabetes. Binge drinkers are more likely to take risks such as driving while intoxicated and to experience falls and other accidents. Older people have a lower tolerance for alcohol. Binge drinking decreases with age and occurs more frequently among men than it does among women. As Exhibit 5 shows, 17.7 percent of Texas men ages 50–64 reported binge drinking in the past 30 days, while 8.0 percent of those in the 65+ group reported similar behavior.


Exhibit 5. Binge Drinking Rates in Texas by Age Group and Sex, 2013



Source: Behavioral Risk Factor Surveillance System (BRFSS), 2013

### Illicit Drug Use Among Older Americans

Exhibit 6. Illicit Drug Use Among Older Americans, 2002–2013



Source: National Survey on Drug Use and Health (NSDUH), 2013  
 Illicit drug use includes illegal drugs and prescription drugs used nonmedically. SAMHSA provides a list of drugs included in its survey in the NSDUH methodological summary.

Nationally, the rate of illicit drug use among older adults ages 50–64 more than doubled from 2002 to 2013. This development partially reflects the entrance into this age group of the baby boom cohort, whose rates of illicit drug use have been higher than those of older cohorts. Although state-specific data are not available, the *Texas Behavioral Health Barometer* is available for download from the Substance Abuse and Mental Health Services Administration (SAMHSA) website ([www.samhsa.gov/data/population-data-nsduh/reports?tab=33](http://www.samhsa.gov/data/population-data-nsduh/reports?tab=33)).


## Admissions to Substance Use Disorder Treatment Among Older Texans

In 2012, there were 4,538 admissions of Texans ages 50 and older to substance use disorder (SUD) treatment in state-funded treatment programs, a rate of 58.0 per 100,000 people ages 50+. This rate was lower than the regional rate and the national average. Men made up 71.1 percent of these admissions. Of all admissions, 68.4 percent were White/Caucasian, 28.8 percent were Black/African American, and 20.0 percent were Hispanic.

The principal sources of referral to treatment among those ages 50 and older were:

Self-Referral	Criminal Justice	Other
45.9%	21.6%	32.6%


**Exhibit 7. SUD Treatment Admissions of Adults Ages 50+ in Texas, Region 6, and the United States by Sex, 2012**



Source: Treatment Episode Data Set (TEDS), 2012<sup>1</sup>  
Data include only those clients reported to SAMHSA.

## SUD Treatment Admissions Among Texans Ages 50+ by Insurance Type

**Exhibit 8. SUD Treatment Admissions of Adults Ages 50+ in Texas, Region 6, and the United States by Insurance Type, 2012**



Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

In Texas, 88.9 percent of older adult admissions to SUD treatment were uninsured, 8.3 percent had Medicaid, 2.1 percent had Medicare, and 0.6 percent had private insurance.

### *SUD Treatment Admissions Among Texans Ages 50+ by Primary Sources of Payment*

Self-Pay	Medicaid	Other
0.0%	0.0%	100.0%


Exhibit 8 and the accompanying text reflect the type of health insurance (if any) clients had at admission; the table above represents primary sources of payment.

<sup>1</sup> TEDS data are collected by states as a condition of the Substance Abuse Prevention and Treatment Block Grant. Guidelines suggest that states report all clients admitted to publicly financed treatment; however, states are inconsistent in applying the guidelines. States may structure and implement different quality controls over the data. For example, states may collect different categories of information to answer TEDS questions. Information is then "walked over" to TEDS definitions.

## Alcohol Use Disorder Treatment Admissions Among Texans Ages 50+

Alcohol was the most frequently cited substance used by older Texans in publicly financed SUD treatment in 2012. It was mentioned as a substance of use in 51.6 percent of admissions among those ages 50+. This was lower than the regional and national rates.

**Exhibit 9. Alcohol Use Treatment Admissions of Adults Ages 50+ in Texas, Region 6, and the United States, 2012**




Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

## SUD Treatment Admissions for Non-Alcohol Substance Use

**Exhibit 10. Non-Alcohol Substance Use Treatment Admissions of Adults Ages 50+ in Texas, Region 6, and the United States, 2012**

Substances other than alcohol were cited as the primary substances of use for 48.4 percent of older adult admissions to publicly funded treatment in Texas.



Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

## Drug-Related Emergency Department Visits Involving Pharmaceutical Misuse and Abuse by Older Adults

SAMHSA periodically releases reports from the Drug Abuse Warning Network (DAWN). DAWN, discontinued in 2011, consisted of a nationwide network of hospital emergency departments (EDs) primarily located in large metropolitan areas. DAWN data consist of professional reviews of ED records to determine the extent to which alcohol and other substance abuse was involved in ED visits. According to the November 25, 2010, *DAWN Report*:


- *In 2004, there were an estimated 115,803 ED visits involving pharmaceutical misuse and abuse by adults aged 50 or older; in 2008, there were 256,097 such visits, representing an increase of 121.1 percent*
- *One fifth (19.7 percent) of ED visits involving pharmaceutical misuse and abuse among older adults were made by persons aged 70 or older*
- *Among ED visits made by older adults, pain relievers were the type of pharmaceutical most commonly involved (43.5 percent), followed by drugs used to treat anxiety or insomnia (31.8 percent) and antidepressants (8.6 percent)*
- *Among patients aged 50 or older who visited the ED for pharmaceutical misuse or abuse, more than half (52.3 percent) were treated and released, and more than one third (37.5 percent) were admitted to the hospital*

Bulleted text in italics above is taken from SAMHSA's Center for Behavioral Health Statistics and Quality. (2010, November 25). *The DAWN Report: Drug-related emergency department visits involving pharmaceutical misuse and abuse by older adults*. Rockville, MD: SAMHSA.

## CO-OCCURRING SUBSTANCE USE AND MENTAL DISORDERS

### Older Texans in SUD Treatment With Co-Occurring Mental Disorders

**Exhibit 11. SUD Treatment Admissions of Adults Ages 50+ With a Co-Occurring Mental Disorder in Region 6 and the United States, 2012**



Source: TEDS, 2012  
Data include only those clients reported to SAMHSA.

The national literature shows a strong relationship between substance use and mental disorders. Studies show that 30–80 percent of individuals with a substance use or mental disorder also have a co-occurring disorder.


States may choose not to report some of the fields to TEDS, including information on SUD treatment admissions with a co-occurring disorder. These data were not reported by Texas in 2012. Therefore, Exhibit 11 shows regional and national figures only.

## MENTAL HEALTH

### Older Texans Reporting Frequent Mental Distress

BRFSS, a household survey conducted in all 50 states and several territories, asks the following question: "Now thinking about your mental health, which includes stress, depression, and problems with emotions, for how many days during the past 30 days was your mental health not good?" Those individuals reporting 14 or more "Yes" days in response to this question experience frequent mental distress (FMD). Exhibit 12 shows that older Texans experience FMD at a rate that is lower than the regional and national rates.


**Exhibit 12. Individuals Reporting Frequent Mental Distress in Texas, Region 6, and the United States, 2013**



Source: BRFSS, 2013

### Older Texans Reporting Frequent Mental Distress by Age Group and Sex

**Exhibit 13. Texans Reporting Frequent Mental Distress by Age Group and Sex, 2013**



Source: BRFSS, 2013

Older men in Texas were more likely to indulge in binge drinking, but older women were more likely to report that they had FMD (14 days or more per 30-day period). As Exhibit 13 shows, 12.9 percent of women in the 50–64 age group and 6.8 percent in the 65+ age group reported FMD, while 8.4 percent of men in the 50–64 age group and 5.2 percent in the 65+ age group reported FMD.

## Other Measures of Mental Health

BRFSS collected other measures showing risk factors for mental and/or physical illness. These included:

- Social and Emotional Support (2010). BRFSS asked, "How often do you get the social and emotional support you need?" The possible responses were always, usually, sometimes, rarely, or never.
- Life Satisfaction (2010). BRFSS asked, "In general, how satisfied are you with your life?" The possible responses were very satisfied, satisfied, dissatisfied, or very dissatisfied.

Exhibit 14 presents the results of these surveys among older Texans.

**Exhibit 14. BRFSS Measures, 2010**


Indicator	Ages 50+	Ages 50–64	Ages 65+
Rarely or never get social or emotional support	12.1%	11.4%	13.3%
Dissatisfied or very dissatisfied	5.2%	6.2%	3.6%

Source: BRFSS, 2010

## Individuals With Frequent Mental Distress Report High Rates of Poor Physical Health

Older Americans who experienced FMD were more likely to report that their physical health was poor (14 days or more in the past 30-day period when physical health was "not good"). As shown in Exhibit 15, although nearly 11 percent of older Americans with no mental distress reported poor physical health, more than 50 percent of those with FMD reported poor physical health.

**Exhibit 15. Individuals Ages 50+ in the United States Reporting Poor Physical Health by Level of Mental Distress, 2013**




Source: BRFSS, 2013

## Relationship Among Mental Distress, Diabetes, Stroke, Heart Attack, High Blood Pressure, and Coronary Disease

Older Americans who experience FMD are more likely to report that they have health problems. People with FMD were more than twice as likely to report having a stroke than those with some or no mental distress. They experienced coronary disease and heart attack at more than 1.6 times, diabetes at more than 1.4 times, and high blood pressure at nearly 1.2 times the rate of those with some or no mental distress.

**Exhibit 16. Individuals Ages 50+ in the United States With Mental Distress and Serious Health Problems, 2013**



Source: BRFSS, 2013

## Older Texans Admitted to State Mental Health Services

Approximately 3.1 percent of the people served by the Texas mental health system were ages 65 and older. This represents more than 9,980 adults.

Source: Center for Mental Health Services (CMHS) Uniform Reporting System (URS) Output Tables, 2014



## DATA SOURCES

**BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM** ([www.cdc.gov/brfss](http://www.cdc.gov/brfss)). BRFSS, managed by CDC, is the largest ongoing telephone health survey system in the world, tracking health conditions and risk behaviors in the United States annually since 1984. Data are collected on a monthly basis in all 50 states, the District of Columbia, and participating territories. BRFSS data are collected by local jurisdictions and reported to CDC.

**CENTERS FOR DISEASE CONTROL AND PREVENTION, NATIONAL CENTER FOR HEALTH STATISTICS, UNDERLYING CAUSE OF DEATH 1999-2013 ON CDC WONDER ONLINE DATABASE, RELEASED 2015** (<http://wonder.cdc.gov/ucd-icd10.html>). The WONDER online database “contains mortality and population counts for all U.S. counties. Data are based on death certificates for U.S. residents. Each death certificate identifies a single underlying cause of death and demographic data. The number of deaths, crude death rates or age-adjusted death rates, and 95% confidence intervals and standard errors for death rates can be obtained by place of residence (total U.S., region, state and county), age group (single-year-of age, 5-year age groups, 10-year age groups and infant age groups), race, Hispanic ethnicity, gender, year, cause-of-death (4-digit ICD-10 code or group of codes), injury intent and injury mechanism, drug/alcohol induced causes and urbanization categories. Data are also available for place of death, month and week day of death, and whether an autopsy was performed.”

**CENTER FOR MENTAL HEALTH SERVICES UNIFORM REPORTING SYSTEM** ([www.samhsa.gov/data/us\\_map](http://www.samhsa.gov/data/us_map)). States that receive CMHS Block Grants are required to report aggregate data to URS. URS reports include information about utilization of mental health services and client demographic and outcome information.

**NATIONAL SURVEY ON DRUG USE AND HEALTH** (<https://nsduhweb.rti.org>). The NSDUH, managed by SAMHSA, is “an annual nationwide survey involving interviews with approximately 70,000 randomly selected individuals aged 12 and older.” NSDUH data are most frequently used by state planners to assess the need for SUD treatment. NSDUH data also include information about mental health conditions.

**TREATMENT EPISODE DATA SET** ([www.samhsa.gov/data/client-level-data-teds/reports?tab=19](http://www.samhsa.gov/data/client-level-data-teds/reports?tab=19)). States that receive Substance Abuse Prevention and Treatment Block Grant funds submit individual client data to TEDS. TEDS includes both admission and discharge data sets, and some 1.5 million admissions are reported annually. TEDS includes information about utilization of SUD treatment services and client demographic and outcome information. TEDS is an admission-based system, and TEDS admissions do not represent individuals. Thus, an individual admitted to treatment twice within a calendar year would be counted as two admissions.

**U.S. CENSUS BUREAU** ([www.census.gov/people](http://www.census.gov/people)). Two main sources of Census Bureau data were used in this report: (1) population estimates and (2) population projections.