

Mapping a Pandemic: SARS-CoV-2 Seropositivity in the United States

Heather Kalish^{*1}, Carleen Klumpp-Thomas^{*2}, Sally Hunsberger^{*3}, Holly Ann Baus⁴, Michael P Fay³, Nalyn Siripong⁵, Jing Wang⁶, Jennifer Hicks¹, Jennifer Mehalko⁷, Jameson Travers², Matthew Drew⁷, Kyle Pauly¹, Jacquelyn Spathies¹, Tran Ngo⁸, Kenneth M. Adusei⁸, Maria Karkanitsa⁸, Jennifer A Croker⁹, Yan Li¹⁰, Barry I. Graubard¹¹, Lindsay Czajkowski⁴, Olivia Belliveau¹², Cheryl Chairez¹², Kelly Snead⁷, Peter Frank⁷, Anandakumar Shunmugavel⁸, Alison Han⁴, Luca T. Giurgea⁴, Luz Angela Rosas⁴, Rachel Bean⁴, Rani Athota⁴, Adriana Cervantes-Medina⁴, Monica Gouzoulis⁴, Brittany Heffelfinger⁴, Shannon Valenti⁵, Rocco Caldararo¹³, Michelle M. Kolberg¹⁴, Andrew Kelly², Reid Simon², Saifullah Shafiq², Vanessa Wall⁷, Susan Reed⁴, Eric W Ford⁹, Ravi Lokwani⁸, John-Paul Denson⁷, Simon Messing⁷, Sam G. Michael², William Gillette⁷, Robert P. Kimberly⁹, Steven E. Reis⁵, Matthew D. Hall², Dominic Esposito⁷, Matthew J. Memoli^{4†}, Kaitlyn Sadtler^{8†}

¹Trans-NIH Shared Resource on Biomedical Engineering and Physical Science, National Institute of Biomedical Imaging and Bioengineering, National Institutes of Health, Bethesda, MD 20894

²National Center for Advancing Translational Sciences, National Institutes of Health, Rockville, MD 20850

³Biostatistics Research Branch, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD 20894

⁴Clinical Studies Unit, Laboratory of Infectious Diseases, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD 20894

⁵Clinical and Translational Science Institute, University of Pittsburgh, Pittsburgh, Pennsylvania

⁶Clinical Monitoring Research Program Directorate, Frederick National Laboratory for Cancer Research, Frederick MD 21702

⁷Protein Expression Laboratory, NCI RAS Initiative, Frederick National Laboratory for Cancer Research, Frederick MD 21702

⁸Section on Immuno-Engineering, National Institute of Biomedical Imaging and Bioengineering, National Institutes of Health, Bethesda, MD 20894

⁹Center for Clinical and Translational Science, School of Medicine, University of Alabama at Birmingham, Birmingham, AL 35294

¹⁰Joint Program in Survey Methodology, Department of Epidemiology and Biostatistics, University of Maryland College Park, College Park, MD 20742

¹¹Division of Cancer Epidemiology & Genetics, Biostatistics Branch, National Cancer Institute, National Institutes of Health, Bethesda MD 20894

¹²Laboratory of Immunoregulation, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD 20894

¹³Clinical Research Directorate, Frederick National Laboratory for Cancer Research, Leidos Biomedical Research, Inc, Frederick MD 21702

¹⁴Division of Clinical Research, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD 20894

*these authors contributed equally

†Address correspondence to: Dr. Matthew Memoli (memolim@nih.gov) or Dr. Kaitlyn Sadtler (kaitlyn.sadtler@nih.gov)

45 **ABSTRACT**

46

47 Asymptomatic SARS-CoV-2 infection and delayed implementation of diagnostics have led to poorly
48 defined viral prevalence rates. To address this, we analyzed seropositivity in US adults who have not
49 previously been diagnosed with COVID-19. Individuals with characteristics that reflect the US population
50 ($n = 11,382$) and who had not previously been diagnosed with COVID-19 were selected by quota sampling
51 from 241,424 volunteers (ClinicalTrials.gov NCT04334954). Enrolled participants provided medical,
52 geographic, demographic, and socioeconomic information and 9,028 blood samples. The majority (88.7%)
53 of samples were collected between May 10th and July 31st, 2020. Samples were analyzed via ELISA for
54 anti-Spike and anti-RBD antibodies. Estimation of seroprevalence was performed by using a weighted
55 analysis to reflect the US population. We detected an undiagnosed seropositivity rate of 4.6% (95% CI: 2.6
56 – 6.5%). There was distinct regional variability, with heightened seropositivity in locations of early
57 outbreaks. Subgroup analysis demonstrated that the highest estimated undiagnosed seropositivity within
58 groups was detected in younger participants (ages 18-45, 5.9%), females (5.5%), Black/African American
59 (14.2%), Hispanic (6.1%), and Urban residents (5.3%), and lower undiagnosed seropositivity in those with
60 chronic diseases. During the first wave of infection over the spring/summer of 2020 an estimate of 4.6% of
61 adults had a prior undiagnosed SARS-CoV-2 infection. These data indicate that there were 4.8 (95% CI:
62 2.8-6.8) undiagnosed cases for every diagnosed case of COVID-19 during this same time period in the
63 United States, and an estimated 16.8 million undiagnosed cases by mid-July 2020.

64 **INTRODUCTION**

65

66 COVID-19, the disease caused by SARS-CoV-2 infection, presents with a spectrum of illness ranging from
67 asymptomatic to severe disease and death. As with most respiratory viral diseases, it is difficult to estimate
68 the true prevalence of the disease during a pandemic and the extent of its spread is only known after
69 extensive study¹⁻³. The majority of patients infected develop robust antibody responses against the viral
70 spike (S), nucleocapsid (N), and envelope (E) proteins that can be detected via serologic testing⁴⁻⁸. Anti-S
71 antibodies persist for months, and can neutralize infection⁹. Frequently, these neutralizing antibodies bind
72 the receptor binding domain (RBD) of the spike protein, but antibodies against the spike S2 domain have
73 also been observed¹⁰⁻¹⁵.

74

75 To characterize the spread of SARS-CoV-2 infection in the United States, we evaluated seropositivity in a
76 national survey of participants who had not previously been diagnosed with SARS-CoV-2 infection. We
77 used quota sampling from a large pool of volunteers to obtain a representative sample and performed
78 statistical weighting to generate prevalence estimates which provide a clear picture of the extent of SARS-
79 CoV-2 infection. To ensure accurate classification of seropositivity, we utilized our dual-antigen ELISA
80 protocol that evaluated IgG and IgM antibodies against both the full spike ectodomain and the RBD^{7,16}.
81 These foundational considerations generated critical data needed to estimate spread during the pandemic
82 and gain insight into the potential future outcomes.

83

84 These results, including the subgroup analysis, give us a previously undescribed view into the spread of the
85 pandemic by more clearly identifying the large numbers of individuals with undiagnosed infections during
86 the initial months of the pandemic. These data are of great importance as we consider the impact vaccination
87 may have on the future course of the pandemic and plan for current and future available vaccines to be
88 administered. In addition, these data can also help us better assess the public health measures taken during
89 the pandemic and how to take the best approaches forward to any future public health emergencies.

90 **METHODS**

91

92 *Study Protocol:*

93

94 This study was designed to determine the seroprevalence of anti-SARS-CoV-2 antibodies in adults 18 years
95 of age or older in the United States who had not been previously diagnosed with COVID-19. The primary
96 endpoint was the weighted estimate of seroprevalence in the US. Secondary endpoints were weighted
97 estimates for subgroups categorized by demographics/risk factors. An initial period enrolled a convenience
98 sample of 593 volunteers prior to the quota sample. Participants across the US (all 50 states and DC) were
99 then enrolled via telephone consent from a pool of volunteers who provided basic demographic data in
100 response to the study announcement. Recruitment calls were made from three sites: NIAID Laboratory of
101 Infectious Diseases Clinical Studies Unit, the University of Pittsburgh CTSI, and the University of Alabama
102 at Birmingham CCTS. Selection of participants is described below. Selected participants were contacted
103 by the study team, consented, and sent a blood microsampling kit and online questionnaire in REDCap
104 (project-redcap.org). For a small subset of participants ($n = 214$) working on the NIH campus, serum was
105 collected via venipuncture. This study (ClinicalTrials.gov NCT04334954) was approved by the National
106 Institutes of Health Institutional Review Board and conducted in accordance with the provisions of the
107 Declaration of Helsinki and Good Clinical Practice guidelines. All participants provided verbal informed
108 consent prior to enrollment.

109

110 *Participant Selection*

111

112 All volunteers were emailed an initial survey to collect basic demographic characteristics. Survey responses
113 were de-identified and aggregated by sub-category of state, type of locality approximated from zip codes,
114 age, sex, race, and ethnicity (**Figure 1**). Target sample sizes for these sub-categories were determined from
115 the U.S. census, and were updated every evening based on the characteristics of people who had already
116 enrolled to assure that individuals in each sub-category were enrolled evenly over time. Within each sub-
117 category, participants were initially assigned a selection probability calculated from the target number as a
118 proportion of the available pool. Specific sub-categories that had insufficient numbers were aggregated to
119 estimate their impact on the overall distribution of the 6 main characteristics. If a particular characteristic
120 had insufficient numbers, sample probabilities were boosted for volunteers who had the characteristic. For
121 each day's call list, the most representative of 20,000 randomly generated lists was used, each list drawn
122 without replacement from the volunteer pool based on the sampling probabilities previously defined.
123 Representativeness was assessed by estimating a weighted sum of squared differences from the desired

124 targets and picking the list with the lowest deviation. Unselected participants were eligible to be called at a
125 later date. This algorithm is designed such that each cohort of invited participants is representative of the
126 diversity of the US population with respect to the 6 sampling variables (see Statistical Supplement Section
127 3.4).

128

129 *Sample Collection:*

130

131 Participants provided blood samples by mail using a Mitra microsampling kit (Neoteryx, Torrance, CA) or
132 standard venipuncture. Microsampling kits contained visual instructions on the sampling process, bandages,
133 gauze, lancets, and four 20 μ l microsampling devices for a total collection of 80 μ l of whole blood.
134 Participants utilized the lancet to draw blood from their fingertip and collect blood onto each of the four
135 microsamplers. Participants returned the dried microsamplers with desiccant via overnight shipping. Those
136 who underwent venipuncture did so in the NIH Clinical Center phlebotomy lab where 18 ml of blood was
137 collected in a serum separator and whole blood tube. Once received in the laboratory serum samples were
138 processed, and microsamplers were stored dry at -80°C until elution and analysis.

139

140 *Serologic Assays:*

141

142 Antibodies from samples were analyzed using ELISA as previously described^{7,16-18}. In order to maintain
143 longitudinal quality control and ensure that the assays remained stable across multiple months of assay
144 implementation, positive and negative controls were included on each assay plate and monitored for
145 stability (**Supplemental Fig. 1**). Seropositivity cut points were defined by evaluating 300 true negative
146 samples and 56 true positive samples. Positivity thresholds were based on the mean optical density
147 (absorbance) plus 3 standard deviations (see Supplemental Materials for details). The final criterion of a
148 Spike⁺ and RBD⁺ for any combination of IgG or IgM gave estimated sensitivity and specificity of 1, with
149 raw values for recombinant antibody results reported in **Supplemental Fig. 2** and **Supplemental Table 1**.
150 Additionally, IgA was evaluated via previously described ELISA to further phenotype the participant's
151 serologic status.

152

153 *Statistical Analysis*

154

155 The previously described iterative quota sampling continuously matched the proportion of people in the
156 study with the census estimated proportion of people in the country on 6 variables (**Table 1, Figure 1**).
157 This ensured that each periodic sample of participants over the course of the study were representative, and

158 the time effects of the pandemic were approximately independent of those 6 variables. Each participant was
159 asked demographic and health-related questions that matched ones on the Behavioral Risk Factor
160 Surveillance System (BRFSS) survey, a large probability-based national survey¹⁹. Responses to those
161 matching questions were used with BRFSS survey data to adjust estimators to account for important criteria
162 that may be related to both selection probability and seropositivity but were not accounted for in the quota
163 sampling. Those adjusted estimators used weighting based on the propensity of being a quota sample versus
164 a BRFSS sample participant and poststratification to US census data. It additionally accounted for
165 sensitivity and specificity. Confidence intervals were calculated for the final seroprevalence estimates
166 accounting for both the variability of the weighting and of the sensitivity and specificity adjustment. The
167 ratio of undiagnosed cases over diagnosed cases was estimated as the final seroprevalence estimate times a
168 factor calculated from the daily national population and diagnosed cases. For more methods and details see
169 Section 3 of the **Supplementary Materials**.

170

171 **RESULTS**

172

173 *Enrollment and Demographic Representation*

174

175 Recruitment took place from April 1, 2020 until August 4, 2020. During that time 11,283 participants were
176 enrolled from a pool of 241,424 volunteers. Of these participants, 214 had blood collected via venipuncture
177 and 11,069 were sent microsamplers. Over 80% of the microsamplers were returned (9,089 participants).
178 Ultimately 9,028 participant samples were analyzed via ELISA for presence of SARS-CoV-2 antibodies.
179 Of those, 8,058 participants had complete clinical questionnaire data and were included in the weighted
180 analysis (**Figure 1**). The majority (>88%) of sample collection occurred within the 11-week period between
181 May 10th to July 31st, 2020 (**Supplemental Fig. 3**). The six major demographic factors used in participant
182 selection are summarized in **Table 1**. Participant sampling was highly representative of the U.S. population.
183 When expanded to include the additional 10 demographic or health related factors captured by the BRFSS,
184 many factors were well matched, but there were some differences: our sample population was more highly
185 educated, employed, and had better access to healthcare (**Table 1**).

186

187 *Estimates of Seroprevalence*

188

189 There were 304 seropositive participants in the analysis set (**Figure 2a,b**). This gave a weighted estimate
190 of 4.6% of the undiagnosed adults in the U.S. population that were seropositive for SARS-CoV-2 (95% CI:
191 2.6% to 6.5%, $n = 8058$ complete testing and survey). Using this average rate over the study period, we

192 estimate that there were 4.8 undiagnosed cases per each diagnosed case over the course of the study (95%
193 CI: 2.8, 6.8). In seropositive participants, 36.51% were IgG⁺IgM⁺IgA⁺, 28.29 % were IgG⁺IgM⁻IgA⁺,
194 17.11% were IgG⁺IgM⁻IgA⁻, 13.16 % were IgG⁻IgM⁺IgA⁻, 4.28 % were IgG⁻IgM⁻IgA⁻, and 0.66 % were
195 IgG⁻IgM⁺IgA⁺ (**Figure 2a-c, Supplemental Fig. 4**).

196
197 We found regional variations of seroprevalence estimates across the US (**Fig. 2d, 3**). The Northeast and
198 Mid-Atlantic Regions showed the highest rates of seropositivity whereas the lowest in the Midwest. Urban
199 areas were estimated to have higher levels of seropositivity (5.3%) compared to rural areas (1.1%
200 seropositivity) at the time samples were collected. Estimates of seroprevalence were calculated for other
201 demographic subgroups (**Figure 3**). The youngest age group, 18-44, had the highest estimated
202 seropositivity (5.9%). Estimated seroprevalence for females was 5.5% and 3.5% in males. The
203 seroprevalence estimate for Black/African Americans was highest at 14.2% followed by participants who
204 self-identified as other/unlisted race (11.1%), American Indian/Alaska Native (6.8%), followed by
205 White/Caucasian (2.5%), while those identifying as Asian displayed the lowest seroprevalence estimate
206 (2.0%).

207
208 Participants who reported a known exposure to a SARS-CoV-2-infected individual had a higher
209 seroprevalence estimate (15.6%) compared to those who did not (2.7%). In comparison to the national
210 average (4.6%), those that worked from home had a lower seropositivity estimate of 3.0%. Those who
211 reported prior vaccination (influenza 3.2% and/or pneumonia 2.3%) had a lower likelihood for undiagnosed
212 seropositivity. Those who had health conditions associated with poor outcomes in SARS-CoV-2 infection,
213 including coronary heart disease, asthma, and diabetes, displayed lower rates of seropositivity (**Figure 4**).
214 Other health conditions were also correlated with a decreased seropositivity rate such as skin cancer, stroke,
215 or arthritis.

216 217 **DISCUSSION**

218
219 This study demonstrates that spread of the SARS-CoV-2 virus in the US during the first six months of the
220 pandemic was more widespread than has been suggested by data reporting diagnostic test-confirmed cases.
221 Similar to responses to other respiratory viruses, such as influenza, many individuals develop asymptomatic
222 or mild disease that is not medically attended and therefore never diagnosed. Our findings indicate that
223 there are nearly five individuals with a previous asymptomatic infection for every diagnosed case.
224 Furthermore, patterns of our seroprevalence data match well with those of diagnosed cases reported during
225 a similar timeframe.²⁰ For example, the greater seropositivity estimated in densely populated urban areas

226 follows the observed initial spread of SARS-CoV-2. In comparison to the national average, we found that
227 the Midwest, South, and West had lower seroprevalences during the study timeframe, which preceded a
228 substantial increase in infections in these regions detected by viral testing.

229

230 Our data suggest that the youngest age group had the highest undiagnosed seroprevalence, which is
231 consistent with observations that they display less severe symptoms than older patients²¹. We also found
232 higher undiagnosed seroprevalence in females, possibly suggesting a higher risk for asymptomatic disease.
233 Participants with chronic diseases that are more likely to be associated with severe clinical manifestations
234 of COVID-19, including diabetes, heart disease, and asthma, had a lower prevalence of asymptomatic
235 SARS-CoV-2 infection in comparison to the national average. Those with known exposure to SARS-CoV-
236 2 infected individuals had a higher estimated incidence of undiagnosed seropositivity. We also found that
237 Black, African American, and Hispanic participants had higher undiagnosed seropositivity, correlating with
238 national data on disease burden in these sub-groups.

239

240 This study is the first to report a representative sample across the US and to evaluate regional, demographic
241 and socioeconomic differences in the prevalence of asymptomatic SARS-CoV-2 infection. In contrast,
242 other reports of seroprevalence data focus on a specific group of individuals or geographic location²². Our
243 results provide new insight into the spread of SARS-CoV-2. Our estimate of the national undiagnosed
244 exposure rate provides information on the scope of infection during the first six months of the pandemic.
245 This work extends findings from smaller foundational studies of limited populations²³⁻³⁷ by generating an
246 accurate estimate of nationwide and subgroup prevalence.

247

248 Our results estimate that there are approximately 4.8 undiagnosed cases (95% CI 2.76-6.81) for every
249 identified case of COVID-19, suggesting a potential 16.8 million undiagnosed cases by mid-July 2020 in
250 addition to the reported 3 million diagnosed cases in the United States. These data suggest a higher level of
251 infection-induced immunity exists in the population and the size of those with this immunity is even greater
252 now as the virus continued to spread in the months since this study was performed. Further long-term
253 studies of immunity in the population will be necessary to further understand durability of response to the
254 vaccine versus infection, how infection-induced immunity impacts vaccine response and performance, and
255 if herd immunity can play a role in controlling SARS-CoV-2 spread. In addition, further subgroup analysis
256 of our data will be useful in clarifying the spread of disease in the presence of public health measures and
257 how we may be able to refine and further target those measures in the future.

258

259

260 *Limitations*

261

262 Although we were able to recruit a cohort with demographics representative of the general US population,
263 our study has several limitations. First, although extensive statistical adjustments were made, our study
264 cohort is based on a non-random volunteer sample which can have selection bias. However, many
265 traditional random sampling studies using probability sampling design have very low response rates, calling
266 into question the advantages of that practice^{38,39}. Our study population also exhibited some differences from
267 the general US population, such as higher education level and access to healthcare that had to be adjusted
268 for with statistical weighting. We utilized both census and behavioral data to weight our results though it is
269 possible that there are variables associated with disease transmission that are not accounted for in our
270 weighting.

271

272 **CONCLUSIONS**

273

274 These data suggest a much larger spread of the COVID-19 pandemic than originally thought and have
275 implications in basic understanding of SARS-CoV-2 spread, epidemiologic characteristics of its spread and
276 prevalence in different communities, and potential impact on decisions involved in vaccine rollout.
277 Continued large-scale surveillance of SARS-CoV-2 immunity is in progress, discriminating infection-based
278 and vaccine-induced antibody responses, and mathematical models will be generated to understand the
279 pandemic, vaccine performance, public health measure efficacy, and providing insight for our approach to
280 handling the next virus with pandemic potential.

281 **ACKNOWLEDGEMENTS**


282

283 Firstly, we would like to thank the participants in this study, without whom it would not have been possible.
284 The authors would like to acknowledge Dr. Corbett and Dr. Graham of the NIAID VRC for their generous
285 donation of coronavirus spike expression plasmid, and Dr. Aaron Schmidt, J Feldman, BM Hauser and T
286 M Caradonna of the Ragon Institute of MGH, MIT, and Harvard for their donation of their RBD expression
287 plasmid. We thank John Crissey and the CRIMSON team (NIAID, NIH) for their assistance and
288 contributions to the study and data management. We thank Dr. Jason McLellan for scholarly discussions
289 regarding the spike S-2P construct. We thank the CCTS Clinical Research Support Program (A. Delbridge,
290 L. Dukes), UAB School of Public Health and its Survey Research Unit (L. Battle, J. Carson, M. DeRamus,
291 T. Fields, T. Graham, T. Jackson, E. Pruitt, A. Underwood, P. Wolff), the University of Pittsburgh Clinical
292 and Translational Science Institute staff and leadership (J. Avolio, L. Bash, S. Clayton, M. Cristinziano, K.
293 Daw, C. Fascetti, E. Gyurisin, J. Huwe, N. Jones, D. Mathias, S. Mathias, A. Mykita, B. Petersen, M.
294 Phillips, C. Rush, E. Shepherd, S. Shetty, A. Socci, L. Stearns, S. Ugbomah, K. Underwood, L. Yasko) and
295 University of Pittsburgh Information Technology (D. McGaughey, S. Ritzman, T. Smith) for their
296 contributions to this work. We would like to thank Mr. Denzel Bernard for his assistance with sample
297 delivery, and Dr. Stephanie Ford-Scheimer for organizational assistance. This research was supported in
298 part by the Intramural Research Program of the NIH, including the National Institute for Biomedical
299 Imaging and Bioengineering, the National Institute of Allergy and Infectious Disease, and the National
300 Center for Advancing Translational Sciences. This project has been funded in part with Federal funds from
301 the National Cancer Institute, National Institutes of Health, under contract number HHSN261200800001E,
302 75N91019D00024, Task Order No. 75N91019F00130, Clinical and Translational Science Awards Program
303 grants UL1TR003096 (UAB) and UL1TR001857 (University of Pittsburgh). The content of this publication
304 does not necessarily reflect the views or policies of the Department of Health and Human Services, nor
305 does mention of trade names, commercial products, or organizations imply endorsement by the U.S.
306 Government. Disclaimer: The NIH, its officers, and employees do not recommend or endorse any company,
307 product, or service.

	US Population (BRFSS)			CoV2 Serosurvey Population	
	n	%	weighted (%)	n	%
Selection Criteria					
Region					
North East	91307	21.19	17.6	1508	16.7
Midwest	67110	15.57	16.97	1445	16.01
Mid-Atlantic	80979	18.79	16.91	1833	20.3
South/Central	60482	14.03	15.35	1293	14.32
Mountain/Southwest	86204	20	15.89	1392	15.42
West/Pacific	44866	10.41	17.27	1557	17.25
Age Group					
18 - 45	125081	28.59	46	3837	42.51
45 - 70	207749	47.49	39.84	3783	41.91
70 - 95	104605	23.91	14.17	1407	15.59
Sex					
Male	197411	45.24	48.66	4318	47.83
Female	238911	54.76	51.34	4710	52.17
Urban/Rural					
Urban	365714	84.9	93.48	8550	94.78
Rural	65234	15.1	6.52	471	5.22
Race					
White only	345710	81	73.41	6986	77.4
Black only	37862	8.87	12.9	830	9.2
Others	43219	10.13	13.69	1210	13.41
Ethnicity					
Hispanic	36941	8.53	17.06	1495	16.56
Not Hispanic	395931	91.47	82.94	7532	83.44
Additional Weighting Criteria					
Children					
Yes	113408	26.21	35.81	2943	32.88
No	319281	73.79	64.19	6009	67.12
Education					
<=HS	151606	34.79	41.07	240	2.68
College	119979	27.53	30.88	1284	14.35
>=College	164229	37.68	28.05	7422	82.96
Homeowner					
Own	305545	70.36	66.49	6635	74.12

Rent	107208	24.69	27.32	1861	20.79
Others	21535	4.96	6.19	456	5.09
Employment					
Employed	219493	50.75	57.74	6364	71.09
NLF	174920	40.45	31.38	2129	23.78
Unemployed	38053	8.8	10.88	459	5.13
Health Insurance					
Yes	400028	91.86	87.85	8697	97.31
No	35433	8.14	12.15	240	2.69
Flu Vaccinated					
Yes	234727	59	50.62	6198	73.73
No	163124	41	49.38	2208	26.27
Cardiovascular Disease					
Yes	52284	12.07	9.07	354	3.98
No	380985	87.93	90.93	8541	96.02
Pulmonary Disease					
Yes	84102	19.33	18.53	1671	18.96
No	350913	80.67	81.47	7140	81.04
Immune Disease					
Yes	170115	39.14	29.29	2039	23.1
No	264571	60.86	70.71	6787	76.9
Diabetes					
Yes	60703	13.9	11.41	482	5.41
No	375876	86.09	88.59	8430	94.59


309 **Table 1: Characteristics of serosurvey population in comparison to United States population.** Census
310 and Behavioral Risk Factor Surveillance System (BRFSS, 2018) data on selection criteria were utilized for
311 quota-based sampling. Other values from BRFSS were utilized for statistical weighting. The comparisons
312 between the estimated proportions in the United States (BRFSS) versus our sample population for the
313 SARS-CoV-2 serosurvey are displayed in this table.



315

316


317 **Figure 1: Serosurvey overview and statistical workflow.** A flow chart of donor recruitment through data
 318 analysis displaying steps in data acquisition and any attrition from data sets if applicable. Key: Ovals =
 319 starts and ends, gray rectangles = subsets of participants in this study, blue parallelograms = individuals
 320 from outside data sets that contribute to adjusted prevalence estimates, blue rounded rectangles = analysis
 321 processes.




322

323 **Figure 2: Geographic distribution of undiagnosed seropositivity in the United States in summer 2020.**

324 Raw serology data for (a) IgG and (b) IgM and (c) IgA against SARS-CoV-2 Spike and Receptor Binding
 325 Domain (RBD). Cut points for positivity are shown as red dashed lines, data are optical density (OD). (d)
 326 Serologic phenotype of antibody presence in seropositive participants (e) US Map showing seropositivity
 327 in six regions surveyed: Northeast = ME, NH, VT, MA, NY, CT, RI, PA, NJ, 7.5% (95% CI: 3.7 – 11.3%);
 328 Midwest = MN, IA, WI, IL, IN, MI, OH, 1.6% (95% CI: 0.06-2.3%); Mid-Atlantic = MD, DE, DC, VA,
 329 WV, KY, TN, NC, SC, GA, 8.6% (1.3 – 15.8%); South/Central = FL, MS, AL, LA, AR, MO, KS, OK,
 330 3.0% (1.2 – 4.5%); Mountain/Southwest = TX, NM, AZ, CO, UT, WY, NE, SD, ND, MT, ID, 4.5% (0.09
 331 – 7.9%); West/Pacific = WA, OR, NV, CA, AK, HI, 1.9% (0.02 – 3.2%). One person in diagram represents
 332 100 participants, orange represents weighted prevalence estimate within the geographic region.



334 **Figure 3: Undiagnosed seroprevalence in main demographic categories.** Six main categories utilized
 335 during quota-based sampling: region, age, sex, race, ethnicity, and urban/rural. Seropositivity estimates of
 336 samples that had a full clinical questionnaire completed and successful sampling. Data are weighted
 337 estimates \pm 95% confidence intervals. Dashed line = weighted national seroprevalence estimate. * = n value
 338 too low to make proper weighted estimate, raw positivity displayed.



340 **Figure 4: Seroprevalence estimates of health and behavioral traits.** Seropositivity estimates of samples
 341 that had a full clinical questionnaire completed and successful sampling. Data are weighted estimates ±
 342 95% confidence intervals. Dashed line = weighted national seroprevalence estimate.

REFERENCES

- 343
344
345 1. Fauci AS, Lane HC, Redfield RR. Covid-19—navigating the uncharted. In: Mass Medical Soc;
346 2020.
- 347 2. Taubenberger JK, Morens DM. The pathology of influenza virus infections. *Annu Rev Pathol Mech*
348 *Dis.* 2008;3:499-522.
- 349 3. Taubenberger JK, Morens DM. 1918 Influenza: the mother of all pandemics. *Revista Biomedica.*
350 2006;17(1):69-79.
- 351 4. Huang AT, Garcia-Carreras B, Hitchings MD, et al. A systematic review of antibody mediated
352 immunity to coronaviruses: antibody kinetics, correlates of protection, and association of antibody
353 responses with severity of disease. *medRxiv.* 2020.
- 354 5. Long Q-X, Liu B-Z, Deng H-J, et al. Antibody responses to SARS-CoV-2 in patients with COVID-
355 19. *Nature medicine.* 2020:1-4.
- 356 6. Ainsworth M, Andersson M, Auckland K, et al. Performance characteristics of five immunoassays
357 for SARS-CoV-2: a head-to-head benchmark comparison. *The Lancet Infectious Diseases.* 2020.
- 358 7. Klumpp-Thomas C, Kalish H, Drew M, et al. Standardization of enzyme-linked immunosorbent
359 assays for serosurveys of the SARS-CoV-2 pandemic using clinical and at-home blood sampling.
360 *medRxiv.*
- 361 8. Stadlbauer D, Amanat F, Chromikova V, et al. SARS-CoV-2 Seroconversion in Humans: A
362 Detailed Protocol for a Serological Assay, Antigen Production, and Test Setup. *Current Protocols*
363 *in Microbiology.* 2020;57(1):e100.
- 364 9. Dan JM, Mateus J, Kato Y, et al. Immunological memory to SARS-CoV-2 assessed for greater
365 than six months after infection. *bioRxiv.* 2020.
- 366 10. Chi X, Yan R, Zhang J, et al. A neutralizing human antibody binds to the N-terminal domain of the
367 Spike protein of SARS-CoV-2. *Science.* 2020;369(6504):650-655.
- 368 11. He Y, Zhou Y, Liu S, et al. Receptor-binding domain of SARS-CoV spike protein induces highly
369 potent neutralizing antibodies: implication for developing subunit vaccine. *Biochemical and*
370 *biophysical research communications.* 2004;324(2):773-781.
- 371 12. Liu L, Wang P, Nair MS, et al. Potent neutralizing antibodies against multiple epitopes on SARS-
372 CoV-2 spike. *Nature.* 2020;584(7821):450-456.
- 373 13. Pinto D, Park Y-J, Beltramello M, et al. Cross-neutralization of SARS-CoV-2 by a human
374 monoclonal SARS-CoV antibody. *Nature.* 2020:1-6.
- 375 14. Shi R, Shan C, Duan X, et al. A human neutralizing antibody targets the receptor binding site of
376 SARS-CoV-2. *Nature.* 2020:1-8.
- 377 15. Wang Z, Lorenzi JC, Muecksch F, et al. Enhanced SARS-CoV-2 Neutralization by Secretory IgA
378 in vitro. *bioRxiv.* 2020.
- 379 16. Hicks J, Klumpp-Thomas C, Kalish H, et al. Serologic cross-reactivity of SARS-CoV-2 with
380 endemic and seasonal Betacoronaviruses. *medRxiv.* 2020.
- 381 17. Esposito D, Mehalko J, Drew M, et al. Optimizing high-yield production of SARS-CoV-2 soluble
382 spike trimers for serology assays. *Protein Expression and Purification.* 2020:105686.
- 383 18. Mehalko J, Drew M, Snead K, et al. Improved production of SARS-CoV-2 spike receptor-binding
384 domain (RBD) for serology assays. *Protein expression and purification.* 2020:105802.
- 385 19. Centers for Disease Control and Prevention (CDC) AG. Behavioral Risk Factor Surveillance
386 System Survey Questionnaire and Data. In: U.S. Department of Health and Human Services
387 CfDcAp, ed2020.
- 388 20. Dong E, Du H, Gardner L. An interactive web-based dashboard to track COVID-19 in real time.
389 *The Lancet infectious diseases.* 2020;20(5):533-534.
- 390 21. Mehra MR, Desai SS, Kuy S, Henry TD, Patel AN. Cardiovascular Disease, Drug Therapy, and
391 Mortality in Covid-19. *New England Journal of Medicine.* 2020;382(25):e102.
- 392 22. Rostami A, Sepidarkish M, Leeftang MMG, et al. SARS-CoV-2 seroprevalence worldwide: a
393 systematic review and meta-analysis. *Clin Microbiol Infect.* 2020.

- 394 23. Caini S, Bellerba F, Corso F, et al. Meta-analysis of diagnostic performance of serological tests for
395 SARS-CoV-2 antibodies up to 25 April 2020 and public health implications. *Euro Surveill.*
396 2020;25(23).
- 397 24. Kontou PI, Braliou GG, Dimou NL, Nikolopoulos G, Bagos PG. Antibody Tests in Detecting
398 SARS-CoV-2 Infection: A Meta-Analysis. *Diagnostics (Basel)*. 2020;10(5).
- 399 25. Lee CY, Lin RTP, Renia L, Ng LFP. Serological Approaches for COVID-19: Epidemiologic
400 Perspective on Surveillance and Control. *Front Immunol*. 2020;11:879.
- 401 26. Lisboa Bastos M, Tavaziva G, Abidi SK, et al. Diagnostic accuracy of serological tests for covid-
402 19: systematic review and meta-analysis. *BMJ*. 2020;370:m2516.
- 403 27. Biggs HM, Harris JB, Breakwell L, et al. Estimated community seroprevalence of SARS-CoV-2
404 antibodies—two Georgia counties, April 28–May 3, 2020. *Morbidity and Mortality Weekly Report*.
405 2020;69(29):965.
- 406 28. Bryan A, Pepper G, Wener MH, et al. Performance characteristics of the Abbott Architect SARS-
407 CoV-2 IgG assay and seroprevalence in Boise, Idaho. *Journal of clinical microbiology*. 2020.
- 408 29. Dietrich ML, Norton EB, Elliott D, et al. SARS-CoV-2 Seroprevalence Rates of Children in
409 Louisiana During the State Stay at Home Order. *medRxiv*. 2020.
- 410 30. Feehan AK, Fort D, Garcia-Diaz J, et al. Seroprevalence of SARS-CoV-2 and infection fatality
411 ratio, Orleans and Jefferson Parishes, Louisiana, USA, May 2020. *Emerging infectious diseases*.
412 2020;26(11):2765.
- 413 31. Havers FP, Reed C, Lim T, et al. Seroprevalence of antibodies to SARS-CoV-2 in 10 sites in the
414 United States, March 23-May 12, 2020. *JAMA*. 2020;180(12):1576-1586.
- 415 32. McLaughlin CC, Doll MK, Morrison KT, et al. High community SARS-CoV-2 antibody
416 seroprevalence in a ski resort community, Blaine County, Idaho, US. preliminary results. *Medrxiv*.
417 2020.
- 418 33. Menachemi N, Yiannoutsos CT, Dixon BE, et al. Population point prevalence of SARS-CoV-2
419 infection based on a statewide random sample—Indiana, April 25–29, 2020. *Morbidity and*
420 *Mortality Weekly Report*. 2020;69(29):960.
- 421 34. Ng DL, Goldgof GM, Shy BR, et al. SARS-CoV-2 seroprevalence and neutralizing activity in
422 donor and patient blood. *Nature communications*. 2020;11(1):1-7.
- 423 35. Rosenberg ES, Tesoriero JM, Rosenthal EM, et al. Cumulative incidence and diagnosis of SARS-
424 CoV-2 infection in New York. *medRxiv*. 2020.
- 425 36. Sood N, Simon P, Ebner P, et al. Seroprevalence of SARS-CoV-2–Specific Antibodies Among
426 Adults in Los Angeles County, California, on April 10-11, 2020. *Jama*. 2020.
- 427 37. Sutton M, Cieslak P, Linder M. Notes from the Field: Seroprevalence Estimates of SARS-CoV-2
428 Infection in Convenience Sample—Oregon, May 11–June 15, 2020. *Morbidity and Mortality*
429 *Weekly Report*. 2020;69(32):1100.
- 430 38. Elliott MR, Valliant R. Inference for nonprobability samples. *Statistical Science*. 2017;32(2):249-
431 264.
- 432 39. Hill D. The dirty little secret pollsters need to own up to. *The Washington Post*. 2020.
433