

**Retention and the Intersection of Structural Inequities  
in a Breastfeeding Intervention Study**

Helen Wilde LaPlant<sup>1</sup>, Confidence Francis-Edoziuno<sup>1</sup>, Zhe Guan<sup>2</sup>, Tumilara Aderibigbe<sup>1</sup>,  
Xiaolin Chang<sup>2</sup>, Ashwag Saad Alhabodal<sup>1</sup>, Kristen Delaney<sup>1</sup>, Dana Scott<sup>3</sup>, Mary Marshall-Crim<sup>4</sup>,  
Idelisa Freytes<sup>5</sup>, Wendy A. Henderson<sup>1</sup>, Stephen Walsh<sup>1</sup>, and Ruth F. Lucas<sup>1</sup>

<sup>1</sup> School of Nursing, University of Connecticut

<sup>2</sup> Department of Statistics, University of Connecticut

<sup>3</sup> Obstetrics and Gynecology Unit, Breast Health and Cancer Genetics, University of Connecticut  
Health Center

<sup>4</sup> Women's Health Services, Lactation Program, Hartford Hospital

<sup>5</sup> Women's Ambulatory Health Services, Hartford Hospital

Author Note

**Helen Wilde LaPlant, MS, IBCLC**, is a Project Manager, School of Nursing, University of Connecticut, Storrs, Connecticut, and Program Manager, IBCLC, Hispanic Health Council, Hartford, Connecticut.

**Confidence Francis-Edoziuno, BSN**, is a PhD student and Research Assistant, School of Nursing, University of Connecticut, Storrs, Connecticut.

**Zhe Guan, MA**, is a PhD student and Research Assistant, Department of Statistics, University of Connecticut, Storrs, Connecticut.

**Tumilara Aderibigbe, PhD**, is a Post-doctoral Fellow at University of Utah, Salt Lake City, Utah. At the time this research was completed, she was a Graduate Assistant, School of Nursing, University of Connecticut, Storrs, Connecticut.

**Xiaolin Chang, PhD**, was a Graduate Assistant, Department of Statistics, University of Connecticut, Storrs, Connecticut at the time of data collection.

**Ashwag Saad Alhabodal, RN, MHA/HE**, is a PhD student and Research Assistant, School of Nursing, University of Connecticut, Storrs, Connecticut.

**Kristen Delaney, MS, APRN, FNP-BC**, was a Research Assistant, School of Nursing, University of Connecticut, Storrs, Connecticut at the time this research was completed.

**Dana Scott, MD, FACOG**, is an Assistant Professor, Obstetrics and Gynecology Unit, University of Connecticut Health Center, Farmington, Connecticut.

**Mary Marshall-Crim, MSN, RN, IBCLC**, is Lactation Manager, Women's Health Services, Lactation Program, Hartford Hospital, Hartford, Connecticut.

**Idelisa Freytes, BSHCA**, is Practice Manager, Women's Ambulatory Health Services, Hartford Hospital, Hartford, Connecticut.

**Wendy A. Henderson, PhD, CRNP, FAASLD, FAAN**, is a Gail and Ralph Reynolds Presidential Distinguished Professor, Department of Biobehavioral Health Sciences, University of Pennsylvania School of Nursing, and at the time of this research was Director of the PhD Program, School of Nursing, University of Connecticut, Storrs, Connecticut.

**Stephen Walsh, ScD**, is Associate Professor, School of Nursing, University of Connecticut, Storrs, Connecticut.

**Ruth Lucas, Ph.D., RNC, CLS, FAAN**, is an Associate Professor, School of Nursing, University of Connecticut, Storrs, Connecticut.

Corresponding author: Ruth Lucas, PhD, RNC, CLS, 231 Glenbrook Road, Unit 4026, University of Connecticut, Storrs, CT 06269-4026 (e-mail: [ruth.lucas@uconn.edu](mailto:ruth.lucas@uconn.edu)).

**Keywords:** Disruptive technology, breastfeeding, smartphone, pain self-management

### Abstract

**Introduction:** Women below the poverty threshold have lower representation and retention in breastfeeding studies.

**Methods:** A secondary analysis of a longitudinal randomized controlled self-management for breast and nipple pain during breastfeeding study. Participants completed online surveys at discharge, weeks 1, 2, 3, 6, 9, 12, 18, and 24, with face-to-face interviews at 6 and 24 weeks. Text messages were sent to participants when modules and surveys were due. Retention was assessed in R with descriptive statistics, Mann-Whitney, Pearson's chi-square, and Cox Proportional Hazard Regression.

**Results:** Two hundred and forty-four women ( $89 \leq \$50,000$  and  $155 > \$50,000$ ) were recruited. Retention rates at 1 (93%), 2 (87%), 6 (82%), 9 (77%) and 24 (72%) weeks. For women of low income compared to those of high income there was a hazard ratio (HR) of 2.5 ( $p=0.0001$ ) for retention. For non-Hispanic Black and Hispanic women compared to the combined non-Hispanic White and Other group, HRs for retention were 3.3 and 2.6 respectively ( $p=0.0001$ ). Adjustment for age in the final hazard regression model of income, age, race and ethnicity decreased the HR for women of low income to 1.6 and HRs for non-Hispanic Black and Hispanic women to 2.1 and 1.9, respectively ( $p=0.0001$ ). However, none of the individual factors in the model achieved statistical significance.

**Discussion:** Retention in breastfeeding studies impacts breastfeeding duration, a key lifelong preventative health behavior. Despite accessible study design, retention of women desiring to breastfeed was adversely affected by the intersection of income, race and ethnicity, and age.

**Keywords:** Disruptive technology, breastfeeding, smartphone, pain self-management

## **Retention and the Intersection of Structural Inequities in a Breastfeeding Intervention Study**

Breastfeeding (BF) is fundamental to lifelong protective health equity, yet women and infants below the poverty threshold disproportionately do not exclusively breastfeed for six months (24 weeks), and thus do not receive this benefit.<sup>1-4</sup> In 2020, 83.1% of women in the United States (US) initiated BF, and 58.2% continued until 6 months.<sup>4</sup> However, for women below the federal poverty threshold, only 75.4 % ever breastfed and 44.2% breastfed to 6 months.<sup>4</sup>

Despite improvement in overall BF outcomes,<sup>5</sup> significant disparities in BF rates exist due to social and structural inequities of income, race, and environment.<sup>2,6,7</sup> Low income has been shown to have a considerable impact on ability to engage in general protective health behaviors.<sup>8,9</sup> As 19.5% of Black women and 17.1% of Hispanic women are below the poverty threshold, they are less likely to have health insurance and access to BF resources,<sup>10-13</sup> putting them at risk of lower BF rates. Cultural, social, and economic factors, such as inequitable access to lactation resources and support, coupled with inadequate knowledge of the benefits of BF, negatively impact BF practice.<sup>14-21</sup> Therefore, strategies to increase access to BF services and reduce the effects of poverty and structural inequities are necessary for BF equity.<sup>6,18,22,23</sup>

A potential strategy for reducing BF disparities is to increase research study participation of underrepresented groups and individuals at risk of suboptimal health outcomes.<sup>24</sup> Research outcomes provide evidence that supports the development of disparity-reducing policies and health interventions.<sup>25</sup> For research and interventions to be beneficial and generalizable, a diverse participant population that reflects the intended user population should be recruited.<sup>26,27</sup> Effective recruitment and retention strategies contribute significantly to the overall success of clinical studies.<sup>28</sup> However, recruitment and retention of persons in low-income groups is challenging,

resulting in under representation in clinical research studies.<sup>26,28,29</sup> A text-based intervention is an accessible means of promoting BF equity.<sup>30</sup> Through the government Lifeline Assistance program, low-income mothers can access smartphones.<sup>31,32</sup> which enables utilization of text-based interventions to promote BF equity. Therefore, the purpose of this paper is to describe retention rates for women above and below the poverty threshold in a BF study.

## Methods

The *Promoting Self-Management of Breast and Nipple Pain with Biomarkers and Technology for Breastfeeding Women (PROMPT)* (R56NR020041, PI, R. Lucas) study took place in partnership with two academic hospitals in the Northeast. The randomized clinical trial tested the efficacy of a text-message and cloud-based patient-informed self-management intervention. The intervention delivers BF information and support through nurse led text-based communication to decrease breast and nipple pain and increase BF self-efficacy and adaptive coping behaviors (Figure 1). For equitable access, text links were delivered via smartphone, which 95% of women of childbearing age possess.<sup>33</sup> The design used a one-finger swipe method, so women may access the intervention during BF. Women were randomized to intervention or attention control groups and followed via surveys and face-to-face interviews over a 24-week period.

## Sampling

The study aimed to enroll 222 participants. The inclusion criteria were (a) 18 - 45 years of age; (b) birth < 48 hours to a singleton infant of > 37 weeks gestational age; (c) intention to breastfeed; (d) received BF basics during antenatal care; (e) access to the internet via own or study provided smartphone; (f) ability to read and write English; (g) lactation consultant or counselor BF assessment. Exclusion criteria were (a) < 18 years of age; (b) history of significant mental

health disorder (e.g., major depression or bipolar disorder); (c) skin conditions on non-dominant forearm (related to pain sensitivity testing),<sup>34,35</sup> and (d) delivery of an infant with medical complications or congenital anomalies.

### ***Participant Recruitment***

To achieve a diverse sample of participants, members of the research team were hired to represent the ethnic and racial diversity of the women who gave birth at each hospital. The recruitment flyer and educational modules were reviewed by the team for language level and inclusive images. Advertisement was via the PI's online lab webpage and flyers provided at prenatal clinics and labor and delivery. Screening for eligible participants occurred 4-6 times per week through review of the postpartum census by the project manager in consultation with the PI. The research team prioritized whom to approach first, based on time from delivery, race and ethnicity, and confirmed eligibility with the nurse providing care for each potential participant. Eligible women were approached; if the woman expressed interest, a study flyer was left for her review and a time for return was agreed upon. The research member commonly returned multiple times due to ongoing clinical and infant care, occasionally the next day before the woman was discharged.

### ***Participant Consent***

Eligible women were consented in their private hospital rooms. During enrollment, each participant was assessed for capacity to provide consent, HIPAA consent was obtained, the consent form was reviewed, and the woman was asked three questions to confirm understanding. Screening was documented via REDCap (Research Electronic Data Capture, v.13), a secure web-based database, and consent documents were stored in the clinical setting. Women were randomized

using an established dynamic minimization algorithm based on six stratifying characteristics (age, race, route of delivery, BF experience, expected BF duration, and intent to return to work) collected during screening.<sup>36,37</sup>

### ***Participant Retention***

After consent, research team members initiated encrypted text-linked surveys via Twilio (a text distribution platform embedded in REDCap) and confirmed participant receipt and completion. In addition, women completed surveys on BF, pain, and maternal wellbeing and underwent quantitative sensory testing. At enrollment, women without a smartphone were provided a study phone with unlimited data to access the 24-week study.

Following discharge, participants completed online surveys at 1, 2, 3, 6, 9, 12, 18, and 24 weeks and received modules on managing BF pain and challenges, postpartum care, infant development, and maternal mental health. Participants returned at 6 and 24 weeks for in-person interviews and quantitative sensory testing at one of three locations easily accessible by public transport. Scheduling was flexible to accommodate participant availability.

Surveys and modules were sent to participants at predetermined intervals via text links. If items were not completed within six hours, a second invitation link was automatically generated. The online platform allowed participants to begin survey completion and save their results to resume later. Completion of survey instruments and modules was monitored daily by the project manager. If items were not being completed within the one-week window, links were re-sent to participants for remaining surveys or modules in the series, with reminder text messages, telephone calls, and lastly, email messages if participants did not respond to previous methods.

Retention was encouraged through progressive monetary incentives, with a bonus upon completion of the study. To maintain engagement, participants received text links to normal infant development and age-appropriate play modules at weeks 4, 8, 12, 16, 20, and 24. At week 7, a text message was sent to encourage continued participation regardless of feeding method.

### ***Data Analysis and Management***

All data were entered directly into REDCap, a password-protected database, and downloaded for analysis using R software.<sup>38</sup> Descriptive statistics (percentages for categorical data and means/standard deviations for interval data) were determined. Plots of retention relative to time were created for various groups of participants using the Kaplan-Meier method and were formally compared between groups using weighted log-rank tests. To address the hypothesis that a combination of income and other common characteristics of BF outcomes would be associated with retention at 1, 2, 3, 6, 9, 12, 18, and 24 weeks, Cox proportional hazards regression modeling was conducted. Model demographics included income, self-reported race and ethnicity (non-Hispanic black, Hispanic, and a combination of non-Hispanic white and other races [Asian, Native Hawaiian or Pacific Islander, American Indian, and more than one race]), age, route of delivery, history of BF, pre-pregnancy body mass index, insurance, return to work, and infant gender.

## **Results**

### ***Sociodemographic Characteristics***

Due to successful recruitment, in just over 15 months our team increased our target sample to 250 participants representative of greater diversity than Connecticut state population.<sup>39</sup> Participants (N=250) were self-reported as 118 (47.2%) non-Hispanic white, 43 (17.2%) non-Hispanic Black, 71 (28.4%) Hispanic, and 18 (7.2%) Other. Those who self-identified as Hispanic



were 41 (57.7 %) White, 13 (18.3%) Black, 11 (15.5%) more than one race, 3 (4.2%) Native Hawaiian or Pacific Islander, 2 (2.8%) American Indian, and 1 (1.4%) Asian. Other included those who self-identified as the following: 12 (66.7%) non-Hispanic Asian, 5 (27.8%) non-Hispanic more than one race, and 1 (5.6%) non-Hispanic Native Hawaiian or Pacific Islander. Two hundred and forty-four participants reported their yearly income, 89 (36.5%)  $\leq$ \$50,000 and 155 (63.5%)  $>$ \$50,000. Other participant characteristics known to affect BF outcomes: age, route of delivery, history of BF, return to work, pre-pregnancy body mass index, education, and infant gender and weight, are reported in Table 1.

### ***Recruitment and Retention***

Retention rates in the study, for 244 women who provided income data, were at 1 (93%), 2 (87%), 3 (85%), and 6 (82%) weeks (Table 2). From weeks 9 through 24, retention rates declined from 77% to 72% (Table 2). Retention over time in the high-income group was significantly higher than in the low-income group ( $p < 0.0001$ ). There was a statistically significant difference in patterns of retention over time between the race and ethnicity categories ( $p < 0.0001$ ). Retention was higher and comparable for non-Hispanic White and Other women, but lower and comparable for non-Hispanic Black and Hispanic women.

To identify individual and combined effects of income and race and ethnicity on retention, Cox Proportional Hazard Regression<sup>40</sup> analyses were conducted. Due to similarity in retention patterns, the race and ethnicity groups were recoded to a combination of non-Hispanic White and Other women, and a second combination of non-Hispanic Black and Hispanic women. These analyses revealed a hazard ratio (HR) of 2.5 for women of low income compared to those of high income ( $p = 0.0001$ ). This HR implies that low-income status is associated with a 150% increase in

the hazard of withdrawal from the study at any point in time. When non-Hispanic Black and Hispanic women were compared to the combined non-Hispanic White and Other group, the HRs were 3.3 and 2.6 respectively ( $p=0.0001$ ).

In a multivariable Cox model, that included both income and race and ethnicity factors, race and ethnicity continued to be statistically significant ( $p=0.04$ ), but income was no longer significant ( $p=0.06$ ). The adjusted HR for low-income women decreased to 1.7. The adjusted HRs for non-Hispanic Black and Hispanic women decreased to 2.4 and 2.0, respectively.

Proportional hazards regression was also used to investigate potential associations between other covariates and retention. For age, the HR was 1.1, when comparing younger women to older women, suggesting that risk of withdrawal increases by 10% with each one year decrease in age ( $p=0.009$ ). For education, women with attainment of high school or below had a HR of 2.0 compared to those with a college degree or more ( $p=0.04$ ). Lack of prior BF experience was associated with a HR of 2.6 relative to women with prior BF experience ( $p=0.051$ ). Other maternal characteristics known to impact BF studies were not significant.

To investigate potential confounding effects, the significant covariates were successively entered into several models that included the retention, income, and race and ethnicity variables. In the model with the income and race and ethnicity factors, none of the covariates remained statistically significant. However, adjustment for age altered the magnitudes of HRs for the income and race and ethnicity effects in a way that suggested it could be a confounding factor. Specifically, the HR for women in the low income versus high income group decreased to 1.6 and HRs for non-Hispanic Black and Hispanic women decreased to 2.1 and 1.9, respectively.

In the final three-factor model of income, self-reported race and ethnicity, and age, the simultaneous statistical test on all three variables was significant ( $p=.0001$ ). However, none of the individual variables achieved statistical significance. Comparison of nested models with and without age established that, with adjustment for age, the combination of the income and race and ethnicity factors was statistically significant ( $p = 0.006$ ). This suggests that it is the overlap of information between income and race and ethnicity that results in the model being significantly predictive of withdrawal from the study. More technically, it implies that a collinearity between the three indicator variables for the income and race and ethnicity factors predicts retention as opposed to withdrawal (Table 3).

## Discussion

Retention in prospective longitudinal studies of a population below the poverty threshold is challenging in BF and other health-related studies.<sup>23,41</sup> The present study over-enrolled women from income and racial and ethnic populations who are historically under-represented in clinical studies<sup>28,42</sup> and are less likely to initiate and sustain BF.<sup>3</sup> Taken independently, women of low income and diverse race and ethnicity had significantly lower retention in the study. However, when income, race and ethnicity, and age are examined together, their intersection significantly decreases the risk of not being retained in the study. The intersection of these factors represents the structural inequities for women who desire to BF yet are unable to be retained in a BF study.

To facilitate equitable participation, our study utilized an accessible text-based BF intervention to support women to meet their BF goals. Similarly, studies with women below the poverty threshold and Hispanic and non-Hispanic Black women found that the use of a text message-based intervention was acceptable to participants, provided equitable access, and resulted in retention in BF studies.<sup>30,43,44</sup> Reflective of structural inequities, women in our study who had

incomes below the poverty threshold and who self-identified as non-Hispanic Black and Hispanic had a 2.1 and 1.9 times higher risk of not being retained in the study than non-Hispanic White and Other women. Despite inclusive text-based support, women were unable to continue in their chosen BF study.

Historically, women who participate in and complete breastfeeding studies report higher breastfeeding rates on average compared to national rates.<sup>20,44,45</sup> Furthermore, women below the poverty threshold and who self-identify as non-Hispanic Black and Hispanic are sub-optimally represented in clinical research.<sup>42</sup> Our study retention reflects these structural inequities, as non-Hispanic Black and Hispanic women had a higher risk of not being retained in the study, with their risk of dropping out not solely accounted for by income. Structural inequities contribute to disparities in BF rates, particularly for non-Hispanic Black and Hispanic women in the US compared with other racial and ethnic groups.<sup>20,46,47</sup> Cultural values, insufficient BF knowledge and support, and ongoing barriers related to social determinants of health contribute to low initiation of BF for non-Hispanic Black women and early formula introduction among Hispanic women, thus endangering long term BF and impeding retention in a breastfeeding study.<sup>14,17–20,46</sup>

Factors associated with positive retention in our study were increasing maternal age and previous BF history. In the 2018 CDC National Immunization Survey, age was a significant predictor of breastfeeding behavior, as women  $\leq 29$  years are less likely to ever breastfeed.<sup>4</sup> In addition, age and prior BF experience are covariates associated with participation in breastfeeding studies.<sup>4,48</sup> The intersection of age and previous BF history reflects complex underlying structural inequities which create barriers to Hispanic and non-Hispanic Black women for participation in BF studies and in meeting BF goals.<sup>12,18,49</sup> Therefore, while text-based BF support is accessible and acceptable, it requires adaptation to address the unique challenges, barriers, and structural

inequities encountered by Hispanic and non-Hispanic Black women in meeting their BF goals.<sup>23,41,50,51</sup>

Key to successful retention of participants in our longitudinal study were strategies proven effective in a diverse population.<sup>24,26,45,52</sup> Skillful implementation of these strategies was driven by our study's diverse and culturally sensitive research team. Recruitment and face-to-face interviews were conducted by team members hired from the participants' communities. In-person visits at 6 and 24 weeks were encouraged by flexible scheduling at a time mutually convenient to participant and research team, at the client's preferred location (hospital or clinic), with easy accessibility to public transportation. Daily assessment of survey completion and follow-up outreach (survey reminder links, and text, telephone, and email messages) were conducted by the project manager to facilitate data collection and participant retention. After observing a pattern of withdrawal after the early weeks of the study, a text message to encourage participant continuation regardless of current feeding method was suggested and implemented. Progressive compensation was also provided to participants, with a bonus for completion.

### **Limitations**

While this study used an accessible BF intervention, the exclusion criteria of non-English speakers and mothers with infants of  $\leq 37$  weeks' gestation may have omitted many women of racial and ethnic diversity who were otherwise eligible. Another limitation was conducting the study in a state with a comparatively high cost of living, reflected in the significant difference in Hispanic and non-Hispanic retention across the study, contributing to BF inequity for women with  $\leq \$50,000$ . Beyond this paper's scope is the effect on retention due to differences in BF pain perception across racial and ethnic groups, the focus of the larger study. To validate the efficacy of a text-based intervention to support BF equity, a larger study, specifically one that includes

Spanish-speaking women and mothers of infants of 37 weeks' gestation, is needed to address underlying structural inequities and provide equitable support for Hispanic and non-Hispanic Black women to reach their BF goals.

## Conclusions

Women of all people groups want to breastfeed. Findings from this study suggest that text-based support is accessible and acceptable for women from diverse income and racial and ethnic backgrounds. However, while women from all income and racial and ethnic groups were successfully enrolled in the current study, Hispanic and non-Hispanic Black women had lower rates of retention, which suggests that there are complex underlying structural inequities, not solely explained by income level. The use of predictor variables, such as age and previous breastfeeding history may inform new strategies for retaining women from lower income and underrepresented racial and ethnic groups in future research of interventions to promote and support breastfeeding. Research should also include collaboration with community partners to aid in identifying and addressing structural inequities to increase retention in future studies and sustain breastfeeding as a positive health outcome.

**Acknowledgements:** Research reported in this publication was supported by the National Institute of Nursing Research of the National Institutes of Health under Award Number R56NR020041 to Dr. Ruth Lucas. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. The authors thank the participants of the PROMPT study for their time and commitment to participation.

**Authorship Contribution Statement:** Author 1: Project administration, Writing - review & editing. Author 2: Writing - review & editing. Author 3: Formal analysis, Visualization. Author 4: Writing - original draft. Author 5: Formal analysis. Author 6: Writing - review & editing, Visualization. Author 7: Investigation. Author 8: Project administration. Author 9: Project administration. Author 10: Project administration. Author 11: Writing - review & editing. Author 12: Formal analysis, Writing – review & editing. Author 13: Conceptualization, Methodology, Supervision, Funding acquisition, Writing - review & editing.

***Conflicts of Interest:*** The authors have no conflicts of interest to report.

***Ethical Conduct of Research:*** The PROMPT study was performed in line with the principles of the Declaration of Helsinki. Approval was granted by the Ethics Committee of University of Connecticut, Hartford Hospital, and University of Connecticut Health Center (6-30-2021/E-HHC-2021-0143). Informed consent was obtained from all participants in the PROMPT study.

***Clinical Trial Registration:*** NCT05262920. Registered 2 March 2022. The first participant was enrolled on 16 March 2022. <https://clinicaltrials.gov/ct2/show/NCT05262920>

## References

1. Abegunde D, Hutchinson P, Anaba U, et al. Socioeconomic inequality in exclusive breastfeeding behavior and ideation factors for social behavioral change in three north-western Nigerian states: a cross-sectional study. *Int J Equity Health* 2021;20(1):172; doi: 10.1186/s12939-021-01504-4.
2. Temple Newhook J, Newhook LA, Midodzi WK, et al. Poverty and breastfeeding: comparing determinants of early breastfeeding cessation incidence in socioeconomically marginalized and privileged populations in the FiNaL study. *Health Equity* 2017;1(1):96–102; doi: 10.1089/heq.2016.0028.
3. CDC. 2022 Breastfeeding Report Card. 2023. Available from: <https://www.cdc.gov/breastfeeding/data/reportcard.htm> [Last accessed: 3/13/2024].
4. CDC. Results: Breastfeeding Rates. 2023. Available from: [https://www.cdc.gov/breastfeeding/data/nis\\_data/results.html](https://www.cdc.gov/breastfeeding/data/nis_data/results.html) [Last accessed: 3/13/2024].
5. Kirksey K. A social history of racial disparities in breastfeeding in the United States. *Soc Sci Med* 1982 2021;289:114365; doi: 10.1016/j.socscimed.2021.114365.
6. Crear-Perry J, Correa-de-Araujo R, Lewis Johnson T, et al. Social and structural determinants of health inequities in maternal health. *J Womens Health* 2002 2021;30(2):230–235; doi: 10.1089/jwh.2020.8882.
7. Pérez-Escamilla R. Breastfeeding in the 21st century: how we can make it work. *Soc Sci Med* 1982 2020;244:112331; doi: 10.1016/j.socscimed.2019.05.036.
8. Hughes MC, Baker TA, Kim H, et al. Health behaviors and related disparities of insured adults with a health care provider in the United States, 2015-2016. *Prev Med* 2019;120:42–49; doi: 10.1016/j.ypmed.2019.01.004.
9. Papageorge NW, Zahn MV, Belot M, et al. Socio-demographic factors associated with self-protecting behavior during the Covid-19 pandemic. *J Popul Econ* 2021;34(2):691–738; doi: 10.1007/s00148-020-00818-x.
10. U.S. Census Bureau. Poverty in the United States: 2022. 2023. Available from: <https://www.census.gov/library/publications/2023/demo/p60-280.html> [Last accessed: 3/13/2024].
11. Mahajan S, Caraballo C, Lu Y, et al. Trends in differences in health status and health care access and affordability by race and ethnicity in the United States, 1999-2018. *JAMA* 2021;326(7):637–648; doi: 10.1001/jama.2021.9907.



12. Schpero WL, Morden NE, Sequist TD, et al. For selected services, Blacks and Hispanics more likely to receive low-value care than Whites. *Health Aff Proj Hope* 2017;36(6):1065–1069; doi: 10.1377/hlthaff.2016.1416.
13. Willner C. Racial and Ethnicity Disparities in Earnings, Employment, and Education in Connecticut. 2022. Available from: <https://www.ctdata.org/acs-racial-and-ethnicity-disparities> [Last accessed: 3/13/2024].
14. Aderibigbe O, Lucas R. Exclusive breastfeeding in African American women: a concept analysis. *J Adv Nurs* 2023;79(5):1699–1713; doi: 10.1111/jan.15301.
15. Asiodu IV, Bugg K, Palmquist AEL. Achieving breastfeeding equity and justice in Black communities: past, present, and future. *Breastfeed Med Off J Acad Breastfeed Med* 2021;16(6):447–451; doi: 10.1089/bfm.2020.0314.
16. Hernández-Cordero S, Pérez-Escamilla R. What will it take to increase breastfeeding? *Matern Child Nutr* 2022;18(S3):e13371; doi: 10.1111/mcn.13371.
17. Kopp SJ, Kelly EA, DeFranco EA. Influence of social determinants of health on breastfeeding intent in the United States. *Birth* 2023;50(4):858–867; doi: 10.1111/birt.12740.
18. Louis-Jacques AF, Marhefka SL, Brumley J, et al. Historical antecedents of breastfeeding for African American women: from the pre-colonial period to the mid-twentieth century. *J Racial Ethn Health Disparities* 2020;7(5):1003–1012; doi: 10.1007/s40615-020-00727-5.
19. Rabb K, Alikhani A, von Ash T, et al. Examining Black women’s breastfeeding experiences in the United States: a scoping review of qualitative studies. *Matern Child Health J* 2023;27(10):1726–1741; doi: 10.1007/s10995-023-03692-7.
20. Rhodes EC, Damio G, LaPlant HW, et al. Promoting equity in breastfeeding through peer counseling: the US Breastfeeding Heritage and Pride program. *Int J Equity Health* 2021;20(1):128; doi: 10.1186/s12939-021-01408-3.
21. Pérez-Escamilla R, Curry L, Minhas D, et al. Scaling up of breastfeeding promotion programs in low- and middle-income countries: the “breastfeeding gear” model. *Adv Nutr Bethesda Md* 2012;3(6):790–800; doi: 10.3945/an.112.002873.
22. Raju TNK. Achieving healthy people 2030 breastfeeding targets in the United States: challenges and opportunities. *J Perinatol* 2023;43(1):74–80; doi: 10.1038/s41372-022-01535-x.
23. Segura-Pérez S, Hromi-Fiedler A, Adnew M, et al. Impact of breastfeeding interventions among United States minority women on breastfeeding outcomes: a systematic review. *Int J Equity Health* 2021;20(1):72; doi: 10.1186/s12939-021-01388-4.
24. Arring NM, Aduse-Poku L, Jiagge E, et al. A scoping review of strategies to increase Black enrollment and retention in cancer clinical trials. *JCO Oncol Pr* 2022;18(9):614–632; doi: 10.1200/OP.21.00863.

25. Thornton RLJ, Glover CM, Cené CW, et al. Evaluating strategies for reducing health disparities by addressing the social determinants of health. *Health Aff (Millwood)* 2016;35(8):1416–1423; doi: 10.1377/hlthaff.2015.1357.
26. Horowitz CR, Sabin T, Ramos M, et al. Successful recruitment and retention of diverse participants in a genomics clinical trial: a good invitation to a great party. *Genet Med Off J Am Coll Med Genet* 2019;21(10):2364–2370; doi: 10.1038/s41436-019-0498-x.
27. Harris MT, Campbell Statler M, Montague E, et al. Identifying and reducing barriers to having a racially and ethnically diverse research population. *Int J Ind Ergon* 2024;99:103527; doi: 10.1016/j.ergon.2023.103527.
28. Taani MH, Zabler B, Fendrich M, et al. Lessons learned for recruitment and retention of low-income African Americans. *Contemp Clin Trials Commun* 2020;17:100533; doi: 10.1016/j.conctc.2020.100533.
29. O'Donoghue J, Luther J, Hoque S, et al. Strategies to improve the recruitment and retention of underserved children and families in clinical trials: a case example of a school-supervised asthma therapy pilot. *Contemp Clin Trials* 2022;120:106884; doi: 10.1016/j.cct.2022.106884.
30. Harari N, Rosenthal MS, Bozzi V, et al. Feasibility and acceptability of a text message intervention used as an adjunct tool by WIC breastfeeding peer counsellors: the LATCH pilot. n.d.; doi: 10.1111/mcn.12488.
31. Lee HJ, Whitacre B. Estimating willingness-to-pay for broadband attributes among low-income consumers. *Telecommun Policy* 2017.
32. FCC. Lifeline Program for Low-Income Consumers | Federal Communications Commission. 2024. Available from: <https://www.fcc.gov/general/lifeline-program-low-income-consumers> [Last accessed: 3/13/2024].
33. Smith A. U.S. Smartphone Use in 2015. 2015. Available from: <https://www.pewresearch.org/internet/2015/04/01/us-smartphone-use-in-2015/> [Last accessed: 3/13/2024].
34. Starkweather AR, Heineman A, Storey S, et al. Methods to measure peripheral and central sensitization using quantitative sensory testing: a focus on individuals with low back pain. *Appl Nurs Res ANR* 2016;29:237–241; doi: 10.1016/j.apnr.2015.03.013.
35. Rolke R, Magerl W, Campbell KA, et al. Quantitative sensory testing: a comprehensive protocol for clinical trials. *Eur J Pain* 2006;10(1):77–77; doi: 10.1016/j.ejpain.2005.02.003.
36. Pocock SJ, Simon R. Sequential treatment assignment with balancing for prognostic factors in the controlled clinical trial. *Biometrics* 1975;31(1):103–115.

37. Jin M, Polis A, Hartzel J. Algorithms for minimization randomization and the implementation with an R package. *Commun Stat - Simul Comput* 2021;50(10):3077–3087; doi: 10.1080/03610918.2019.1619765.
38. R Core Team. R: A language and environment for statistical computing. MSOR Connect 2014.
39. U.S. Census Bureau. QuickFacts: Connecticut; United States. 2022. Available from: <https://www.census.gov/quickfacts/fact/table/CT,US/PST045222> [Last accessed: 3/13/2024].
40. Lee E, Wang JW. *Statistical Methods for Survival Data Analysis*. 4th ed. Wiley; 2013.
41. Bender W, Levine L, Durnwald C. Text message-based breastfeeding support compared with usual care: a randomized controlled trial. *Obstet Gynecol* 2022;140(5):853–860; doi: 10.1097/AOG.0000000000004961.
42. Geller SE, Koch AR, Roesch P, et al. The more things change, the more they stay the same: a study to evaluate compliance with inclusion and assessment of women and minorities in randomized controlled trials. *Acad Med J Assoc Am Med Coll* 2018;93(4):630–635; doi: 10.1097/ACM.0000000000002027.
43. Burns E, Schmied V. “The right help at the right time”: positive constructions of peer and professional support for breastfeeding. *Women Birth J Aust Coll Midwives* 2017;30(5):389–397; doi: 10.1016/j.wombi.2017.03.002.
44. Demirci JR, Suffoletto B, Doman J, et al. The development and evaluation of a text message program to prevent perceived insufficient milk among first-time mothers: retrospective analysis of a randomized controlled trial. *JMIR MHealth UHealth* 2020;8(4):e17328; doi: 10.2196/17328.
45. Lucas R, Zhang Y, Walsh SJ, et al. Efficacy of a breastfeeding pain self-management intervention: a pilot randomized controlled trial. *Nurs Res* 2019;68(2):E1–E10; doi: 10.1097/NNR.0000000000000336.
46. Bartick M, Reyes C. Las dos cosas: an analysis of attitudes of Latina women on non-exclusive breastfeeding. *Breastfeed Med Off J Acad Breastfeed Med* 2012;7(1):19–24; doi: 10.1089/bfm.2011.0039.
47. Tran V, Masterson AR, Frieson T, et al. Barriers and facilitators to exclusive breastfeeding among Black mothers: a qualitative study utilizing a modified Barrier Analysis approach. *Matern Child Nutr* 2022;19(1):e13428; doi: 10.1111/mcn.13428.
48. Feltner C, Weber RP, Stuebe A, et al. *Breastfeeding Programs and Policies, Breastfeeding Uptake, and Maternal Health Outcomes in Developed Countries*. AHRQ Comparative Effectiveness Reviews. Agency for Healthcare Research and Quality (US): Rockville (MD); 2018.

49. Dreyfus B, Kuri L, Ferri M, et al. Understanding Hispanic/Latino participation in clinical trials and observational studies, and strategies to increase participation: a targeted literature review. *J Health Care Poor Underserved* 2023;34(1):399–424; doi: 10.1353/hpu.2023.0026.
50. Patchen L, Ellis L, Harrington CB, et al. Engaging African American parents to develop a mobile health technology for breastfeeding: KULEA-NET. *J Hum Lact* 2020;36(3):448–460; doi: 10.1177/0890334420930208.
51. Martinez-Brockman JL, Harari N, Segura-Pérez S, et al. Impact of the Lactation Advice Through Texting Can Help (LATCH) trial on time to first contact and exclusive breastfeeding among WIC participants. *J Nutr Educ Behav* 2018;50(1):33-42.e1; doi: 10.1016/j.jneb.2017.09.001.
52. Teague S, Youssef GJ, Macdonald JA, et al. Retention strategies in longitudinal cohort studies: a systematic review and meta-analysis. *BMC Med Res Methodol* 2018;18(1):151; doi: 10.1186/s12874-018-0586-7.

## RETENTION AND STRUCTURAL INEQUITIES

**Table 1**  
*Demographic Characteristics (n=250)*

	<i>n</i>	%
Age, mean (SD)	30.24 (4.91)	
Race/Ethnicity		
Non-Hispanic White	118	47.2
Non-Hispanic Black	43	17.2
Hispanic	71	28.4
Other	18	7.2
Income		
≤ \$50,000	89	35.6
> \$50,000	155	62.0
Missing	6	2.4
History of breastfeeding		
Yes	130	52.0
No	120	48.0
Insurance		
Medicaid	91	36.4
Private	139	55.6
Self-pay/uninsured_and_other	14	5.6
Missing	6	2.4
Education		
Any College or above	207	82.8
High School or below	43	17.2
Infant Gender		
Male	132	52.8
Female	111	44.4
Missing	7	2.8
Infant Weight (kg) mean (SD)	117.182 (28.872)	
Pre-pregnancy BMI mean (SD)	27.53 (6.45)	
Return to work		
≤6 weeks	9	3.6
>6 weeks	241	96.4
Route of delivery		
Vaginal	167	66.8
Cesarean section	76	30.4
Missing	7	2.8

**Table 2***Retention of Key Characteristics at Baseline, 1, 2, 3, 6, 9, 12, 18, and 24 Weeks*

Variable	Baseline	Week 1	Week 2	Week 3	Week 6	Week 9	Week 12	Week 18	Week 24	P value
	<i>n</i>	<i>n</i>	<i>n</i>	<i>n</i>	<i>n</i>	<i>n</i>	<i>n</i>	<i>n</i>	<i>n</i>	
Total Sample	244	225	211	205	200	187	184	179	175	
Income										0.0001
≤ \$50,000	89	77	73	70	66	56	56	52	50	
> \$50,000	155	148	138	135	134	131	128	127	125	
Race and Ethnicity										0.0001
Non-Hispanic Black	40	36	33	30	28	24	23	23	21	
Non-Hispanic White	117	111	107	106	103	102	101	98	96	
Hispanic	70	62	55	53	53	46	45	43	43	
Other	17	16	16	16	16	15	15	15	15	
Age mean ( <i>SD</i> )	30.29 (4.94)	30.4 (4.97)	30.5 (4.93)	30.58 (4.9)	30.66 (4.92)	30.81 (4.83)	30.84 (4.75)	30.92 (4.76)	30.95 (4.74)	0.0001

## RETENTION AND STRUCTURAL INEQUITIES

23

**Table 3**

*Multiple Cox regression model of characteristics of retention to 24 weeks*

Fixed Effects	Coefficient	Adjusted Hazard Ratio Model*				P
		Exp (Coef)	Exp (-Coef)	95% CI		
				LL	UL	
Income						0.140
≤ \$50,000	Reference	Reference	Reference			
>\$50,000	-0.445	0.641	0.301	-1.036	0.146	
Race						0.080
Non-Hispanic White and Other	Reference	Reference	Reference			
Hispanic	0.648	1.912	0.324	0.014	1.282	0.045
Non-Hispanic Black	0.753	2.123	0.375	0.017	1.489	0.045
Age	-0.029	0.971	0.028	0.083	0.025	0.292

\* $p=0.0001$

